APPLIED AND ENVIRONMENTAL MICROBIOLOGY, June 2001, p. 2421–2429 Vol. 67, No. 6
0099-2240/01/$04.00+0 DOI: 10.1128/AEM.67.6.2421–2429.2001
Copyright © 2001, American Society for Microbiology. All Rights Reserved.

Genotypes Associated with Virulence in Environmental Isolates of *Vibrio cholerae*

IRMA N. G. RIVERA,[1,2,3]* JONGSIK CHUN,[4] ANWAR HUQ,[1,5] R. BRAD SACK,[6] AND RITA R. COLWELL[1,5]

Center of Marine Biotechnology, University of Maryland Biotechnology Institute, Baltimore, Maryland 21202[1]; Department of Cell and Molecular Biology, University of Maryland, College Park, Maryland 20742[5]; Departamento de Microbiologia, Imunologia e Parasitologia, Escola Paulista de Medicina,[2] and Departamento de Microbiologia, Instituto de Ciências Biomédicas, Universidade de São Paulo, CEP 05508-900,[3] São Paulo, Brazil; Microbiology Department, Seoul National University, Seoul, Korea 151-742[4]; and School of Hygiene and Public Health, Johns Hopkins University, Baltimore, Maryland 21205[6]

Received 30 October 2000/Accepted 5 March 2001

***Vibrio cholerae* is an autochthonous inhabitant of riverine and estuarine environments and also is a facultative pathogen for humans. Genotyping can be useful in assessing the risk of contracting cholera, intestinal, or extraintestinal infections via drinking water and/or seafood. In this study, environmental isolates of *V. cholerae* were examined for the presence of *ctxA, hlyA, ompU, stn/sto, tcpA, tcpI, toxR*, and *zot* genes, using multiplex PCR. Based on *tcpA* and *hlyA* gene comparisons, the strains could be grouped into Classical and El Tor biotypes. The *toxR, hlyA*, and *ompU* genes were present in 100, 98.6, and 87.0% of the *V. cholerae* isolates, respectively. The CTX genetic element and toxin-coregulated pilus El Tor (*tcpA* ET) gene were present in all toxigenic *V. cholerae* O1 and *V. cholerae* O139 strains examined in this study. Three of four nontoxigenic *V. cholerae* O1 strains contained *tcpA* ET. Interestingly, among the isolates of *V. cholerae* non-O1/non-O139, two had *tcpA* Classical, nine contained *tcpA* El Tor, three showed homology with both biotype genes, and four carried the *ctxA* gene. The *stn/sto* genes were present in 28.2% of the non-O1/non-O139 strains, in 10.5% of the toxigenic *V. cholerae* O1, and in 14.3% of the O139 serogroups. Except for *stn/sto* genes, all of the other genes studied occurred with high frequency in toxigenic *V. cholerae* O1 and O139 strains. Based on results of this study, surveillance of non-O1/non-O139 *V. cholerae* in the aquatic environment, combined with genotype monitoring using *ctxA, stn/sto*, and *tcpA* ET genes, could be valuable in human health risk assessment.**

Toxigenic *Vibrio cholerae* O1 and *V. cholerae* O139 are etiological agents of epidemic cholera. However, both *V. cholerae* O1 strains that do not produce cholera toxin, i.e., that are nontoxigenic (NT), and non-O1/non-O139 strains have also been associated with cholera, gastroenteritis, septicemia, and/or extraintestinal infections (49, 51, 52, 63, 66, 71, 81). Outbreaks of cholera were reported in Brazil during the third (1853 to 1854), fourth (1866 to 1868), and fifth (1893 to 1895) pandemics (3). In the early 1970s, when cholera spread to Africa and Southern Europe, it was forecast to arrive in countries across the Atlantic as well. This prompted the establishment of a surveillance program in S. Paulo State, Brazil, by the WHO (World Health Organization) and CETESB (Companhia de Tecnologia de Saneamento Ambiental, S.P.-Brazil—State Agency for Environmental Control). Sewage samples were monitored for *V. cholerae* in the community, and 12,867 samples were collected. From these samples, four NT *V. cholerae* O1 strains were isolated, the first *V. cholerae* O1 NT strains to be isolated in Brazil. The isolates were from sewage samples collected in 1978, 1980, and 1983 (42). Non-O1 *V. cholerae* strains were subsequently isolated from sewage (77.3%), seawater (40.4%), and freshwater (33.3%) samples collected in 1982 and 1983, at a time when no cases of cholera or gastroenteritis had been reported in S. Paulo State (41). At the same time, in Rio de Janeiro State, non-O1 *V. cholerae* was isolated from 12% of seawater and oyster samples (64).

The seventh pandemic reached the Americas on 29 January 1991 in Lima, Peru, and spread rapidly to the Peruvian Northern Andean and Amazon regions (74). Brazil reported its first case of cholera on 8 April 1991 in Tocantins. Cholera cases then occurred in the Amapa, Amazonas, Maranhão, Mato Grosso, Pará, and Rondonia States. However, in S. Paulo State, a more developed region of the country, the highest incidence of cholera in 1994 was 0.23 per 100,000 inhabitants compared to 33.4, the national average that same year (47).

The pathogenicity of *V. cholerae* O1 and O139 strains depends on a combination of properties, including enterotoxin (cholera toxin [CT], *ctxA*) and the ability to adhere to, and colonize, the small intestine (colonization factor, *tcpA*) (27). The major virulence-associated factors are present in clusters (23), with at least three regions in the *V. cholerae* chromosome. The first is the CTX genetic element (45), which has now been reported to comprise the genome of a filamentous bacteriophage (CTXΦ) (80). The second region is a large pathogenicity island for *V. cholerae* (VPI) (35) that encodes a toxin-coregulated pilus (TCP) gene cluster, a type IV pilus that functions as an essential colonization factor (75) and acts as CTXΦ receptor (35). The third gene cluster, the RTX toxin gene cluster, was described by Lin et al. (40) in a *V. cholerae* O1 El Tor strain and encodes cytotoxic activity for Hep-2 cells in vitro. However, the implication of RTX in pathogenesis has yet to be confirmed (16). Other factors have been associated with enteropathogenicity and include an El Tor-like hemolysin (*hlyA*)

* Corresponding author. Mailing address: Departamento de Microbiologia, ICB-USP, 1374, Lineu Prestes Av.—Edif. ICB II, USP-S.P., São Paulo, Brazil CEP 05508-900. Phone: 55-11-38187205. Fax: 55-11-38187354. E-mail: igrivera@usp.br.

Downloaded from http://aem.asm.org/ on December 27, 2016 by University of North Texas Libraries

TABLE 1. Genotypic traits of toxigenic and NT *V. cholerae* O1, *V. cholerae* O139, and *V. mimicus* isolates examined in this study

Strain (no. of strains)	Country	Reference(s)	Yr	Source	Code (*n*)	Genotype (presence [+] or absence [−] of genes)							
						ctxA	*hlyA*ET	*ompU*	*stn/sto*	*tcpA*ET	*tcpI*	*toxR*	*zot*
Toxigenic *V. cholerae* O1 (19)	Brazil	61	1991–1994	Sewage	RC7, RC221, RC226, CT2578, CT8514, CT31644, CT25017, CT24082 (8)	+	+	+	−	+	+	+	+
	Peru	61	1991, 1993	Clinical	RC8, IAL 1941 (2)	+	+	+	−	+	+	+	+
	Brazil	61	1993	Sewage	RC243, CT15861 (2)	+	+	+	−	+	−	+	+
	Chile		1991	Clinical	RC11	+	+	+	−	+	+	+	+
	Mexico		1991	Clinical	RC25	+	+	+	−	+	+	+	+
	Brazil	41	1994	River	CT7021	+	+	+	−	+	+	+	+
	Brazil	61	1992	Clinical	CTMACM14	+	+	+	−	+	−	+	+
	Brazil	61	1993	Sewage	RC224	+	+	+	+	+	+	+	+
	Peru		1991	Clinical	RC24	+	+	+	+	+	+	+	+
	Brazil	61	1993	Sewage	CT15989	+	+	+	*[b]	+	+	+	+
Total positives						19	19	19	2	19	16	19	19
NT *V. cholerae* O1 (4)	Brazil	41	1978	Sewage	TM45	−	+	+	−	+	+	+	−
	Brazil	41, 61	1978	Sewage	RC229, TM207831 (2)	−	+	+	−	+	−	+	−
	Brazil	41	1980	Sewage	RC231	−	+	+	−	−	+	+	−
Total positives						0	4	4	0	3	2	4	0
V. cholerae O139 (7)	India	61	1993	Clinical	RC4, RC30, RC120, RC138, RC139 (5)	+	+	+	−	+	+	+	+
	Bangladesh		1993	Clinical	RC46	+	+	+	+	+	−	+	+
	India	61	1993	Clinical	IGV	+	+	+	−	+	−	+	+
Total positives						7	7	7	1	7	5	7	7
V. mimicus (5)	Louisiana			Environ[a]	RC54	−	−	+	+	−	−	−	−
	Louisiana			Environ[a]	RC55	+	+	−	+	−	+	−	+
	Louisiana			Environ[a]	RC56	+	+	−	−	+	+	−	+
	Louisiana			Environ[a]	RC57	+	+	−	+	−	−	−	−
	Louisiana			Environ[a]	RC59	+	−	+	−	−	−	−	+
Total positives						4	3	2	3	1	2	0	3

[a] Environ, environmental.
[b] *, Amplicon with a different size.

(82), heat-stable enterotoxin (*stn/sto*) (1, 22, 55), hemagglutinins (14), neuraminidase (*nanH*) (20), a new CT (33), outer membrane protein (*ompU*) (72), Shiga-like toxin (*stx*) (33), a ToxR regulatory protein (46), and a zonula occludens toxin (*zot*) (18).

V. cholerae can be found in the environment both as a free-living bacterium and in association with zooplankton (30). Therefore, not surprisingly, non-O1/non-O139 *V. cholerae* is frequently isolated from the aquatic environment and seafood (5, 8, 9, 29, 30, 31, 32, 41, 43, 44, 78). In fact, *V. cholerae* is a heterogeneous species, with 206 serotypes identified to date (G. B. Nair, personal communication).

The emergence of the new serogroup O139 as a second etiologic agent of cholera epidemics (48), along with the discovery of horizontal and vertical genetic transfer by phages (80) and the elucidation of pathogenicity islands and other mobile genetic elements (36), has revived interest in the non-O1/non-O139 *V. cholerae* strains that had been previously implicated in cholera-like epidemics (2, 13, 51, 71, 81). In addition, the possible conversion of non-O1 to O1 serotype has provided added interest (10).

Cholera surveillance is now under way in many countries, based primarily on detection of *V. cholerae* O1 and O139 and determining the presence of cholera toxin using biological and molecular methods. However, virulence-associated factors in *V. cholerae* isolates from environmental sources are of concern.

The primary objective of this study was to evaluate the presence of virulence-associated factors in *V. cholerae* populations as potential pathogenic markers suitable for environmental monitoring. Virulence-associated factors studied here included cholera toxin (*ctxA*), hemolysin (*hlyA*), non-O1 heat-stable enterotoxin (*stn/sto*), outer membrane protein (*ompU*), TCP (*tcpA* and *tcpI*), ToxR regulatory protein (*toxR*), and zonula occludens toxin (*zot*).

MATERIALS AND METHODS

Bacterial strains. A total of 69 *V. cholerae* isolates were included in this study. *V. cholerae* O1 strains comprised 19 toxigenic clinical and environmental isolates from Brazil (14 isolates), Peru (3 isolates), Chile (1 isolate), and Mexico (1 isolate) and four nontoxigenic *V. cholerae* O1 isolates from Brazil (1978 to 1980). Thirty-nine environmental isolates of non-O1/non-O139 *V. cholerae* from Brazil, seven *V. cholerae* O139 clinical isolates from India, and five *V. mimicus* environmental isolates from Louisiana (United States) were included in the study (Tables 1 and 2). The Brazilian environmental strains were isolated in the CETESB Laboratory, S. Paulo, Brazil. All of the isolates are part of a culture collection (RRC) at the Center of Marine Biotechnology (Baltimore, Md.). Frozen stock cultures were subcultured on Luria-Bertani (LB) broth (Difco Laboratories, Detroit, Mich.), streaked onto LB agar, and then onto TCBS agar (Oxoid) to verify purity.

Downloaded from http://aem.asm.org/ on December 27, 2016 by University of North Texas Libraries

TABLE 2. Genotypic traits of non-O1/non-O139 *V. cholerae* Brazilian isolates examined in this study

Country (no.)	Yr	Reference(s)	Source	Code (*n*)	Genotype (presence [+] or absence [−] of genes)									
					ctxA	*hlyA*ET	*hlyA*Class	*ompU*	*stn*/*sto*	*tcpA*ET	*tcpA*Class	*tcpI*	*toxR*	*zot*
Brazil (9)	1992	43	Mussels	RC236, RC237, RC238, RC244 (4)	−	+	−	+	−	−	−	+	+	−
	1992	43, 61	Mussels	RC245, GM39 (2)	−	+	−	+	−	−	−	−	+	−
	1992	43, 61	Mussels	RC239, GM32 (2)	−	+	−	+	−	+	−	+	+	−
	1992	44	Oysters	RC247	−	−	−	+	−	−	−	−	+	−
Brazil (9)	1981, 1982	—	Seawater	RC63, RC64, RC65 (3)	−	+	−	+	+	−	−	+	+	−
	1982	—	Seawater	RC62	−	+	−	+	+	−	+	−	+	−
	1983	61	Seawater	TM16589	−	+	−	+	−	−	−	+	+	−
	1982	—	Seawater	RC61	+	+	−	+	+	−	+	+	+	−
	1981	41	Seawater	RC253	+	+	−	+	−	+	−	−	+	−
	1982	—	Seawater	TMA52	−	+	−	+	+	+	+	+	+	−
	1982	—	Seawater	RC60	+	−	+	+	+	+	+	+	+	−
Brazil (7)	1982	—	Sediment	RC68	−	+	−	+	*[a]	−	−	−	+	−
	1981	61	Sediment	TM50022	−	+	−	+	−	+	−	−	+	−
	1983	—	Sediment	RC71	−	+	−	+	−	+	−	+	+	−
	1983	—	Sediment	RC70	−	+	−	+	+	−	−	−	+	−
	1982	—	Sediment	RC66	−	+	−	+	+	+	−	−	+	−
	1982	—	Sediment	RC67	−	+	−	+	+	−	−	+	+	−
	1983	—	Sediment	RC69	−	+	−	+	+	+	+	+	+	−
Brazil (14)	1977, 1978, 1979	41, 61	Wastewater	RC235, RC248, TM41338, IG4 (4)	−	+	−	−	−	−	−	−	+	−
	1977, 1978	41	Wastewater	RC242, RC251 (2)	−	+	−	*[a]	−	−	−	−	+	−
	1979, 1992	41	Wastewater	RC234, RC252 (2)	−	+	−	+	−	−	−	+	+	−
	1992	41	River	RC241	−	+	−	−	−	−	−	+	+	−
	1983	41	Wastewater	RC240	−	+	−	−	−	+	−	+	+	−
	1979	61	Wastewater	TM31152	−	+	−	*[a]	−	−	−	+	+	−
	1992	61	Wastewater	CT3481	−	+	−	+	−	−	−	−	+	−
	1977	41	Wastewater	RC233	+	+	−	+	−	+	−	−	+	−
	1982	41	Wastewater	RC246	−	+	−	+	−	+	−	+	+	−
Total strains (39)					4	37	1	30	11	12	5	22	39	0

[a] *, Amplicon with different size.

Positive and negative controls. *V. cholerae* O1 Classical ATCC 11623, *V. cholerae* O1 El Tor ATCC 14033 (*ctxAB* negative), *V. cholerae* O1 Classical ATCC 14035, *V. cholerae* non-O1 ATCC 14547, *V. cholerae* non-O1 ATCC 25872 (*ctxAB*$^+$), *V. cholerae* non-O1 ATCC 25874 (*ctxAB*$^+$), *V. mimicus* ATCC 33653, *V. cholerae* O22, and *V. cholerae* O31 were used as positive controls, and *Escherichia coli* was used as a negative control.

Chromosomal DNA preparation. DNA was extracted by the CTAB (cetyltrimethylammonium bromide) method previously described (62). DNA extracts were resuspended in Tris-EDTA (10 mM Tris-HCl, 0.10 mM EDTA [pH 8.0]) buffer and stored at 4°C for further analysis. Dilutions of template DNA were made with sterile distilled water to obtain a concentration of ca. 100 ng/μl.

PCR primers and amplification conditions. The oligonucleotide primers for each of the selected virulence-associated factors were designed based on available GenBank sequences for *V. cholerae* O1 Classical and *V. cholerae* O1 E1 Tor for all genes, except the *stn*/*sto* genes, for which *V. cholerae* non-O1 and *V. cholerae* O1 sequences were used. The sequence positions and accession numbers of the sequences or sources are listed in Table 3. Oligonucleotide primers were synthesized by Genosys Biotechnologies, Inc.

Multiplex PCR. The following reagents were added to each sample PCR mixture: 2.5 μl of 10× amplification buffer A (100 mM Tris-HCl [pH 8.3], 500 mM KCl, 15 mM $MgCl_2$, 0.01% [wt/vol] gelatin) (Promega, Madison, Wis.); 0.5 μl each of 2.5 mM dATP, dCTP, dGTP, and dTTP (Promega); 1.0 μl each of forward and reverse primers (20 μM); 0.125 μl of *Taq* DNA polymerase at 5 U/μl (Promega); and Milli-Q water (to a final volume of 24 μl). PCR was carried out in 0.2-ml microcentrifuge tubes with 24 μl of the PCR mixture and 1 μl (ca. 0.10 μg) of template DNA. The solution was mixed, centrifuged briefly, and placed in an automated Peltier thermal cycler (PTC-200; M. J. Research).

PCR amplification conditions were as follows: denaturation at 94°C for 2 min, annealing for 1 min at the temperatures shown in Table 3, and extension at 72°C, as given in Table 3; with a final extension step at 72°C for 10 min at the end of 30 cycles, followed by maintenance at 4°C.

PCR products were separated by agarose gel electrophoresis (1.4%) in 1× TAE buffer (0.04 Tris-acetate, 0.001 M EDTA [pH 8.0]), stained with ethidium bromide, and visualized by using UV light.

RESULTS

Multiplex PCR was carried out, using positive and negative controls and the primers designed for the genes studied. The size of each amplicon was verified. Optimal PCR conditions were determined for *ctxA*/*ompU*-PCR, *hlyA* (Classical and El Tor), *zot*/*toxR*-PCR, and *tcpA* (Classical and El Tor), using multiplex PCR. For *tcpI* and *stn*/*sto* genes, simple PCR was used. The multiplex and simple PCR products obtained for each gene investigated are shown in Fig. 1, and the corresponding amplicon sizes are given in Table 3.

The virulence-associated factors for specific *V. cholerae* serogroups are summarized in Tables 1 and 2.

All of the *V. cholerae* O1, O139, and non-O1/non-O139 strains, regardless of whether they were toxigenic or NT, were found to possess the *tox*R regulatory sequence, a gene absent in the five *V. mimicus* strains tested.

CT and zonula occludens toxin (ZOT) were present in all (100%) of the toxigenic *V. cholerae* O1 and O139 strains tested and were absent in NT *V. cholerae* O1. In *V. cholerae* non-O1/non-O139 strains, we found four strains (RC60, RC61, RC233, and RC253) to be *ctxA* positive and *zot* negative. These isolates

Downloaded from http://aem.asm.org/ on December 27, 2016 by University of North Texas Libraries

TABLE 3. Primers used in this study

Gene(s), primers, and sequences (5′ to 3′)	Amplicon size (bp)	PCR conditions[a]	Source or reference; accession no.
ctxA and *ompU* (CT subunit A and outer membrane protein)		60–1	
94F, CGG GCA GAT TCT AGA CCT CCT G	564 (*ctxA*)		19
614R, CGA TGA TCT TGG AGC ATT CCC AC			
80F, ACG CTG ACG GAA TCA ACC AAA G	869 (*ompU*)		This study; AE004149, AF253529, U73751
906R, GCG GAA GTT TGG CTT GAA GTA G			
zot and *toxR* (zonula occludens toxin and operon ToxR)		60–1	
225F, TCG CTT AAC GAT GGC GCG TTT T	947 (*zot*)		This study; AE004224, AF123049, AF175708, M8363
1129R, AAC CCC GTT TCA CTT CTA CCC A			
101F, CCT TCG ATC CCC TAA GCA ATA C	779 (*toxR*)		This study; M21249, AE004179
837R, AGG GTT AGC AAC GAT GCG TAA G			
tcpA (TCP A [Classical and El Tor])		60–1	
72F, CAC GAT AAG AAA ACC GGT CAA GAG	451 (El Tor)		Modified from reference 39; AE004168, UO9807, X64098, X09807
477R, CGA AAG CAC CTT CTT TCA CGT TG			
647R, TTA CCA AAT GCA ACG CCG AAT G	620 (Classical)		
tcpI (TCP I)		60–3	
132F, TAG CCT TAG TTC TCA GCA GGC A	862 (*tcpI*)		This study; AE004168, L25659, X64098
951R, GGC AAT AGT GTC GAG CTC GTT A			
hlyA (hemolysin [Classical and El Tor])		60–1	
489F, GGC AAA CAG CGA AAC AAA TAC C	481 (El Tor)		Modified from reference 69; AE004362, AF117834, M36855, X51746, Y00557
744F, GAG CCG GCA TTC ATC TGA AT	738/727 (ET/Clas)		
1184R, CTC AGC GGG CTA ATA CGG TTT A			
stn/*sto* (non-O1 heat-stable enterotoxin)		55–1	
67F, TCG CAT TTA GCC AAA CAG TAG AAA	172 (*stn*/*sto*)		This study; M85198, X74108
194R, GCT GGA TTG CAA CAT ATT TCG C			

[a] Temperature of annealing (°C)–time of extension (min).

were positive for *ctxA*, using multiplex PCR assay, with a 564-bp amplicon identical to that of the O1 strains. The size of *ctxA* amplicons was confirmed by sequencing (data not shown).

The oligonucleotide primers targeting *tcpA* exploited sequence differences between the *tcpA* of the El Tor (ET) and Classical *V. cholerae* biotypes. All toxigenic *V. cholerae* O1 and O139 and three of four nontoxigenic *V. cholerae* O1 isolates examined showed amplicons of the same size as that obtained for the *tcpA* El Tor gene. Of 39 non-O1/non-O139 *V. cholerae* strains, 14 yielded amplicons, using *tcpA* primers. The amplicon size (451 bp) of nine strains (RC66, RC71, RC233, RC239, RC240, RC246, RC253, GM32, and TM50022) was identical to that obtained for *V. cholerae* El Tor ATCC 14033, while the amplicon size (620 bp) of two strains (RC61 and RC62) was identical to that obtained for *V. cholerae* O1 Classical ATCC 14035. Interestingly, three strains carried both genes (RC60, RC69, and TMA52) (Table 2).

The *tcpI* gene was frequently found in toxigenic *V. cholerae* O1 (84.2%) and O139 (71.4%) serogroups. However, this gene was also present in 50% of the NT *V. cholerae* O1 and in 56.4% of the non-O1/non-O139 serogroups.

The *hlyA* El Tor gene was found in all toxigenic and NT *V. cholerae* O1 and in *V. cholerae* O139. Among non-O1/non-O139 serogroup strains, 94.9% showed homology to El Tor hemolysin, 2.6% were associated with Classical hemolysin, and 2.5% were negative for both genes. The amplified fragment sizes were 727 bp, specifically for the Classical biotype (ATCC 11623 and ATCC 14035) and both 481 bp and 738 bp for the El Tor biotype. Occasionally, a larger fragment (~1.4 kb) was observed in the El Tor biotype strains.

The *ompU* gene was found in all strains of toxigenic and NT *V. cholerae* O1, *V. cholerae* O139, and in 76.9% of the environmental non-O1/non-O139 *V. cholerae* isolates.

Genes homologous to *stn*/*sto* were observed in toxigenic *V. cholerae* O1 (2 of 19), *V. cholerae* O139 (1 of 7), and non-O1/non-O139 *V. cholerae* (11 of 39) strains, yielding an amplicon size of 172 bp. However, we observed an amplicon of 800 bp in one isolate from sediment (RC68).

The genotypes found in each *V. cholerae* serogroup are listed in Table 4. The genotype most frequently observed in clinical toxigenic *V. cholerae* O1 (68.4%) and O139 (71.4%) isolates was *ctxA hlyA*ET *ompU tcpA*ET *tcpI toxR zot*. Other genotypes found in toxigenic *V. cholerae* O1 strains were similar, but the *tcpI* and/or *stn*/*sto* genes were absent. In non-O1/non-O139 *V. cholerae* serogroups, the genotypes were diverse; however, the most frequent genotypes were *hlyA*ET *ompU tcpI toxR* (7 of 39) and *hlyA toxR* (6 of 39).

The presence of virulence-associated genes in non-O1/non-O139 *V. cholerae* isolates was analyzed by origin, i.e., water and sediment for marine ecosystems, mussels, and wastewater, and the results are summarized in Table 5.

DISCUSSION

The majority of *V. cholerae* strains in the environment are considered to be harmless estuarine microorganisms. However, specific strains appear to have evolved that cause disease in humans by effectively colonizing the small intestine and releasing a potent enterotoxin. Eight chromosomal regions coding for virulence-associated factors in *V. cholerae* (*ctxA,*

Downloaded from http://aem.asm.org/ on December 27, 2016 by University of North Texas Libraries



FIG. 1. Agarose gel electrophoresis of amplicons obtained using multiplex PCR. Lane M, molecular weight marker (lambda DNA *Hin*dIII); lanes 1, 3, 5, 11, and 14, *V. cholerae* O1 El Tor nontoxigenic ATCC 14033; lanes 2, 6, 9, 12, and 15, *V. cholerae* O1 Classical ATCC 14035; lane 4, RC224, *V. cholerae* O1 El Tor (Brazilian isolate from sewage, 1993); lane 7, TMA52, *V. cholerae* non-O1/non-O139 (Brazilian isolate from seawater, 1982); lane 8, negative control; lane 10, RC46, *V. cholerae* O139 (*tcpI* negative); lane 13, RC60, *V. cholerae* non-O1/non-O139 (Brazilian isolate from seawater, 1982); lane 16, RC66 (Brazilian isolate from sediment, 1982).

hlyA, ompU, stn/sto, tcpA, tcpI, toxR, and *zot*) were included in this study.

Distribution of virulence-associated genes in *V. cholerae*. The major CTX genetic element has the structure of a compound transposon, with a 4.5-kb central core region (*ctxAB, zot, ace, orfU*, and *cep*) that encodes for both CT and for functions required for virion morphogenesis and is flanked by one or more copies of a 2.7-kb repetitive sequence that encodes functions required for regulation, replication, and integration of CTXϕ (36, 80). In this region, we tested for the presence of two genes: CT subunit A (*ctxA*) and ZOT (*zot*). Interestingly, we found four of the non-O1/non-O139 *V. cholerae* isolates contained *ctxA* gene but not the *zot* gene. The ZOT was described by Fasano et al. (18) as a toxin that increases permeability of the small intestinal mucosa by affecting the structure of the intercellular tight junction, or zonula occludens. However, Waldor and Mekalanos (80) found that *zot* and *orfU* correspond to genes involved in CTXΦ morphogenesis and that the biological activity previously designated "zonula occludens toxin" is probably not directly associated with the *zot* gene product, unless its product possesses a dual function. Additionally, the presence of the *zot* gene and the absence of the *ctx* gene in *V. cholerae* and *V. mimicus* strains has been reported (7, 21, 38). To explore the CTXΦ genome integrated in the *V. cholerae* chromosome, sequencing of the CTX genetic element is under way.

The *V. cholerae* pathogenicity island (VPI) is 39.5 kb and contains genes associated with virulence (TCP-ACF cluster), the regulation of virulence (*toxT* and *tcpP/H*), the regulation of chemotaxis (*tcpI* or *acfB*), and mobility (*int* and *orfI*) (24, 36). Because the genes encoding TCP have been suggested to be part of a larger genetic element consisting of a cluster of genes, we looked for the presence of both *tcpA* and *tcpI* genes. The *tcpA*ET gene was present in all toxigenic *V. cholerae* O1 and O139 isolates, showing that despite different lipopolysaccharide structures, the two share common TCP-associated antigens (60). The *tcpA*ET gene was also found in 3 of 4 strains of NT *V. cholerae* O1 and in 9 of 39 non-O1/non-O139 Brazilian strains. Interestingly, in the non-O1/non-O139 serogroup, two strains had *tcpA* genes of the Classical biotype and three had *tcpA* gene of both biotypes, in agreement with previous reports (6, 54). These results suggest that these strains may have a selective advantage over nonpathogenic strains, with an ability to colonize the human intestine, and become toxigenic (35). However, while the El Tor TcpA pilin is 82% identical to TcpA of the Classical biotype (60), there are at least four major variants of *tcpA* genes that probably evolved in parallel, though independently, from a common ancestral gene (54). Sequencing of *tcpA* El Tor amplicons demonstrated the fragment size to be 451 bp, and the amplicon sizes for *tcpA* Classical was 620 bp in the *V. cholerae* non-O1/non-O139 strains (data not shown). These variations could indicate other functional significance as well, because TcpA also acts as a coat protein of the bacteriophage VPΦ, produced by vibrios containing VPI (37, 77).

In this study, the presence of strains containing *tcpA* genes, either identical to those of biotype El Tor or of Classical, is in agreement with results obtained in Australia, where *V. cholerae* serogroup O6 was shown to contain the *tcpA*ET gene and another strain of serogroup O23 presented as *tcpA* Classical, both strains having been isolated from water (67).

TcpI, an integral inner membrane protein, involved in environmental sensing and signal transduction (chemosensors), negatively regulates the synthesis of the major pilin subunit of TCP (TcpA) (24, 76). Harkey et al. (24) suggested that regulators such as TcpI, that act downstream of ToxR and ToxT may function to fine-tune the expression of the TCP virulence determinant throughout the pathogenic cycle of *V. cholerae*. However, our results showing the presence of *tcpI* gene in 65.5% of *V. cholerae* strains suggests that the importance of this gene may be physiological and not pathogenesis alone.

Outer membrane protein, OmpU, was reported to be a potential adherence factor for *V. cholerae* (72). However, later studies suggested that OmpU is not involved in the adhesion of *V. cholerae* to the intestinal epithelium (53). In this study, the gene was present in 87% of the *V. cholerae* strains tested, except in nine non-O1/non-O139 *V. cholerae* strains isolated from wastewater. These findings suggest that this gene may be mainly physiological in its activity.

The hemolysin traditionally has been employed to differen-

Downloaded from http://aem.asm.org/ on December 27, 2016 by University of North Texas Libraries

TABLE 4. Genotypes of *V. cholerae* isolates listed by serogroup

Genotypes	No. of isolates				
	O1	NT O1	O139	Non-O1/ non-O139	Total
ctxA$^+$ *hlyA*$^+$ET *ompU*$^+$ *tcpA*$^+$ET *tcpI*$^+$ *toxR*$^+$ *zot*$^+$	13	0	5	0	18
ctxA$^+$ *hlyA*$^+$ET *ompU*$^+$ *tcpA*$^+$ET *toxR*$^+$ *zot*$^+$	3	0	1	0	4
ctxA$^+$ *hlyA*$^+$ET *ompU*$^+$ *stn/sto*$^+$ *tcpA*$^+$ET *tcpI*$^+$ *toxR*$^+$ *zot*$^+$	2+1[a]	0	0	0	3
ctxA$^+$ *hlyA*$^+$ET *ompU*$^+$ *tcpA*$^+$ET *toxR*$^+$	0	0	0	2	2
ctxA$^+$ *hlyA*$^+$ET *ompU*$^+$ *stn/sto*$^+$ *tcpA*$^+$ET *toxR*$^+$ *zot*$^+$	0	0	1	0	1
ctxA$^+$ *hlyA*$^+$ET *ompU*$^+$ *stn/sto*$^+$ *tcpA*$^+$Class *tcpI*$^+$ *toxR*$^+$	0	0	0	1	1
ctxA$^+$ *hlyA*$^+$Class *ompU*$^+$ *stn/sto*$^+$ *tcpA*$^+$ET/Class *tcpI*$^+$ *toxR*$^+$	0	0	0	1	1
hlyA$^+$ET *ompU*$^+$ *tcpI*$^+$ *toxR*$^+$	0	1	0	7	8
hlyA$^+$ET *toxR*$^+$	0	0	0	6	6
hlyA$^+$ET *ompU*$^+$ *tcpA*$^+$ET *tcpI*$^+$ *toxR*$^+$	0	1	0	4	5
hlyA$^+$ET *ompU*$^+$ *stn/sto*$^+$ *tcpI*$^+$ *toxR*$^+$	0	0	0	4	4
hlyA$^+$ET *ompU*$^+$ *toxR*$^+$	0	0	0	4	4
hlyA$^+$ET *ompU*$^+$ *tcpA*$^+$ET *toxR*$^+$	0	2	0	1	3
hlyA$^+$ET *ompU*$^+$ *stn/sto*$^+$ *tcpA*$^+$ET/Class *tcpI*$^+$ *toxR*$^+$	0	0	0	2	2
hlyA$^+$ET *tcpI*$^+$ *toxR*$^+$	0	0	0	2	2
hlyA$^+$ET *ompU*$^+$ *stn/sto*$^+$ *tcpA*$^+$Class *toxR*$^+$	0	0	0	1	1
hlyA$^+$ET *ompU*$^+$ *stn/sto*$^+$ *tcpA*$^+$ET *toxR*$^+$	0	0	0	1	1
hlyA$^+$ET *tcpA*$^+$ET *tcpI*$^+$ *toxR*$^+$	0	0	0	1	1

[a] 1, i.e., an *stn/sto* amplicon with another size.

tiate between the two biotypes of *V. cholerae* O1. The sequence of the Classical biotype has an 11-bp deletion within the *hlyA* structural gene, compared to the El Tor biotype (57). Using information on these genes, we designed primers to differentiate both biotypes and verified the presence of this gene in 98.6% of the *V. cholerae* strains, in agreement with previous reports (4, 68, 71, 82, 83). The majority belonged to the El Tor biotype, regardless of the hemolytic phenotype (67 of 69). The multiplex PCR for hemolysin effectively distinguished the two biotypes in non-O1/non-O139 *V. cholerae* strains compared with *V. cholerae* O1 Classical ATCC 11623 and ATCC 14035 and *V. cholerae* O1 El Tor ATCC 14033. Interestingly, the *hlyA* gene is located on chromosome 2 (26).

V. cholerae non-O1/non-O139 strains may also produce a 17-amino-acid heat-stable enterotoxin (NAG-ST) (*stn*) that is closely related to the heat-stable toxins produced by enterotoxigenic *E. coli* and other enteric pathogens (1, 55). Also, a heat-stable enterotoxin in *V. cholerae* O1 strains (*sto*) has been described (22). We designed primers based on the *stn/sto* genes associated with heat-stable enterotoxin production and found the genes to be homologous to those of toxigenic *V. cholerae* O1 (2 of 19), *V. cholerae* O139 (1 of 7), and non-O1/non-O139 *V. cholerae* (11 of 39) strains. This is the first report showing higher values than those reported in other studies, using hybridization with a NAG-ST probe, in *V. cholerae* non-O1/non-O139 population (12, 28, 56, 58). Interestingly, the *stn/sto* gene occurred more frequently in isolates from seawater (7 of 9) and sediment (4 of 7) and was absent in isolates from sewage (0 of 14) or oysters and mussels (0 of 9). It should be mentioned, however, that Caldini et al. (5) reported the presence of the *sto* gene in 12.7% (19 of 150) of the *V. cholerae* non-O1 isolates from freshwater in the river basin of central Italy, but the incidence of NAG-ST in *V. cholerae* non-O1 was not clearly established. Results of a study involving human volunteers demonstrated that, besides the production of NAG-ST, the virulence of non-O1/non-O139 *V. cholerae* depends on its ability to colonize the intestine (50).

The regulation and expression of genes for growth and survival depend on the regulon ToxR, coordinately regulated by a cascade mechanism involving three known components: ToxR, ToxS, and ToxT (15). ToxR, a 32-kDa transmembrane protein, is the master regulator, and its expression is dependent upon environmental growth conditions (incubation temperature, pH, osmolarity, bile salts, oxygen tension, hydrostatic pressure, and amino acid composition of the medium) (15, 46). The *toxR* gene encodes a transcriptional activator controlling CT gene

TABLE 5. Distribution of virulence-associated genes among non-O1/non-O139 *V. cholerae* isolates listed by source

Brazilian source	*n*	Virulence-associated gene-positive isolates[a]										
		ctxA	*hlyA*		*stn/sto*	*ompU*	*tcpA*			*tcpI*	*toxR*	*zot*
			ET	C			ET	C	ETC			
Mussels	9	0	8	0	0	9	2	0	0	6	9	0
Seawater	9	3	8	1	7	9	1	2	2	7	9	0
Sediment	7	0	7	0	4	7	3	0	1	3	7	0
Wastewater	14	1	14	0	0	5	3	0	0	6	14	0
Total	39	4	37	1	11	30	9	2	3	22	39	0

[a] C, Classical; ETC, both ET and Classical.

Downloaded from http://aem.asm.org/ on December 27, 2016 by University of North Texas Libraries

expression (*ctxA*), TCP biogenesis (*tcpA*), outer membrane protein expression (*ompU*), and at least 17 distinct genes in O139 and O1 strains (15, 27, 46). In this study, the presence of the *toxR* gene was verified in all *V. cholerae* studied, regardless of serogroup or source of isolation, a finding in agreement with previous reports (21, 71).

The presence of *toxR* (100%), *hlyA* (98.6%), *ompU* (87%), and *tcpI* (65.5%) genes in the *V. cholerae* isolates suggest that they are required for functioning of the organism in the environment and are not solely related to pathogenesis.

Five *V. mimicus* environmental strains were included in our study because of their close relationship to *V. cholerae*. None carried the *toxR* gene. The presence of CT in four of the five *V. mimicus* strains was confirmed, while the *zot* gene was found in only three of the strains, results similar to those reported earlier by Chowdhury et al. (7). The *stn*/*sto* gene was found in three of five environmental strains, as reported elsewhere, using PCR (79). In contrast, Pal et al. (56) reported the presence of NAG-ST gene in 13.7 and 22.6% of *V. mimicus* isolates from environmental and clinical sources, respectively, and Ramamurthy et al. (59) suggested that *V. mimicus* may act as a genetic reservoir for these genes.

Multiplex PCR was found to be sensitive and specific for assessing the pathogenicity of clinical and environmental *V. cholerae* isolates. We tested multiplex PCR for *ctxA*/*tcpA* El Tor genes using four primers—94F, 614R, 72F, and 477R—as a primary screening for *V. cholerae* during epidemiological surveillance with good success (data not shown).

***V. cholerae* genotypes.** A single factor cannot explain enteropathogenicity. In this study eight virulence-associated genes were detected in *V. cholerae* strains isolated from both clinical and environment sources. Previous findings, in an earlier study of 172 *V. cholerae* non-O1/non-O139 environmental isolates from seawater and sediment samples collected in São Paulo, Brazil, showed that 60.4% of the strains were *hlyAC*$^+$, *nanH*$^+$, and *toxR*$^+$ and *ctxAB*, *elt* (33), and *zot* negative as determined by using probes and radioactive hybridization. The frequency of occurrence of the genes among the strains tested was 98.8% for *toxR*, 97.1% for *hlyAC*, 66.9% for *nanH*, 5.8% for *slt*, 1.2% for *zot*, and 0.6% for *ctxAB* (D. E. Alvarado, V. H. Pellizari, T. A. T. Gomes, and I. N. G. Rivera, Abstr. 8th Int. Symp. Microb. Ecol. p. 89, 1998). In this study, we included *V. cholerae* O1 toxigenic and NT strains and *V. cholerae* O139 strains, using multiplex PCR, finding that 100% of the *V. cholerae* O1 and O139 strains carried *ctxA*, *hlyA*ET, *ompU*, *tcpA*ET, *toxR*, *zot*, and *tcpI* and sometimes the *stn*/*sto* gene.

Non-O1/non-O139 *V. cholerae* strains tested revealed the genotype *hlyA*ET$^+$ *ompU*$^+$ *tcpI*$^+$ *toxR*$^+$ and negative for *ctxA, stn*/*sto, tcpAET*, and *zot* for isolates from mussels (4 of 9), wastewater (2 of 14), and seawater (1 of 9). Interestingly, we observed the genotype *hlyA*ET$^+$ *toxR*$^+$ and negative for *ctxA*, *ompU*, *stn*/*sto*, *tcpA*ET, and *zot* in 6 of 14 non-O1/non-O139 *V. cholerae* isolates from wastewater, a finding similar to the virulence pattern obtained for 15 clinical strains associated with an unusual upsurge of cholera-like disease in India (71) and in non-O1/non-O139 *V. cholerae* strains isolated from volunteers in a vaccine trial in Peru (13).

A low frequency of the combination *ctxA*$^+$ *tcpA*Class$^+$ (1 of 39), and *ctxA*$^+$ *tcpA*ET$^+$ (3 of 39) was observed in environmental non-O1/non-O139 *V. cholerae* strains, as reported earlier (21). However, we found 23.1% (9 of 39) of CT-negative, but TcpAET-positive strains, potentially convertible to toxigenic strains by CTXΦ, either inside the host intestine or in the environment.

V. cholerae is autochthonous to estuarine and coastal environments (nutrient poor) and also colonize the human intestine (nutrient rich). In the study reported here, we observed a varied incidence of virulence-associated factors in *V. cholerae* isolates, depending on the source or ecosystem (seawater, seafood, wastewater, and clinical specimens). The response to changing conditions occurs by expressing appropriate sets of genes that promote growth in each niche.

Aquatic and marine ecosystems are subjected to large spatial and temporal nutrient fluxes arising from seasonal and geographic variations in temperature, salinity, nutrient input, pH, oxygen tension, etc. (65). The mechanisms by which environmental conditions affect the coordinated expression of virulence factors by *V. cholerae* remain poorly understood. Hase and Mekalanos (25) proposed a model where both ToxR and -S and TcpP and -H are involved in sensing various environmental and internal stimuli and are required for the production of TCP in *V. cholerae*. Furthermore, the *V. cholerae* genome encodes 43 methyl-accepting chemotaxis proteins that regulate swimming behavior in response to aminoacids, sugars, and oxygen (16).

Non-O1/non-O139 *V. cholerae* and risk assessment. Non-O1/non-O139 *V. cholerae* strains can no longer be ignored. The rationale for continuous monitoring is based first on the emergence of serogroup O139 (Bengal) in Bangladesh (CT positive) and Argentina (CT negative), each of which clearly evolved independently (73). Second, the sixth pandemic, the seventh pandemic, and U.S. Gulf Coast isolates represent three different clones, each independently evolved from environmental non-O1 *V. cholerae* isolates (34). Third, other epidemic serogroups have emerged, including *V. cholerae* O31 (50), O37 (11), O53 (52), O10, and O12 (13, 71). Finally, the emergence of a new clone of the *V. cholerae* O1 El Tor in Calcutta, India (70), and in Bangladesh (17) has been reported. The possibility exists that additional new strains of toxigenic *V. cholerae* with epidemic potential may emerge in the future.

We reported earlier an ERIC-PCR method for detecting emergent pathogenic *V. cholerae* strains, whereby specific fingerprints for pathogenic strains (CT and ZOT toxin positive) were different from nonpathogenic strains (61). In this study, we report probable parental toxigenic genotypes present in small numbers in the Brazilian environment. While these strains did not cause epidemics, there are environmental factors that may change, enhance multiplication or dominance, or select for genotypes of *V. cholerae*, and the strains themselves need to be tested for potential selective advantage under selected environmental conditions.

ACKNOWLEDGMENTS

Irma N. G. Rivera was supported by a Fellowship from UNESCO-ASM during the research period and from CNPQ (National Council for Research Support [Brazil]) during the postdoctoral period.This research was supported by NIH (grant 1R01A139129-01) and EPA (grant R824995-01).

We thank P. S. Sanchez and M. I. Sato from the State Agency for Environmental Control-CETESB, S. Paulo, Brazil, for providing the strains in this study.

Downloaded from http://aem.asm.org/ on December 27, 2016 by University of North Texas Libraries

REFERENCES

1. **Arita, M. T., T. Takeda, T. Honda, and T. Miwatani.** 1986. Purification and characterization of *Vibrio cholerae* non-O1 heat-stable enterotoxin. Infect. Immun. **52:**45–49.
2. **Bagchi, K., P. Echevarria, J. D. Arthur, O. Sethabutr, O. Serichantalergs, and C. W. Hoge.** 1993. Epidemic of diarrhea caused by *Vibrio cholerae* non-O1 that produced heat-stable toxin among Khmers in a camp in Thailand. J. Clin. Microbiol. **31:**1315–1317.
3. **Barua, D.** 1992. History of cholera, p. 1–35. *In* D. Barua and W. B. Greenough III (ed.), Cholera. Plenum Medical Book Company, New York, N.Y.
4. **Brown, M. H., and P. A. Manning.** 1985. Haemolysin genes of *Vibrio cholerae*: presence of homologous DNA in non-haemolytic O1 and haemolytic non-O1 strains. FEMS Microbiol. Lett. **30:**197–201.
5. **Caldini, G., A. Neri, S. Cresti, V. Boddi, G. M. Rossolini, and E. Lanciotti.** 1997. High prevalence of *Vibrio cholerae* non-O1 carrying heat-stable-enterotoxin-climate river basin of central Italy. Appl. Environ. Microbiol. **63:** 2934–2939.
6. **Chakraborty, S., A. K. Mukhopadhyay, R. K. Bhadra, A. N. Ghosh, R. Mitra, T. Shimada, S. Yamasaki, S. M. Faruque, Y. Takeda, R. R. Colwell, and G. B. Nair.** 2000. Virulence genes in environmental strains of *Vibrio cholerae*. Appl. Environ. Microbiol. **66:**4022–4028.
7. **Chowdhury, M. A. R., R. T. Hill, and R. R. Colwell.** 1994. A gene for the enterotoxin zonula occludens toxin is present in *V. mimicus* and *V. cholerae* O139. FEMS Microbiol. Lett. **119:**377–380.
8. **Colwell, R. R., J. B. Kaper, and S. W. Joseph.** 1977. *Vibrio cholerae* and *Vibrio parahaemolyticus* and other vibrios: occurrence and distribution in Chesapeake Bay. Science **198:**394–396.
9. **Colwell, R. R., and W. M. Spira.** 1992. The ecology of *Vibrio cholerae*, p. 107–127. *In* D. Barua and W. B. Greenough III (ed.), Cholera. Plenum Medical Book Company, New York, N.Y.
10. **Colwell, R. R., A. Huq, M. A. R. Chowdhury, P. R. Brayton, and B. Xu.** 1995. Serogroup conversion of *Vibrio cholerae*. Can. J. Microbiol. **41:**946–950.
11. **Dakin, S. P. H., D. J. Howel, R. G. A. Sutton, M. E. O'Keffe, and P. Thoms.** 1994. Gastroenteritis due to non-agglutinable (non-cholera) vibrios. Med. J. Aust. **2:**490.
12. **Dalsgaard, A., O. Serichantalergs, T. Shimada, O. Sethabutr, and P. Echevarria.** 1995. Prevalence of *Vibrio cholerae* with heat-stable enterotoxin (NAG-ST) and cholera toxin genes; restriction fragment length polymorphisms of NAG-ST genes among *V. cholerae* O serogroups from a major shrimp production area in Thailand. J. Med. Microbiol. **43:**216–220.
13. **Dalsgaard, A., M. J. Albert, D. N. Taylor, T. Shimada, R. Meza, O. Serichantalergs, and P. Echevarria.** 1995. Characterization of *Vibrio cholerae* non-O1 serogroups obtained from an outbreak of diarrhea in Lima, Peru. J. Clin. Microbiol. **33:**2715–2722.
14. **Datta-Roy, K., K. Banerjee, S. P. De, and A. C. Ghose.** 1986. Comparative study of expression of hemagglutinins, hemolysins, and enterotoxins by clinical and environmental isolates of non-O1 of *Vibrio cholerae* in relation to their enteropathogenicity. Appl. Environ. Microbiol. **52:**875–879.
15. **DiRita, V. J.** 1992. Co-ordinate control of virulence gene expression by ToxR in *Vibrio cholerae*. Mol. Microbiol. **6:**451–458.
16. **DiRita, V. J.** 2000. Genomics happens. Science **289:**1488.
17. **Faruque, S. M., M. J. Albert, and J. J. Mekalanos.** 1998. Epidemiology, genetics and ecology of toxigenic *Vibrio cholerae*. Microbiol. Mol. Biol. Rev. **62:**1301–1314.
18. **Fasano, A., B. Baudry, D. W. Pumplin, S. S. Wasserman, B. D. Tall, J. M. Ketley, and J. B. Kaper.** 1991. *Vibrio cholerae* produces a second enterotoxin which affects intestinal tight junctions. Proc. Nat. Acad. Sci. USA **88:**5242–5246.
19. **Fields, P. I., T. Popovic, K. Wachsmuth, and O. Olsvik.** 1992. Use of polymerase chain reaction for detection ot toxigenic *Vibrio cholerae* O1 strains from the Latin American cholera epidemics. J. Clin. Microbiol. **30:**2128–2121.
20. **Galen, J. E., J. M. Ketley, A. Fasano, S. H. Richardson, S. S. Wasserman, and J. B. Kaper.** 1992. Role of *Vibrio cholerae* neuraminidase in the function of cholera toxin. Infec. Immun. **60:**406–415.
21. **Ghosh, C., R. K. Nandy, S. K. Dasgupta, G. B. Nair, R. H. Hall, and A. C. Ghose.** 1997. A search for cholera toxin (CT), toxin coregulated pilus (TCP), the regulatory element ToxR and other virulence factors in non-O1/non-O139 *Vibrio cholerae*. Microb. Pathog. **22:**199–208.
22. **Guglielmetti, P., L. Bravo, A. Zanchi, R. Montè, G. Lombardi, and G. M. Rossolini.** 1994. Detection of the *Vibrio cholerae* heat-stable enterotoxin gene by polymerase chain reaction. Mol. Cell. Probes **8:**101–106.
23. **Hacker, J., G. Blum-Oehler, I. Mühldorfer, and H. Tschäpe.** 1997. Pathogenicity islands of virulent bacteria: structure, function and impact on microbial evolution. Mol. Microbiol. **23:**1089–1097.
24. **Harkey, C. W., K. D. Everiss, and K. M. Peterson.** 1994. The *Vibrio cholerae* toxin-coregulated-pilus gene *tcpI* encodes a homolog of methyl-accepting chemotaxis proteins. Infect. Immun. **62:**2669–2678.
25. **Häse, C. C., and J. J. Mekalanos.** 1998. TcpP protein is a positive regulator of virulence gene expression. Proc. Natl. Acad. Sci. USA **95:**730–734.
26. **Heidelberg, J. F., J. A. Eisen, W. C. Nelson, R. A. Clayton, M. L. Gwinn, R. J. Dodson, D. H. Haft, E. K. Hickey, J. D. Peterson, L. Umayam, S. R. Gill, K. E. Nelson, T. D. Read, H. Tettelin, D. Richardson, M. D. Ermolaeva, J. Vamathevan, S. Bass, H. Qin, I. Dragoi, P. Sellers, L. McDonald, T. Utterback, R. D. Fleishmann, W. C. Nierman, O. White, S. L. Salzberg, H. O. Smith, R. R. Colwell, J. J. Mekalanos, J. C. Venter, and C. M. Fraser.** 2000. DNA sequence of both chromosomes of the cholera pathogen *Vibrio cholerae*. Nature **406:**477–484.
27. **Herrington, D. A., R. H. Hall, G. Losonsky, J. J. Mekalanos, R. K. Taylor, and M. M. Levine.** 1988. Toxin, toxin-coregulated pili and the *toxR* regulon are essential for *Vibrio cholerae* pathogenesis in humans. J. Exp. Med. **168:** 1487–1492.
28. **Hoge, C. W., O. Sethabutr, L. Bodhidatta, P. Echevarria, D. C. Robertson, and J. G. Morris, Jr.** 1990. Use of a synthetic oligonucleotide probe to detect strains of non-serovar O1 *Vibrio cholerae* carrying the gene for heat-stable enterotoxin (NAG-ST). J. Clin. Microbiol. **28:**1473–1476.
29. **Huq, A., E. B. Small, P. A. West, M. I. Huq, R. Rahman, and R. R. Colwell.** 1983. Ecology of *V. cholerae* with special reference to planktonic crustacean copepods. Appl. Environ. Microbiol. **45:**275–283.
30. **Huq, A., and R. R. Colwell.** 1996. Vibrios in the marine and estuarine environment: tracking *Vibrio cholerae*. Ecosystem Health **2:**198–214.
31. **Islam, M. S., B. S. Drasar, and R. B. Sack.** 1994. The aquatic environment as reservoir of *Vibrio cholerae*: a review. J. Diarrhoeal Dis. Res. **11:**197–206.
32. **Kaper, J., H. Lockman, R. R. Colwell, and S. W. Joseph.** 1979. Ecology, serology and enterotoxin production of *Vibrio cholerae* in Chesapeake Bay. Appl. Environ. Microbiol. **37:**91–103.
33. **Kaper, J. B., A. Fasano, and M. Trucksis.** 1994. Toxins of *Vibrio cholerae*, p. 145–176. *In* I. K. Wachsmuth, P. A. Blake, and O. Olsvik (ed.), *Vibrio cholerae* and cholera: molecular to global perspectives. ASM Press, Washington, D.C.
34. **Karaolis, D. K. R., R. Lan, and P. R. Reeves.** 1995. The sixth and seventh cholera pandemic are due to independent clones separately derived from environmental, non-toxigenic, non-O1 *Vibrio cholerae*. J. Bacteriol. **177:** 3193–3198.
35. **Karaolis, D. K. R., J. A. Johnson, C. C. Bailey, E. C. Boedeker, J. B. Kaper, and P. R. Reeves.** 1998. A *Vibrio cholerae* pathogenicity island associated with epidemic and pandemic strains. Proc. Natl. Acad. Sci. USA **95:**3134–3139.
36. **Karaolis, D. K. R., and J. B. Kaper.** 1999. Pathogenicity islands and other mobile virulence elements of *Vibrio cholerae*, p. 167–187. *In* J. B. Kaper and J. Hacker (ed.), Pathogenicity islands and other mobile virulence elements. ASM Press, Washington, D.C.
37. **Karaolis, D. K. R., S. Somara, D. R. Maneval, Jr., J. A. Johnson, and J. B. Kaper.** 1999. A bacteriophage encoding a pathogenicity island, a type-IV pilus and a phage receptor in cholera bacteria. Nature **399:**375–379.
38. **Karasawa, T., T. Mihara, H. Kurazono, B. G. Nair, S. Garg, T. Ramamurthy, and Y. Takeda.** 1993. Distribution of the *zot* (zonula occludens toxin) gene among strains of *Vibrio cholerae* O1 and non-O1. FEMS Microbiol. Lett. **106:**143–146.
39. **Keasler, S. P., and R. H. Hall.** 1993. Detecting and biotyping *Vibrio cholerae* O1 with multiplex chain reaction. Lancet **341:**1661.
40. **Lin, W., K. J. Fullner, R. Clayton. J. A. Sexton, M. B. Rogers, K. E. Calia, S. B. Calderwood, C. Fraser, and J. J. Mekalanos.** 1999. Identification of a *Vibrio cholerae* RTX toxin gene cluster that is tightly linked to the cholera toxin prophage. Proc. Natl. Acad. Sci. USA **96:**1071–1076.
41. **Martins, M. T.** 1988. Ecologia de *Vibrio cholerae* no ecossistema aquático. Tese Livre Docente ICB-USP, Universidade de São Paulo, São Paulo, Brazil. (In Portuguese.).
42. **Martins, M. T., G. V. A. Pessôa, P. S. Sanchez, M. I. Z. Sato, C. A. Coimbrão, C. K. Monteiro, and E. Marques.** 1991. Occurrence of *V. cholerae* O1 nontoxigenic in wastewaters from São Paulo, Brazil. Wat. Sci. Tech. **24:**363–366.
43. **Matté, G. R., M. H. Matté, M. I. Z. Sato, P. S. Sanchez, I. G. Rivera, and M. T. Martins.** 1994. Potentially pathogenic vibrios associated with mussels from a tropical region on the Atlantic Coast of Brazil. J. Appl. Bacteriol. **77:**281–287.
44. **Matte, G. R., M. H. Matté, I. G. Rivera, and M. T. Martins.** 1994. Distribution of potentially vibrios in oysters from a tropical region. J. Food Prot. **57:**870–873.
45. **Mekalanos, J. J.** 1985. Cholera toxin: genetic analysis, regulation, and role in pathogenesis. Curr. Top. Microbiol. Immunol. **118:**97–118.
46. **Miller, V. L., R. K. Taylor, and J. J. Mekalanos.** 1987. Cholera toxin transcriptional activator ToxR is a transmembrane DNA binding protein. Cell **48:**271–279.
47. **Ministerio de Saúde-Fundação Nacional de Saúde.** 1999. MS/FNS, 1999—Boletin epidemiológico CENEPI, 1999. Ministerio de Saúde-Fundação Nacional de Saúde, Centro de Vigilância Epidemiologica, São Paulo, Brazil.
48. **Mooi, F. R., and E. M. Bik.** 1997. The evolution of epidemic *Vibrio cholerae* strains. Trends Microbiol. **5:**161–165.
49. **Morris, J. G., Jr., J. L. Picardi, S. Lieb, J. V. Lee, A. Roberts, M. Hood, R. A. Guun, and P. Blake.** 1984. Isolation of nontoxigenic *Vibrio cholerae* O group 1 from a patient with severe gastrointestinal disease. J. Clin Microbiol. **19:**196–197.
50. **Morris, J. G., Jr., T. Takeda, B. D. Tall, G. A. Losonsky, S. K. Bhattacharya,**

Downloaded from http://aem.asm.org/ on December 27, 2016 by University of North Texas Libraries

B. D. Forrest, B. A. Kay, and M. Nishibuchi. 1990. Experimental non-O1 group *Vibrio cholerae* gastroenteritis in humans. J. Clin. Investig. **85:**697–705.

51. **Morris, J. G.** 1994. Non-O1 group 1 *Vibrio cholerae* strains not associated with epidemic disease, p. 103–116. *In* I. K. Wachsmuth, P. A. Blake, and O. Olsvik (ed.), *Vibrio cholerae* and cholera: molecular to global perspectives. ASM Press, Washington, D.C.
52. **Mukhopadhyay, A. K., P. K. Saha, S. Garg, S. K. Bhattacharya, T. Shimada, T. Takeda, Y. Takeda, and G. B. Nair.** 1995. Distribution and virulence of *Vibrio cholerae* belonging to serogroups other than O1 and O139: a nation survey. Epidemiol. Infect. **114:**65–70.
53. **Nakasone, N., and Iwanaga, M.** 1998. Characterization of outer membrane protein OmpU of *Vibrio cholerae* O1. Infect. Immun. **66:**4726–4728.
54. **Nandi, B., R. K. Nandy, A. C. P. Vicente, and A. C. Ghose.** 2000. Molecular characterization of a new variant of toxin-coregulated pilus protein (TcpA) in a toxigenic non-O1/non-O139 strain of *Vibrio cholerae*. Infect. Immun **68:**948–952.
55. **Ogawa, A., J. Kato, H. Watanabe, B. G. Nair, and T. Takeda.** 1990. Cloning and nucleotide sequence of a heat-stable enterotoxin gene from *Vibrio cholerae* non-O1 isolated from a patient with traveler's diarrhea. Infect. Immun. **58:**3325–3329.
56. **Pal, A., T. Ramamurthy, R. K. Bhadra, T. Takeda, T. Shimada, Y. Takeda, G. B. Nair, S. C. Pal, and S. Chakrabarty.** 1992. Reassessment of the prevalence of heat-stable enterotoxin (NAG-ST) among environmental *Vibrio cholerae* non-O1 strains isolated from Calcutta, India, by using a NAG-ST DNA probe. Appl. Environ. Microbiol. **58:**2485–2489.
57. **Rader, A. E., and J. R. Murphy.** 1988. Nucleotide sequences and comparison of the hemolysin determinants of *Vibrio cholerae* El Tor RV79 (Hly^+) and RV79 (Hly^-) and Classical 569B (Hly^-). Infect. Immun. **56:**1414–1419.
58. **Ramamurthy, T., P. K. Bag, A. Pal, S. K. Bhattacharya, T. Shimada, T. Takeda, T. Karasawa, H. Kurasono, Y. Takeda, and G. B. Nair.** 1993. Virulence patterns of *Vibrio cholerae* non-O1 strains isolated from hospitalised patients with acute diarrhoea in Calcutta, India. J. Med. Microbiol. **39:**310–317.
59. **Ramamurthy, T., M. J. Albert, A. Huq, R. R. Colwell, Y. Takeda, T. Takeda, T. Shimada, B. K. Mandal, and G. B. Nair.** 1994. *Vibrio mimicus* with multiple toxin types isolated from human and environmental sources. J. Med. Microbiol. **40:**194–196.
60. **Rhine, J. A., and R. K. Taylor.** 1994. TcpA pilin sequences and colonization requirements for O1 and O139 *Vibrio cholerae*. Mol. Microbiol. **13:**1013–1020.
61. **Rivera, I. G., M. A. R. Chowdhury, A. Huq, D. Jacobs, M. T. Martins, and R. R. Colwell.** 1995. Enterobacterial repetitive intergenic consensus sequences and the PCR to generate fingerprints of genomic DNAs from *Vibrio cholerae* O1, O139, and non-O1 strains. Appl. Environ. Microbiol. **61:**2898–2904.
62. **Rivera, I. N. G., M. A. R. Chowdhury, P. S. Sanchez, M. I. Sato, A. Huq, R. R. Colwell, and M. T. Martins.** 1995. Detection of cholera (ctx) and zonula occludens (zot) toxin genes in *Vibrio cholerae* O1, O139, and non-O1 strains. World J. Microbiol. Biotech. **11:**572–577.
63. **Rodrigue, D. C., T. Popovic, and I. K. Wachsmuth.** 1994. Nontoxigenic *Vibrio cholerae* O1 infections in the United States, p. 69–76. *In* I. K. Wachsmuth, P. A. Blake, and O. Olsvik (ed.), *Vibrio cholerae* and cholera: molecular to global perspectives. ASM Press, Washington, D.C.
64. **Rodrigues, D. P., and E. Hofer.** 1986. *Vibrio* species from the water-oyster ecosystem of Sepetitiba Bay in Rio de Janeiro State, Brazil. Rev. Microbiol. **17:**332–338.
65. **Roszak, D. B., and R. R. Colwell.** 1987. Survival strategies of bacteria in the natural environment. Microbiol. Rev. **51:**365–379.
66. **Saha, P. K., H. Koley, A. K. Mukhopadhyay, S. K. Bhattacharya, G. B. Nair, B. S. Ramakrishnan, S. Krishnan, T. Takeda, and Y. Takeda.** 1996. Nontoxigenic *Vibrio cholerae* O1 serotype Inaba Biotype El Tor associated with a cluster of cases of cholera in Southern India. J. Clin. Microbiol. **34:**1114–1117.
67. **Said, B., H. R. Smith, S. M. Scotland, and B. Rowe.** 1995. Detection and differentiation of the gene for toxin co-regulated pili (*tcpA*) in *Vibrio cholerae* non-O1 using the polymerase chain reaction. FEMS Microbiol. Lett. **125:**205–210.
68. **Sakazaki, R., C. Z. Gomes, and M. Sebald.** 1967. Taxonomical studies of the so-called NAG vibrios. Jpn. J. Med. Sci. Biol. **20:**265–280.
69. **Shangkuan, Y. H., Y. S. Show, and T. M. Wang.** 1995. Multiplex polymerase chain reaction to detect toxigenic *Vibrio cholerae* and to biotype *Vibrio cholerae* O1. J. Appl. Bacteriol. **79:**264–273.
70. **Sharma, C., G. B. Nair, A. K. Mukhopadhyay, S. K. Bhattacharya, R. K. Ghosh, and A. Ghosh.** 1997. Molecular characterization of *Vibrio cholerae* O1 biotype El Tor strains isolates between 1992 and 1995 in Calcutta, India: evidence for the emergence of a new clone of the El Tor biotype. J. Infect. Dis. **175:**1134–1141.
71. **Sharma, C., M. Thungapathra, A. Ghosh, A. K. Mukhopadhyay, A. Basu, R. Mitra, I. Basu, S. K. Bhattacharya, T. Shimada, T. Ramamurthy, T. Takeda, S. Yamasaki, Y. Takeda, and G. B. Nair.** 1998. Molecular analysis of non-O1, non-O139 *Vibrio cholerae* associated with an unusual upsurge in the incidence of cholera-like disease in Calcutta, India. J. Clin. Microbiol. **36:**756–763.
72. **Sperandio, V., C. Bailey, J. A. Giron, V. J. DiRita, W. D. Silveira, A. L. Vettore, and J. B. Kaper.** 1996. Cloning and characterization of the gene encoding the OmpU outer membrane protein of *Vibrio cholerae*. Infect. Immun. **64:**5406–5409.
73. **Stroeher, U. H., K. E. Jedani, B. K. Dredge, R. Morona, M. H. Brown, L. E. Karageorgos, M. J. Albert, and P. A. Manning.** 1997. Genetic rearrangements in the *rfb* regions of *Vibrio cholerae* O1 and O139. Proc. Natl. Acad. Sci. USA **92:**10374–10378.
74. **Tauxe, R., L. Seminario, R. Tapia, and M. Libel.** 1994. The Latin American epidemic, p. 321–344. *In* I. K. Wachsmuth, P. A. Blake, and O. Olsvik (ed.), *Vibrio cholerae* and cholera: molecular to global perspectives. ASM Press, Washington, D.C.
75. **Taylor, R. K., V. L. Miller, D. B. Furlong, and J. J. Mekalanos.** 1987. Use of *phoA* gene fusions to identify a pilus colonization factor coordinately regulated with cholera toxin. Proc. Natl. Acad. Sci. USA **84:**2833–2837.
76. **Taylor, R. K.** 1989. Genetic studies of enterotoxin and other potential virulence factors of *Vibrio cholerae*, p. 309–329. *In* D. A. Hopwood and K. F. Chater (ed.), Genetics of bacterial diversity. Academic Press, Ltd., London, England.
77. **Taylor, R. K.** 1999. Virus on virus infects bacterium. Nature **399:**312–313.
78. **Twedt, R. M., J. M. Madden, J. M. Hunt, D. W. Francis, J. T. Peeler, A. P. Duran, W. O. Hebert, S. G. McCay, C. N. Roderick, G. T. Spite, and T. J. Wazenski.** 1981. Characterization of *Vibrio cholerae* isolated from oysters. Appl. Environ. Microbiol. **41:**1475–1478.
79. **Vicente, A. C., A. M. Coelho, and C. A. Salles.** 1997. Detection of *Vibrio cholerae* and *V. mimicus* heat-stable toxin gene sequence by PCR. J. Med. Microbiol. **46:**398–402.
80. **Waldor, M. K., and J. J. Mekalanos.** 1996. Lysogenic conversion by a filamentous phage encoding cholera toxin. Science **272:**1910–1914.
81. **World Health Organization.** 1969. Outbreak of gastroenteritis by nonagglutinable (NAG) vibrio. Wkly. Epidemiol. Rev. **44:**1–28.
82. **Yamamoto, K., M. Al-Omani, T. Honda, Y. Takeda, and T. Miwatani.** 1984. Non-O1 *Vibrio cholerae* hemolysin: purification, partial characterization, and immunological relatedness to El Tor hemolysin. Infect. Immun. **45:**192–196.
83. **Yamamoto, K., Y. Ichinose, N. Nakasone, M. Tanabe, M. Nagahama, J. Sakurai, and M. Iwanaga.** 1986. Identity of hemolysins produced by *Vibrio cholerae* non-O1 and *V. cholerae* O1, biotype El Tor. Infect. Immun. **51:**927–931.

Downloaded from http://aem.asm.org/ on December 27, 2016 by University of North Texas Libraries