

# Amino Acid Profiles and Presumptive Nutritional Assessment of Single-Cell Protein from Certain Lactobacilli<sup>1</sup>

M. D. ERDMAN, W. G. BERGEN, AND C. ADINARAYANA REDDY\*

Departments of Animal Husbandry and Microbiology and Public Health, Michigan State University, East Lansing, Michigan 48824

Received for publication 10 December 1976

The amino acid profiles, modified essential amino acid (MEAA) indexes, and in vitro pepsin digestibilities were determined for single-cell protein (SCP) from certain industrially important lactobacilli. For the three parameters examined, substantial differences were seen between different *Lactobacillus* species and between strains within a given species. SCP from all of the lactobacilli examined appeared relatively high in MEAA indexes and pepsin digestibility. SCP from *L. acidophilus* 3205 and *L. fermenti* 3954 had the highest MEAA indexes, whereas *L. bulgaricus* 2217 and *L. thermophilus* 3863 had the highest percentage of digestible crude protein. SCP from *L. plantarum* strains had the lowest MEAA indexes. The essential amino acid compositions of SCP from different lactobacilli appear comparable to that of Food and Agriculture Organization reference protein and SCP from other sources.

Lactic acid-producing bacteria of the genus *Lactobacillus* have been widely used in the past for the industrial production of lactic acid from whey, corn, potatoes, sulfite waste liquor, and a number of other materials rich in carbohydrates (1, 7, 8, 12, 13, 17). Lactobacilli are extensively used in the dairy industry for producing yogurt, cheese, buttermilk, and a variety of other fermented milk products, as well as in most pickle fermentations (6, 26). Reddy et al. (20) recently developed a process for the fermentative conversion of whey into a ruminant feed supplement in which approximately 7% of the crude protein is derived from lactic acid bacteria. In spite of the widespread use and importance of lactobacilli in various fermentation processes for the production of food or feed products, sparse data are available regarding the amino acid profiles of single-cell protein (SCP) derived from different lactobacilli, except for one report in 1945 by Camien et al. (5). These workers, by using microbiological assays, determined the essential amino acid contents of three species of lactobacilli. No data were presented on the nutritional quality of the SCP from these organisms. The present study was initiated to determine the amino acid composition of the SCP from 12 strains of industrially important lactobacilli, representing seven species, and to obtain a presumptive assessment of the nutritional value of SCP from these

organisms by determining the modified essential amino acid (MEAA) indexes and percent digestibility of the crude protein (total nitrogen  $\times$  6.25).

## MATERIALS AND METHODS

**Bacteria.** *Lactobacillus acidophilus* 3205 and 3532, *L. bulgaricus* 3533, *L. fermenti* 8954 and 3957, *L. plantarum* 3074, and *L. thermophilus* 3863 were obtained from the Institute for Fermentation, Osaka, Japan. *L. casei* 14435 and *L. plantarum* 14431 and 8014 were obtained from the American Type Culture Collection, Rockville, Md. *L. bulgaricus* 2217 was obtained from Chris Hansen Laboratories, Milwaukee, Wis., and *L. delbrueckii* B-443, supplied by R. Costilow, was originally obtained from the Northern Regional Research Laboratory, Agricultural Research Service, U.S. Department of Agriculture, Peoria, Ill. All stock cultures were maintained on Trypticase soy agar deeps and transferred once every 4 weeks to tubes of the same medium. All tubes, after inoculation, were incubated at 37°C for 24 h and then stored at 4°C until used.

**Preparation of cells.** A small amount of growth from 24-h Trypticase soy agar slant culture was transferred aseptically with an inoculating needle to 10 ml of sterile Trypticase soy broth (TSB) in screw-cap tubes (18 by 150 mm) and were incubated at 37°C for 24 h. These TSB cultures were used to inoculate 500 ml of TSB contained in 1-liter, foam-plugged Erlenmeyer flasks which, after incubation as described above, were used to inoculate 3.5 liters of TSB contained in 4-liter, foam-plugged Erlenmeyer flasks.

**Harvesting and protein extraction.** After 24 h of

<sup>1</sup> Journal article no. 7884, Michigan Agricultural Experiment Station.

incubation at 37°C, cells from the 4-liter cultures were harvested by centrifugation at  $45,900 \times g$  for 15 min with a centrifuge (Dupont-Sorvall model RC-5) equipped with a continuous-flow centrifuge head. The centrifuged cell paste was washed in 10 volumes of 0.2 M sodium phosphate buffer (pH 7.0) and centrifuged ( $25,000 \times g$ ) for 15 min. The supernatant was discarded, and the cell pellet was washed twice in 10 volumes of distilled water and collected by centrifugation as described above. The washed cell pellet was collected in a glass vial, resuspended in 10 volumes of distilled water, shell-frozen, and lyophilized (Cryolizer freeze dryer, model B64, New Brunswick Scientific Co., New Brunswick, N.J.). For cold trichloroacetic acid extraction, about 1.7 g of lyophilized cells was suspended in 100 ml of 5.0% (wt/vol) trichloroacetic acid, incubated for 30 min at 25°C, and centrifuged at  $25,000 \times g$  for 15 min. The supernatant was discarded, and the pellet was dried (60°C) in a forced-air oven to constant weight. Cold trichloroacetic acid extraction was used for extracting SCP from eight strains of lactobacilli representing six species.

For hot trichloroacetic acid extraction, cells were harvested as described above, washed with 200 ml of distilled water, and collected by centrifugation at  $25,000 \times g$  for 15 min. The washed cells were suspended in 200 ml of 5.0% trichloroacetic acid, incubated at 90°C for 30 min, and centrifuged at  $25,000 \times g$  for 15 min. The supernatant was discarded, and the precipitated protein was suspended in 50 ml of distilled water, shell-frozen, and lyophilized. Hot trichloroacetic acid extraction was used for extracting SCP from four species of lactobacilli, because this procedure is reported to extract nonprotein nitrogen more completely than cold trichloroacetic acid extraction (21, 22). As expected, hot trichloroacetic acid extraction in general resulted in a higher amino acid concentration, as compared with the cold trichloroacetic acid extraction of the same SCP.

**Protein hydrolysis.** Total nitrogen (TN) was determined on triplicate samples of the SCP from each organism by the micro-Kjeldahl technique (25), with HgO as a catalyst. Crude protein (CP) was defined as  $TN \times 6.25$ . Two aliquots containing 10 mg of CP from each of the SCP sources were placed in separate glass screw-cap vials. One milliliter of a solution containing 1  $\mu$ mol each of norleucine and *S*- $\beta$ -(4-pyridylethyl)-L-cysteine (internal standards) and 9 ml of 6.4 N HCl was added to each glass vial. The vials were flushed with nitrogen, sealed, and heated at 121°C and 1.06-kg/cm<sup>2</sup> pressure for 24 h. The cooled acid hydrolysates were filtered through a fluted filter paper (Whatman no. 2). The filtrates were evaporated to dryness (60°C) in a Buchler flash evaporator (model PF, A. H. Thomas Co., Philadelphia, Pa.), resuspended in 10 ml of distilled water, and evaporated to dryness two more times. Four milliliters of 0.05 M citrate buffer (pH 2.0) containing 0.3 N LiCl was added to the dried acid hydrolysates, and the solutions were transferred to glass screw-cap vials and stored at -4°C until analyzed for amino acids with an amino acid analyzer (Technicon TSM-1, Technicon Corp., Tarrytown, N. Y.).

**In vitro pepsin digestibility.** In vitro pepsin digestibility was determined by the procedure of the As-

sociation of Official Agricultural Chemists (2), as modified below. Smaller protein digestions were carried out in 50 ml of Oak Ridge-type polycarbonate centrifuge tubes. After pepsin treatment, the indigestible CP was precipitated by treatment with 5% trichloroacetic acid for 30 min at 25°C, followed by centrifugation at  $48,200 \times g$  for 30 min. TN was determined on duplicate samples of supernatant and pellet.

**MEAA index.** MEAA index was determined by the procedure for Oser (16), as modified by Mitchell (15), except that tryptophan concentrations were not determined and were not included in the calculations of the MEAA indexes.

## RESULTS AND DISCUSSION

**Amino acid composition.** The results presented in Table 1 show that there are substantial differences in amino acid composition of SCP from different *Lactobacillus* species. SCP from *L. plantarum* and *L. fermenti* 3957 differed from that of the other lactobacilli in that it had a lower amount of lysine. SCP from *L. plantarum* 14431 and 3074 had an approximately three times greater concentration of the essential amino acid, isoleucine, as compared with the concentration of the same amino acid in SCP from *L. plantarum* 8014 as well as in other *Lactobacillus* species. Aspartic acid concentration was quite high in SCP from all lactobacilli, with the exception of that from *L. fermenti* 3957 and *L. plantarum* 14431 and 3074. High levels of aspartic acid have also been reported in rumen bacteria (3).

In Table 1, the amino acid composition of the SCP from lactobacilli is compared with that of the Food and Agriculture Organization (FAO) reference protein (27), certain other SCP (11, 23), and conventional foods (4, 9, 15, 23). The results showed that the concentration of essential amino acids in SCP from most lactobacilli was equal to or greater than those in the FAO reference protein, suggesting that lactobacilli are a good protein source for human adults. However, SCP from *L. plantarum* 8014 was low in leucine. The amino acid profile of the latter strain was substantially different from that of the other two *L. plantarum* strains examined in that it had a lower concentration of leucine and isoleucine and a higher concentration of lysine.

Most SCP produced to date are deficient in methionine (8, 11, 24). SCP from lactobacilli examined in this study were also low in methionine, compared with the amount in FAO reference protein. However, lactobacilli SCP were relatively superior to soybean protein and wheat flour in their methionine content. Methionine is somewhat labile to acid hydrolysis (14). Therefore, the values presented in Table 1 for

TABLE 1. Amino acid profiles of SCP from lactobacilli, FAO reference protein, and certain SCP and conventional foods<sup>a</sup>

Protein source <sup>b</sup>	Essential amino acids										Nonessential amino acids							Source
	ILE	LEU	LYS	MET	PHE	THR	VAL	TYR	ALA	ARG	ASP	Half-CYS	GLU	GLY	HIS	PRO	SER	
Cold trichloroacetic acid extraction	4.3	7.4	10.4	1.9	3.3	4.0	4.9	2.5	9.0	5.1	9.7	0.2	11.1	4.2	2.4	3.5	2.5	
	4.5	6.1	9.3	2.2	3.2	4.3	5.8	3.3	6.0	4.5	10.5	0.4	9.1	3.5	2.2	3.6	2.6	
	4.2	6.5	7.9	1.9	3.2	4.3	4.9	2.7	7.2	4.0	10.0	0.3	9.8	4.0	1.9	3.0	2.3	
	5.0	6.8	10.2	2.1	4.0	4.2	5.4	3.2	6.8	5.3	11.0	0.5	12.3	4.4	2.7	2.9	3.1	
	4.4	6.3	7.1	2.0	3.1	3.6	4.9	2.4	8.9	5.1	7.7	0.1	10.3	4.2	2.2	2.6	2.7	
	12.3	5.4	4.5	1.8	2.6	3.4	4.2	2.2	9.3	3.0	7.0	0.0	11.5	3.5	1.4	3.1	2.4	
Hot trichloroacetic acid extraction	13.2	5.7	4.9	1.8	2.9	3.3	4.9	2.5	9.8	3.3	7.8	0.2	11.9	3.4	1.7	2.5	2.5	
	4.1	6.4	9.0	1.9	3.2	3.7	4.8	2.6	8.4	4.3	10.6	0.2	9.7	4.2	2.2	2.4	2.3	
	6.4	8.2	10.3	2.1	4.2	4.8	6.4	3.5	8.4	4.9	13.3	0.3	12.4	4.4	2.5	3.4	2.8	
	5.5	7.6	9.6	1.9	4.2	4.4	5.9	3.3	7.5	4.6	14.6	0.5	14.6	4.3	2.0	3.3	2.8	
	4.6	8.0	10.2	2.2	4.1	6.2	6.7	4.2	7.6	4.3	13.5	0.4	10.9	4.4	2.4	3.1	3.2	
	4.6	1.5	9.0	1.9	3.6	4.3	5.5	3.0	7.2	4.3	11.8	0.3	10.4	3.6	2.2	3.0	2.5	
FAO reference protein	4.0	7.0	5.5	3.5 <sup>c</sup>	6.0 <sup>d</sup>	4.0	5.0										27	
	4.7	11.2	6.8	1.9	4.4	5.4	10.7	2.7									23	
	5.6	5.9	5.2	1.1	2.8	3.8	5.2										5	
	6.2	6.8	7.7	1.1	3.5	4.9	5.8										5	
	7.0	7.5	6.9	1.3	4.1	4.9	6.8										5	
	4.6	7.0	7.7	1.7	4.1	4.8	5.3										11	
	5.3	8.2	8.6	2.5	4.1	4.4	5.5	3.4									9	
	5.9	8.8	7.8	3.2	5.5	4.9	7.1										15	
	5.8	7.6	6.6	1.1	4.8	3.9	5.2	3.2									4	
	4.2	7.0	1.9	1.5	5.5	2.7	4.1										23	

<sup>a</sup> Amino acid concentrations are expressed as grams of amino acid per 16 g of N.  
<sup>b</sup> SCP was extracted with 10% (wt/vol) trichloroacetic acid at 25° for organisms under cold trichloroacetic acid. For organisms listed under hot trichloroacetic acid, SCP was extracted with 10% (wt/vol) trichloroacetic acid at 90°C; details are given in Materials and Methods.  
<sup>c</sup> Value reported is the sum of values for methionine + cystine.  
<sup>d</sup> Value reported is the sum of values for phenylalanine + tyrosine.

methionine underestimate the actual methionine concentration in the SCP of the lactobacilli.

Lysine is the most limiting essential amino acid in cereals, but it is present in higher concentration in animal protein sources such as beef and egg. The lysine concentration in SCP from *L. thermophilus*, *L. bulgaricus* 2217, *L. acidophilus* 3532 and 3205, *L. delbrueckii*, *L. fermenti* 3954, *L. plantarum* 8014, and *L. casei* strains surpassed that in beef or whole egg and was approximately twice as much as that present in *L. plantarum* 14431 and 3074.

Camien et al. (5) previously reported the essential amino acids of *L. plantarum* 8014 and 8041, *L. casei* 7469, and *L. fermenti* 9338. Defatted lactobacilli were acid-hydrolyzed by refluxing cells for 20 h in 10 volumes of 8 N HCl, and the amino acid content was determined by microbiological assay. The procedures used were substantially different from ours, and this may account for the small differences observed in essential amino acids reported by Camien et al. These workers suggested that proteins in different strains of microorganisms do not differ markedly in amino acid composition. Our results indicated differences in amino acid composition between SCP from different species of lactobacilli and among strains within a given species. Similar variations in amino acid compositions of SCP from different organisms have been reported by other investigators (19, 24).

**In vitro** protein digestibility and MEAA index. The amino acid concentrations present in a protein source may have no relationship to the biological availability of these amino acids (18). To make a preliminary assessment of the nutritional value of the SCP from lactobacilli, the MEAA index and *in vitro* pepsin digestibility were determined for each SCP type (Table 2). The MEAA index is a measure of the levels of all the essential amino acids in the test protein versus those in whole egg protein (18). Considerable differences in MEAA indexes and digestible crude protein (CP) for the SCP from different lactobacilli were observed. The SCP from *L. acidophilus* 3205 and *L. fermenti* 3954 appeared to be relatively superior to other lactobacilli, with MEAA indexes of 86 and 85, respectively. SCP from *L. bulgaricus* 2217 and *L. thermophilus* 3863 had the highest digestible CP values, 89.2 and 88.1, respectively, but had relatively low MEAA indexes, 76 and 69, respectively. MEAA indexes and digestible CP values were relatively high for *L. casei* 14435 and *L. delbrueckii* B-443. SCP from *L. plantarum* strains had lower MEAA indexes and digestible CP values, suggesting that they are lower-quality

TABLE 2. MEAA index and percent digestible CP of SCP from certain lactobacilli

SCP source	MEAA index	Digestible CP (%)
Casein	91	98.5 <sup>a</sup> ± 0.2 <sup>b</sup>
<i>L. acidophilus</i> 3532	73	79.3 ± 0.5
<i>L. acidophilus</i> 3205	86	83.7 ± 0.5
<i>L. bulgaricus</i> 2217	76	89.2 ± 0.1
<i>L. bulgaricus</i> 3533	69	81.3 ± 0.4
<i>L. casei</i> 14435	80	82.3 ± 0.2
<i>L. delbrueckii</i> B-443	80	82.5 ± 0.2
<i>L. fermenti</i> 3954	85	86.5 ± 0.5
<i>L. fermenti</i> 3957	69	81.6 ± 0.5
<i>L. plantarum</i> 14431	59	79.9 ± 0.8
<i>L. plantarum</i> 8014	62	80.6 ± 0.3
<i>L. thermophilus</i> 3863	69	88.1 ± 0.3

<sup>a</sup> Calculated from reference 10. Values giving standard deviation greater than 1.0 were not included in the calculation of the mean percent digestible crude protein.

<sup>b</sup> Standard error of the mean.

protein sources than casein and the SCP from other lactobacilli.

Lactobacilli are commonly used in industry for the production of fermented food or feed products. The lactobacilli SCP in these fermented foods generally constitutes only a small percentage of the total protein. However, in any one of these processes, it would be more desirable to use a strain of *Lactobacillus* that has a higher quality SCP than a strain with a SCP of poorer quality, other considerations being equal. Reddy et al. (20) recently used *L. bulgaricus* 2217 for the fermentation of whey and produced a product called fermented ammoniated condensed whey (FACW), which is rich in CP (55% CP) and was shown to be an efficacious nitrogen supplement for cattle. Approximately 7% of the CP in FACW comes from *L. bulgaricus*, 17% is from whey proteins, and 70% is from ammonium lactate. Considering the quantity of SCP from *L. bulgaricus* 2217 in this product and its relatively superior amino acid profile, MEAA index, and percent digestible CP, it appears possible that *L. bulgaricus* SCP and whey proteins (mostly lactalbumin, which has a high biological value for humans) could be separated from ammonium lactate in FACW and used as a source of protein for humans. These studies are currently in progress in our laboratory.

#### ACKNOWLEDGMENTS

This research was supported by grants from Calor Agriculture Research, Okemos, Mich., and the MSU Foundation, Michigan State University, East Lansing, Mich.

#### LITERATURE CITED

- Anderson, A. A., and J. E. Graves. 1942. d-Lactic acid fermentation of Jerusalem artichokes. *Ind. Eng. Chem.* 34:1522-1526.

2. Association of Official Agricultural Chemists. 1975. Official methods of analysis, 12th ed., p. 133-135. Association of Official Agricultural Chemists, Washington, D.C.
3. Bergen, W. G., D. B. Purser, and J. H. Cline. 1968. Determination of limiting amino acids of rumen-isolated microbial proteins fed to rats. *J. Dairy Sci.* 51:1698-1700.
4. Callihan, C. D., and C. F. Dunlap. 1969. The economics of microbial proteins produced from cellulosic wastes. *Compost Sci.* 10:6-12.
5. Camien, M. N., A. J. Salle, and M. S. Dunn. 1945. Investigations of amino acids, peptides and proteins. XXII. Percentages of some amino acids in *Lactobacilli*. *Arch. Biochem.* 8:67-78.
6. Carr, J. C., C. V. Cutting, and G. C. Whiting (eds.). 1973. Lactic acid bacteria in beverages and food. Academic Press Inc., New York, N.Y.
7. Cordon, T. C., R. H. Treadway, M. D. Walsh, and M. F. Osborne. 1950. Lactic acid from potatoes. *Ind. Eng. Chem.* 42:1833-1836.
8. Fred, E. B., and W. H. Peterson. 1921. Fermentation process for the production of acetic and lactic acids from corncobs. *Ind. Eng. Chem.* 13:211-213.
9. Food and Agriculture Organization of the United Nations. 1959. Protein requirements, p. 26. FAO nutritional studies, no. 16. Rome, Italy.
10. Food and Agriculture Organization of the United Nations. 1970. Amino acid content of foods and biological data on proteins. FAO nutritional studies, no. 24. Rome, Italy.
11. Kihlberg, R. 1972. The microbe as a source of food. *Annu. Rev. Microbiol.* 26:427-466.
12. Leonard, R. H., W. H. Peterson, and M. J. Johnson. 1948. Lactic acid from fermentation of sulfite waste liquor. *Ind. Eng. Chem.* 40:57-67.
13. Marth, E. H. 1970. Fermentation products from whey, p. 43-81. *In* B. H. Webb and E. O. Whittier (ed.), *Byproducts from milk*. Avi Publishing Co., Westport, Conn.
14. Menden, E., and H. D. Cremer. 1970. Laboratory methods for the evaluation of changes in protein quality, p. 123-161. *In* A. A. Albanese (ed.), *Newer methods of nutritional biochemistry with applications and interpretations*. Academic Press Inc., New York, N.Y.
15. Mitchell, H. H. 1954. Biological value of proteins and amino acid interrelationships, p. 13-28. *In* H. Spector, M. S. Peterson, and T. E. Friedman (ed.), *Symposium on methods for the evaluation of nutritional adequacy and status*, p. 13. National Research Council, Washington, D.C.
16. Oser, B. L. 1951. Method for integrating essential amino acid content in the nutritional evaluation of protein. *J. Am. Diet. Assoc.* 27:396-402.
17. Prescott, S. C., and C. G. Dunn. 1959. The production of lactic acid by fermentation, p. 304-369. *In* *Industrial Microbiology*, 3rd ed. McGraw-Hill Book Co., Inc., New York, N.Y.
18. Protein-Calorie Advisory Group of the United Nations System. 1974. Fourth meeting of the PAG *ad hoc* working group on single cell protein. *PAG Bull.* 4:9-31.
19. Purser, D. R., and S. M. Buechler. 1966. Amino acid composition of rumen organisms. *J. Dairy Sci.* 49:81-84.
20. Reddy, C. A., H. E. Henderson, and M. D. Erdman. 1976. Bacterial fermentation of cheese whey for production of a ruminant feed supplement rich in crude protein. *Appl. Environ. Microbiol.* 32:769-776.
21. Schmidt, G., and S. J. Thannhauser. 1945. A method for the determination of deoxyribonucleic acid, ribonucleic acid and phosphoproteins in animal tissues. *J. Biol. Chem.* 161:83-89.
22. Schneider, W. C. 1945. Phosphorus compounds in animal tissues. Extraction and elimination of desoxypentose nucleic acid and of pentose nucleic acid. *J. Biol. Chem.* 161:293-303.
23. Srinivasan, V. A., and Y. W. Han. 1969. Utilization of bagasse. *Adv. Chem. Ser.* 95:447-460.
24. Tannenbaum, S. R., and D. I. C. Wang (ed.). 1975. *Single-cell protein II*. The MIT Press, Cambridge, Mass.
25. Wall, L. L., and C. W. Gehrke. 1975. An automated total protein nitrogen method. *J. Assoc. Off. Anal. Chemists.* 58:1221-1226.
26. Webb, B. H., and E. O. Whittier (ed.). 1970. *By-products from milk*. Avi Publishing Co., Westport, Conn.
27. World Health Organization. 1973. Energy and protein requirements, p. 1-118. *In* *Technical Report Series no. 522*, FAO nutrition meetings report series no. 52. World Health Organization, Geneva.