

NOTES

Only Part of the Protoxin Gene of *Bacillus thuringiensis* subsp. *berliner* 1715 Is Necessary for Insecticidal Activity†

HIROETSU WABIKO, GARY A. HELD,‡ AND LEE A. BULLA, JR.*

Department of Biochemistry, College of Agriculture, University of Wyoming, Laramie, Wyoming 82071

Received 22 October 1984/Accepted 12 December 1984

***Escherichia coli* strains harboring deletion mutations of the insecticidal protoxin gene of *Bacillus thuringiensis* subsp. *berliner* 1715 were constructed. Although these strains did not produce intact protoxin, cell extracts from one of the mutants were extremely toxic to tobacco hornworm (*Manduca sexta*) larvae, indicating that only a part of the protoxin gene is required for insecticidal activity.**

The gram-positive bacterium *Bacillus thuringiensis* produces an insecticidal proteinaceous parasporal crystal during sporulation (4). The crystal is composed of a monomeric glycoprotein subunit ($M_r = 134$ kilodalton [kDa]), a protoxin (1) that is converted to a lower-molecular-weight toxin ($M_r = 68$ kDa) upon ingestion by a susceptible insect (1, 2, 5, 9). The toxin possesses the same specific toxicity as the protoxin (2, 3), suggesting that a part of the protoxin is inconsequential to insecticidal activity.

To genetically characterize the protoxin and toxin and to better understand the regulation of their syntheses, chromosomal and plasmid genes encoding for protoxin have been isolated from several *B. thuringiensis* subspecies (6, 7, 11). In the present paper, we describe an *Escherichia coli* clone that contains only a part of the protoxin gene of *B. thuringiensis* subsp. *berliner* 1715 but that synthesizes proteins whose toxicity is virtually identical to that of native protoxin.

The protoxin gene was isolated from a 42-megadalton plasmid of *B. thuringiensis* subsp. *berliner* 1715 (kindly provided by G. Rapoport, Institut Pasteur) as described by Klier et al. (7) except that we used pUC13 (14) as a cloning vector and *E. coli* JM83 (14) as a host. The clone, designated pHW13, contains 14.5 kilobase pairs (kbp) of *Bam*HI insert derived from the *B. thuringiensis* subsp. *berliner* 1715 plasmid. A physical map is presented in Fig. 1. It has been shown (7) that the major coding region of the protoxin resides in a 3.0-kbp *Pst*I-*Sst*I subfragment of a 5.2-kbp *Sst*I fragment (Fig. 1). Therefore, for further subcloning, pHW13 DNA was digested completely with either *Pst*I or *Sst*I and religated with T4 DNA ligase. *E. coli* JM83 was transformed with the ligated DNAs with CaCl_2 (10), and transformants were screened for ampicillin ($100 \mu\text{g ml}^{-1}$) resistance on L-agar plates. Thus, clones pP1 from *Pst*I digestion and pS11 from *Sst*I digestion were obtained (Fig. 1). Both clones contained the 3.0-kbp *Pst*I-*Sst*I insert. Intact protoxin was produced by the pP1 clone but not by pS11 (see below).

To further truncate pP1 DNA, the pP1 clone was sequentially deleted beginning at the *Pst*I site by partial digestion with *Hind*III. Although it has been reported (7) that the 14.5-kbp *Bam*HI fragment does not contain a *Hind*III cleavage site, we found three such sites, two of which were located in pP1 DNA (Fig. 1). The 14.5-kbp *Bam*HI insert in pHW13 is probably different from that in pBT42-1 isolated by Klier et al. (7), although the restriction pattern, other than that for *Hind*III, is quite similar. An excess amount of pP1 DNA (about $20 \mu\text{g}$) was cleaved with *Hind*III (8 U) for a limited period (20 min) at 37°C . The cleavage products were separated by gel electrophoresis with 0.6% low-gelling-temperature agarose (Sigma Chemical Co., St. Louis, Mo.). Portions of agarose containing relevant sizes of DNA were cut and melted at 65°C , and the DNA was extracted by phenol treatment at 37°C and ethanol precipitated. DNA obtained in such a manner was self-ligated and used to transform strain JM83. Ampicillin-resistant transformants were selected as described above, and clones pH2 and pH1 were obtained (Fig. 1).

To examine the expression of the genes subcloned in *E. coli* with respect to protoxin antigen production and insect toxicity, we prepared cell extracts of *E. coli*. Early-stationary-phase cells (grown in L broth containing $100 \mu\text{g}$ of ampicillin ml^{-1} at 37°C) of strain JM83 harboring each subcloned plasmid were concentrated 25-fold in a solution of lysozyme (1 mg ml^{-1} in 10 mM EDTA, pH 8.5), incubated for 30 min at 25°C , and then sonicated (Branson sonifier model 350, output control 3, 30% duty, 10 bursts). The resulting lysates were adjusted to pH 11.7 with 1 N NaOH. After the lysates were stirred for 2.5 h at 25°C , 1/20 volume of 2 M Tris hydrochloride (pH 8.3) was added to them, and they were cleared by centrifugation ($13,000 \times g$ for 5 min). The proteins in the extracts were separated by sodium dodecyl sulfate-7% polyacrylamide gel electrophoresis as described by Laemmli (8). The proteins in the gel were transferred electrophoretically (5 W for 3 h) to a nitrocellulose filter (Schleicher & Schuell, Inc., Keene, N.H.). The presence of protoxin was detected by enzyme-linked immunochromatography (12), with the following modifications. Gelatin (3% [wt/vol]) was used instead of Tween 20 to prevent nonspecific binding of the antibody to the nitrocellulose filter. Phosphate buffer plus saline was replaced by 20 mM Tris hydrochloride (pH 7.5) and 500 mM NaCl through-

* Corresponding author.

† Contribution no. 1336 of the Wyoming Agricultural Experiment Station, Laramie, WY 82071.

‡ Present address: Health Effects Research Laboratory (MD67), Environmental Protection Agency, Research Triangle Park, NC 27711.




FIG. 1. Plasmids containing *B. thuringiensis* subsp. *berliner* 1715 protoxin gene and its deletion mutations. The plasmid pHW13 carried a 14.5-kbp *Bam*HI fragment derived from a 42-megadalton *B. thuringiensis* subsp. *berliner* 1715 plasmid. It is shown as a linearized form with a unique *Sal*I (Sa) site in pUC13 as reference. Subclone pP1 was obtained from pHW13 by *Pst*I digestion. Deletion derivatives pS11, pH1, and pH2 were constructed from pP1 by using either *Sst*I or *Hind*III digestion as described in the text. Solid bar, *B. thuringiensis* subsp. *berliner* DNA; open bar, pUC13 DNA; cleavage sites, B, *Bam*HI; H, *Hind*III; P, *Pst*I; S, *Sst*I; wavy bar, predicted protoxin gene locus. Numbers represent kilobase pairs of DNA.




FIG. 2. Immunochemical detection of proteins in *E. coli* cell extracts. Early-stationary-phase cells of strain JM83 harboring each plasmid as described in the legend to Fig. 1 were concentrated 25-fold. Extracts were prepared as described in the text. About 80

μg of protein was separated by sodium dodecyl sulfate-7% polyacrylamide gel electrophoresis. After the proteins were transferred to a nitrocellulose filter, the filter was treated with antibody against *B. thuringiensis* subsp. *kurstaki* solubilized crystal which is known to react with *B. thuringiensis* subsp. *berliner* solubilized protoxin (13). The proteins that reacted with the antibody were stained as described in the text. Lanes: 1, pP1; 2, pH2; 3, pH1; 4, pS11; 5, pUC13. Molecular masses (kilodaltons) of the proteins indicated by arrows are: A, 140; B, 100; C, 65; D, 54. These values were estimated by comparing their relative mobilities with those of the following standard proteins (obtained from Bio-Rad Laboratories, Richmond, Calif.) stained with Coomassie blue: myoglobin, 200 kDa; β-galactosidase, 116 kDa; phosphorylase b; 93 kDa; bovine serum albumin, 66 kDa; ovalbumin, 45 kDa. The symbol ● represents proteins that reacted with antibody from the pP1 clone, except protein A (lane 1).

Apparently, the 54-kDa protein cross-reacted nonspecifically with the antibody, because this protein was present in every clone including strain JM83 containing pUC13 alone (lane 5). The eight proteins in question could be degradation products of the intact protoxin. In the case of the pH2 clone, a 100-kDa protein was produced (Fig. 2, band B, lane 2) instead of the 140-kDa protein. If this 100-kDa protein is a truncated protoxin gene product, the size of the deletion in the pH2 protoxin gene should correspond to approximately 1 kbp of DNA. Therefore, we believe that one terminal of the protoxin gene resides at about 0.3 kbp from the *Pst*I site in the pP1 clone (Fig. 1). Furthermore, a 65-kDa protein (Fig. 2, band C, lanes 1 and 2) in deletion mutant pH2 is more prominent than in pP1. This protein may be incomplete protoxin or a degradation product of the complete protoxin. It should be noted that the size of this 65-kDa protein is very close to that of the toxin ($M_r = 68$ kDa; see reference 3). When further deletion was accomplished, such as in pH1, none of the specific proteins mentioned above was obvious (Fig. 2, lane 3). No protoxin antigen was produced by the pS11 clone (Fig. 2, lane 4). Because the 3.0-kbp *Pst*I-*Sst*I fragment was common to both pP1 and pS11 (Fig. 1) and the pP1 clone produced intact protoxin, we believe that the other terminal of the gene is outside the *Sst*I fragment of pS11 (Fig. 1).

Insect toxicity of the *E. coli* cell extracts was tested with tobacco hornworm (*Manduca sexta*) larvae by the method of Bulla et al. (3). One larva was placed on 10 ml of solid artificial diet (Carolina Biological Supply Co., Burlington, N.C.) in a single cup (4 by 7 cm) onto which the cell extracts had been uniformly distributed. The cups containing larvae were incubated at 25°C. The extracts from pHW13, pP1, and pH2 were toxic. In fact, the specific toxicity of each clone, deduced from the amount of protoxin in *E. coli* cell extracts (0.1 to 0.5% of the total protein), was comparable to that of solubilized parasporal crystal (50% lethal concentration, 1.36 ng cm⁻², as calculated by Tyrell et al. [13]). Conversely, no toxicity was observed with extracts of pH1, pS11, and pUC13 clones under the same conditions. This result demonstrates that the pH2 clone has a toxicity nearly identical to that of the original pHW13 plasmid and to that of the pP1 clone and that only part of the protoxin gene is required for toxicity. Possibly, the 100- or 65-kDa protein or both were responsible for the toxicity of the pH2 clone. Because pH1 did not exhibit toxicity and because the 0.9-kbp *Hind*III fragment was absent in this clone (Fig. 1), this particular fragment apparently contains the site that delimits the portion of DNA required for insecticidal activity in the entire protoxin gene. We are currently attempting to localize more precisely the region of DNA that contains the protoxin gene.

This research was funded by grants from Stauffer Chemical Co., Richmond, Calif., and the Agency for International Development, U.S. Department of Agriculture, Washington, D.C.

LITERATURE CITED

1. Bulla, L. A., Jr., D. B. Bechtel, K. J. Kramer, Y. I. Shethna, A. I. Aronson, and P. C. Fitz-James. 1980. Ultrastructure, physiology and biochemistry of *Bacillus thuringiensis*. *Crit. Rev. Microbiol.* **8**:147-203.
2. Bulla, L. A., Jr., L. I. Davidson, K. J. Kramer, and B. J. Jones. 1979. Purification of the insecticidal toxin from the parasporal crystal of *Bacillus thuringiensis* subsp. *kurstaki*. *Biochem. Biophys. Res. Commun.* **91**:1123-1130.
3. Bulla, L. A., Jr., K. J. Kramer, D. J. Cox, B. L. Jones, L. I. Davidson, and G. L. Lookhart. 1981. Purification and characterization of the entomocidal protoxin of *Bacillus thuringiensis*. *Biol. Chem.* **256**:3000-3004.
4. Bulla, L. A., Jr., K. J. Kramer, and L. I. Davidson. 1977. Characterization of the entomocidal parasporal crystal of *Bacillus thuringiensis*. *J. Bacteriol.* **130**:375-383.
5. Cooksey, K. E. 1971. The protein crystal toxin of *Bacillus thuringiensis*: biochemistry and mode of action, p. 247-274. In H. D. Burges and N. W. Hussey (ed.), *Microbial control of insects and mites*. Academic Press, Inc., New York.
6. Held, G. A., L. A. Bulla, Jr., E. Ferrari, J. Hoch, A. I. Aronson, and S. A. Minnich. 1982. Cloning and localization of the lepidopteran protoxin gene of *Bacillus thuringiensis* subsp. *kurstaki*. *Proc. Natl. Acad. Sci. U.S.A.* **79**:6065-6069.
7. Klier, A., F. Fargette, J. Ribier, and G. Rapoport. 1982. Cloning and expression of the crystal protein genes from *Bacillus thuringiensis* strain *berliner* 1715. *EMBO J.* **1**:791-799.
8. Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (London)* **227**:680-685.
9. Lilley, M., R. N. Ruffell, and H. J. Somerville. 1980. Purification of the insecticidal toxin in crystals of *Bacillus thuringiensis*. *J. Gen. Microbiol.* **118**:1-11.
10. Mandel, M., and A. Higa. 1970. Calcium-dependent bacteriophage DNA infection. *J. Mol. Biol.* **53**:159-162.
11. Schnepf, H. E., and H. R. Whiteley. 1981. Cloning and expression of the *Bacillus thuringiensis* crystal protein gene in *Escherichia coli*. *Proc. Natl. Acad. Sci. U.S.A.* **78**:2893-2897.
12. Tsang, V. C. W., J. M. Perata, and A. R. Simons. 1983. Enzyme-linked immunoelectrotransfer blot techniques (EITB) for studying the specificities of antigens and antibodies separated by gel electrophoresis. *Methods Enzymol.* **92**:377-391.
13. Tyrell, D. J., L. A. Bulla, Jr., R. E. Andrews, Jr., K. J. Kramer, L. I. Davidson, and P. Nordin. 1981. Comparative biochemistry of entomocidal parasporal crystals of selected *Bacillus thuringiensis* strains. *J. Bacteriol.* **145**:1052-1062.
14. Vieira, J., and J. Messing. 1982. The pUC plasmids, an M13mp⁺ derived system for insertion mutagenesis and sequencing with synthetic universal primers. *Gene* **19**:259-268.