

Bacterial and Fungal Numbers in Ruminal and Cecal Contents of the Blue Duiker (*Cephalophus monticola*)†

BURK A. DEHORITY^{1*} AND GABRIELLA A. VARGA²

Department of Animal Science, Ohio Agricultural Research and Development Center, The Ohio State University, Wooster, Ohio 44691,¹ and Department of Dairy and Animal Science, The Pennsylvania State University, University Park, Pennsylvania 16802²

Received 21 September 1990/Accepted 10 December 1990

Total and cellulolytic bacterial and fungal numbers were determined in ruminal and cecal contents of 20 blue duikers (*Cephalophus monticola*). The animals were equally divided by sex and fed two diets, either high roughage or high concentrate. The mean concentration for total bacterial numbers in the rumen was 26.0×10^8 /g of contents, with values ranging from 2×10^8 /g to 93×10^8 /g. Cellulolytic numbers averaged 6.0×10^8 /g with a range of 1.5×10^8 /g to 24.0×10^8 /g. No differences related to sex or diet were found. In contrast, total bacterial numbers in the cecum differed between diets ($P < 0.02$), i.e., $1,046 \times 10^6$ bacteria per g for animals fed the high-forage diet compared with 166×10^6 /g for those fed the high-concentrate diet. Cellulolytic bacterial counts in the cecal contents averaged 3.1 and 7.0% of the total counts for the high-forage and high-concentrate diets, respectively. Low concentrations of fungi were found in both ruminal and cecal contents of some, but not all, animals. Unexpectedly, concentrations of bacteria and fungi in the rumen and cecum were highly correlated with their total numbers (concentration multiplied by total weight of contents).

The blue duiker (*Cephalophus monticola*), which weighs 4 to 6 kg at maturity, is probably one of the smallest ruminants in the world. It is native to central and southern Africa and appears to feed on grasses, immature leaves, and fruit. A small number of these animals was brought to The Penn State University in 1982; however, enough animals for this type of survey were not available until recently. The present study reports bacterial and fungal numbers in the ruminal and cecal contents of duikers fed a high-roughage or high-concentrate diet.

MATERIALS AND METHODS

Animals. Twenty blue duikers (*C. monticola*), 10 males and 10 females, were used in this study. Four animals were brought to the autopsy room each morning from the housing facility. A veterinarian administered a tranquilizer; and the duikers were killed by lethal injection. Ruminal and cecal contents were collected upon autopsy.

Diets. Two pelleted diets, a high-roughage diet (57.2% neutral detergent fiber) and a high-concentrate diet (26.8% neutral detergent fiber), were each fed to 10 blue duikers, five males and five females. The diets were administered at 90% of ad libitum intake for at least two months prior to killing of the animals. The compositions of the diets are given in Table 1.

Enumeration of bacteria. Total and cellulolytic bacterial numbers were estimated simultaneously in the same medium by using the most-probable-number (MPN) procedure described by Dehority et al. (7). Ruminal and cecal contents were diluted as described previously (5, 10). After 14 days of incubation, cellulolytic bacterial numbers were estimated by visual loss of cellulose and total bacterial numbers were estimated on the basis of a decrease in medium pH.

Enumeration of fungi. Both total and cellulolytic fungal

numbers in ruminal and cecal contents were also determined with an MPN procedure. The procedure has been described by Obispo (20) and is similar to the MPN procedure of Dehority et al. (7). The composition of the basal medium (7) was changed by adding 0.2% Trypticase–0.05% yeast extract–0.45 ml of VFA mixture (3) per 100 ml of medium and decreasing the amount of ruminal fluid to 20%. A filter-sterilized antibiotic solution was added before use to inhibit bacterial growth. Final antibiotic concentrations in the medium were 2,000 U of penicillin per ml and 130 U of streptomycin per ml.

Statistical analysis. Data were analyzed by Harvey least-squares analysis of variance (11).

RESULTS

The total and cellulolytic bacterial numbers in the ruminal contents of blue duikers are presented in Table 2. Since neither concentration, total numbers, nor the percentage of the population which was cellulolytic was affected by diet or sex, the data were compiled as overall means. Table 3 presents the same parameters for cecal contents. Both concentration and total numbers of bacteria were significantly higher ($P < 0.02$) in cecal contents of duikers fed the high-forage diet.

Fungal numbers in ruminal contents were not affected by diet or sex (Table 4). However, the percentage of fungi which were cellulolytic was significantly higher ($P < 0.05$) in males than in females. No cellulolytic fungi were found in female duikers.

Cecal contents contained very low numbers of fungi, averaging less than 10% of the concentration found in the rumina (Table 5). No significant differences were associated with diet or sex. It should be pointed out that only one animal, a male fed the high-concentrate diet, harbored any cellulolytic fungi in its cecum.

The sporadic occurrence and low numbers of fungi in ruminal and cecal contents raised a question about the adequacy of the medium. After the fungal assays had been

* Corresponding author.

† Paper 260-90 from the Ohio Agricultural Research and Development Center, The Ohio State University.

TABLE 1. Compositions of pelleted diets fed to blue duikers

Ingredient	% of diet ^a	
	High concentrate	High forage
Alfalfa meal	25	50
Ground oats	20	5
Ground corn	18.25	2.25
Wheat middlings	15	25
Soybean meal	7.75	6.75
Molasses	7.5	4.0
Distiller's grains with solubles		2.5
Meat-poultry byproduct	5	
Salt	0.3	0.3
Sodium bicarbonate	0.8	
Vitamin premix ^b	0.39	0.8
Limestone		0.25
Lignin binder		3.0

^a Dry matter basis.

^b Vitamin E-20000, 0.15%; trace minerals, 0.1%; Se, 0.05%; vitamin A-30000, 0.03%; vitamin D₃, 6M, 0.01%; Tylan 10, 0.05%.

incubated for at least 10 days, six MPN tubes from the highest dilution (10^{-6}), one from each of six animals, were inoculated with a 10^{-2} dilution of cattle ruminal contents. All tubes showed marked fungal growth after several days, indicating that the medium was satisfactory.

The relationship between total numbers of bacteria which occur in the rumen and cecum and their concentration per gram of contents are shown in Fig. 1. Correlation coefficients for these data were 0.93 and 0.97 for the rumen and cecum, respectively. Similar correlations were calculated with the fungal data, resulting in coefficients of 0.96 for the rumen and 0.98 for the cecum. The weight of ruminal or cecal contents was poorly correlated with bacterial concentrations ($r = 0.15$ for the rumen and $r = -0.34$ for the cecum).

DISCUSSION

In a recent study, we reported ruminal bacterial concentrations from domestic cattle and sheep determined by using these same MPN procedures (7). On the basis of data from 12 animals (8 calves and 4 sheep), total and cellulolytic bacterial numbers per gram of ruminal contents averaged 15.12×10^9 and 7.46×10^8 , respectively, for animals fed either a high-roughage or high-concentrate diet. The mean concentration of total ruminal bacteria was considerably lower in the blue duikers, whereas the concentration of cellulolytic organisms was much closer to the numbers observed in domestic ruminants.

Cowan et al. (4) compared the nutrient digestibilities of

TABLE 2. Total and cellulolytic bacterial numbers in ruminal contents of blue duikers

Value	Concn ($10^8/g$)		No. ^a (10^{10})		% Cellulolytic bacteria
	Total	Cellulolytic	Total	Cellulolytic	
Mean ^b	26.0	6.0	68.2	16.5	33.1
Range	2-93	1.5-24	4-312	1-55	6-100
SE	5.3	1.2	15.8	3.5	6.3

^a Concentration multiplied by total weight of ruminal contents.

^b Twenty animals, 10 males and 10 females, were used. Five animals of each sex were fed a high-roughage diet, and five received a high-concentrate diet.

TABLE 3. Total and cellulolytic bacterial numbers in cecal contents of blue duikers

Diet and value	Concn ($10^6/g$)		No. ^a (10^8)		% Cellulolytic bacteria
	Total	Cellulolytic ^b	Total	Cellulolytic	
High forage					
Mean ^c	1,046.3 ^d	25.2	238.1 ^d	5.1	3.1
Range	93-2,400	0-240	30-634	0-48	0-26.7
High concentrate					
Mean ^c	165.6 ^e	1.7	37.5 ^e	0.5	7.0
Range	7-430	0-9	3-111	0-3	0-58.1
SE	232.2	17.0	50.4	3.4	4.4

^a Concentration multiplied by total weight of cecal contents.

^b Cellulolytic bacteria were not detected in four animals, one male and one female, fed each diet.

^c Ten animals, five males and five females, were used.

^{d,e} Means in the same column followed by different superscript letters differ significantly ($P < 0.02$).

rabbit pellets, alfalfa hay, and fresh frozen alfalfa by blue duikers and sheep. They found that dry matter and neutral detergent fiber digestibilities were significantly lower ($P < 0.05$) in duikers; however, acid detergent fiber digestibilities were similar for the rabbit pellet and alfalfa hay diets. The bacterial concentrations observed in the present study appear to reflect these digestibilities, i.e., lower concentrations of total bacteria and similar concentrations of cellulolytic organisms.

Concentrations of viable bacteria in the ruminal contents of both domestic and wild ruminants vary considerably, ranging from about $3 \times 10^7/g$ to $300 \times 10^9/g$ for animals fed a wide range of diets (8, 14, 17, 21, 22, 24, 26). Cellulolytic numbers are much lower, ranging from $4 \times 10^5/g$ to $1.1 \times 10^9/g$. The present data fall within these reported ranges. Although blue duikers have not been studied previously, Giesecke and van Gylswyk (8) have reported viable counts of ruminal bacteria from wild impalas and springboks, both of which are in the same subfamily as blue duikers (*Antilopinae*). Total and cellulolytic counts per gram of ruminal contents were $23 \times 10^8/g$ and $0.94 \times 10^8/g$ for impalas and $55 \times 10^8/g$ and $1.1 \times 10^8/g$ for springboks. Although total bacterial concentrations in blue duikers are fairly similar, cellulolytic bacterial numbers are considerably higher. This may be a reflection of the higher quality of the fiber in the

TABLE 4. Total and cellulolytic fungal numbers in ruminal contents of blue duikers

Value	Concn ($10^2/g$)		No. ^a (10^4)		% Cellulolytic fungi ^b	
	Total ^c	Cellulolytic ^d	Total	Cellulolytic	Male	Female
Mean ^e	32.5	25.1	53.2	33.9	33.5	0
Range	0-350	0-350	0-481	0-481	0-100	0
SE	17.7	17.2	26.4	23.5	10.7	10.7

^a Concentration multiplied by total weight of ruminal contents.

^b The difference ($P < 0.05$) between males and females was significant. There were 10 animals in each group. Five animals of each sex were fed a high-roughage diet, and five received a high-concentrate diet.

^c Fungi were detected in six males and four females.

^d Cellulolytic fungi were present in only four males.

^e The data are based on 20 animals, 10 males and 10 females, except for those under % cellulolytic fungi.

TABLE 5. Total and cellulolytic fungal numbers in cecal contents of blue duikers

Value	Concn (per g)		No. ^a (10 ²)		% Cellulolytic fungi
	Total ^b	Cellulolytic ^c	Total	Cellulolytic	
Mean ^d	230	5.5	61.2	10.6	1.9
Range	0-930	0-110	0-274	0-13	0-38
SE	60.4	5.5	17.0	0.6	1.9

^a Concentration multiplied by total weight of cecal contents.

^b Fungi were detected in nine males and nine females.

^c Only one animal, a male, contained cellulolytic fungi, at a concentration of 110/g.

^d Twenty animals, 10 males and 10 females, were used. Five animals of each sex were fed a high-roughage diet, and five received a high-concentrate diet.

diets fed to the duikers compared with that of animals in the wild.

Information on cecal bacteria in ruminants is limited. Four studies have reported concentrations of total viable bacteria ranging from $4.3 \times 10^5/g$ to $15.8 \times 10^8/g$ (13, 16, 19, 25). Cellulolytic numbers, reported in only two studies, varied between 6×10^6 and 1×10^8 (13, 16). Lewis and Dehority (15) measured viable total and cellulolytic bacterial concentrations in sheep fed 100% orchardgrass hay, 60% cracked corn-40% orchardgrass hay, or 80% cracked corn-20% orchardgrass hay. The concentrations were $6.3 \times 10^8/g$, $497.4 \times 10^8/g$, and $496.0 \times 10^8/g$ of cecal contents for total viable bacteria and $1.2 \times 10^6/g$, $0.6 \times 10^6/g$, and $0.4 \times 10^6/g$ for cellulolytic bacteria, respectively. The increase in total

bacterial numbers resulting from addition of corn to the diet, either 60 or 80%, was significant ($P < 0.01$). The present results for total and cellulolytic bacterial concentrations in the ceca of high-forage diet-fed duikers, $10.5 \times 10^8/g$ and $25 \times 10^6/g$, are within what appears to be the normal ranges for forage-fed ruminants. However, total numbers were decreased by 85% in the high-concentrate diet-fed duikers compared with a 78-fold increase from feeding of concentrates in the study of Lewis and Dehority (15). Although the concentration of cellulolytic organisms was higher in the duikers, it followed a similar pattern by decreasing when concentrates were added to the diet. In both studies, animals were fed at 90% of ad libitum intake.

In ruminants, feeding of concentrates (15) or soluble carbohydrates (16) slightly lowered the cecal pH, but all reported values are between pHs 6 and 7. One might speculate that in the study by Lewis and Dehority (15), the increased concentration of bacteria in cecal contents which resulted from feeding of corn was probably caused by the availability of more energy in the cecum. Unfortunately, they did not report ruminal pH values. In the present study, the mean ruminal and cecal pH values were essentially the same for both diets, i.e., 6.2 and 8.0, respectively. These data offer little insight into the significant decrease in cecal bacterial concentrations caused by feeding with concentrates.

Fungal concentrations in ruminal contents from the duikers were not affected by diet; this appears to agree with previous studies of cattle and sheep. Reported values range from 64/g or ml to 44,000/g or ml (1, 9, 12, 23), which is fairly similar to the range of 0 to 35,000/g in the present study. Akin et al. (1) reported cellulolytic fungal concentrations ranging from 36/g to 7,360/g, constituting 5 to 93% of the total population. Only four duikers, all males, contained cellulolytic fungi, with concentrations of 90/g, 150/g, 15,000/g, and 35,000/g, which constituted 100, 35, 100, and 100%, respectively, of the total fungi. The animal containing 90 fungi per g of ruminal contents was fed the high-roughage diet, and the other three received the high-concentrate diet. The significant difference between sexes in percent cellulolytic organisms appears to be due to chance.

Fungi were detected in cecal contents of 18 duikers, nine males and nine females, with concentrations ranging from 40/g to 930/g. These values are similar to those reported by Grenet et al. (9) for cows fed a variety of diets, i.e., 70/g to 1,933/g. Only one animal, a male, contained cellulolytic fungi, and they constituted 38% of the total fungi. It is interesting that fungi occurred in the cecal contents of nine animals that were apparently devoid of ruminal fungi (less than 4/g). One of the animals without cecal fungi did contain 15,000 fungi per g of ruminal contents.

Bauchop (2) and Milne et al. (18) examined ruminal contents and feces from a large number of herbivores for the presence of anaerobic fungi. Two species, impalas and blackbucks, were from the same subfamily as duikers, *Antilopinae*. Fungi were found in the ruminal contents of impalas but were not detected in blackbuck feces.

The high correlations between total numbers of bacteria and fungi and their concentrations in contents of ingesta were unexpected (Fig. 1). Although little information is available, it has been suggested that weight of contents is probably the major factor contributing to total microbial numbers. For example, a significant difference in protozoal concentrations between animals fed low- and high-roughage diets was found; however, when total protozoal numbers were calculated, the differences were no longer significant




FIG. 1. Regression of total bacterial numbers in rumina and ceca (concentration times weight of contents) upon bacterial concentrations.

(6). Obviously, concentrations and total numbers are not independent of each other; however, the low correlation observed between concentration and content weight indicates that for these data, concentration is the major factor determining total numbers.

ACKNOWLEDGMENTS

We thank Pat Tirabasso for technical assistance in both preparation of media and inoculation of bacterial and fungal assays. Thanks also to A. L. Hattel, R. Smith, Terry Whitsel, Che-Ming Yang, R. Mothersbaugh, and Sandra Dehority for help with collection of ruminal and cecal samples.

Salaries and research support for this work were provided by State and Federal funds appropriated to the Ohio Agricultural Research and Development Center, The Ohio State University, and the Pennsylvania Agricultural Experiment Station. Travel support for this project was supplied by a University Small Research Grant from the Office of Research and Graduate Studies, The Ohio State University.

REFERENCES

- Akin, D. E., W. S. Borneman, and W. R. Windham. 1988. Rumen fungi: morphological types from Georgia cattle and the attack on forage cell walls. *Biosystems* **21**:385-391.
- Bauchop, T. 1981. The anaerobic fungi in rumen fibre digestion. *Agric. Environ.* **6**:339-348.
- Caldwell, D. R., and M. P. Bryant. 1966. Medium without rumen fluid for nonselective enumeration and isolation of rumen bacteria. *Appl. Microbiol.* **14**:794-801.
- Cowan, R. L., M. H. Michalec, G. A. Varga, and P. J. Wangness. 1986. Comparison of nutrient digestibility by the blue duiker (*Cephalophus monticola*) and sheep. *J. Anim. Sci.* **63**(Suppl. 1):419.
- Dehority, B. A. 1969. Pectin-fermenting bacteria isolated from the bovine rumen. *J. Bacteriol.* **99**:189-196.
- Dehority, B. A. 1978. Specificity of rumen ciliate protozoa in cattle and sheep. *J. Protozool.* **25**:509-513.
- Dehority, B. A., P. A. Tirabasso, and A. P. Grifo, Jr. 1989. Most-probable-number procedures for enumerating ruminal bacteria, including the simultaneous estimation of total and cellulolytic numbers in one medium. *Appl. Environ. Microbiol.* **55**:2789-2792.
- Giesecke, D., and N. O. van Gylswyk. 1975. A study of feeding types and certain rumen functions in six species of South African wild ruminants. *J. Agric. Sci.* **85**:75-83.
- Grenet, E., G. Fonty, J. Jamot, and F. Bonnemoy. 1989. Influence of diet and monensin on development of anaerobic fungi in the rumen, cecum, and feces of cows. *Appl. Environ. Microbiol.* **55**:2360-2364.
- Grubb, J. A., and B. A. Dehority. 1976. Variation in colony counts of total viable anaerobic rumen bacteria as influenced by media and cultural methods. *Appl. Environ. Microbiol.* **31**:262-267.
- Harvey, W. R. 1982. Mixed model capabilities of LSML 76. *J. Anim. Sci.* **54**:1279-1285.
- Joblin, K. N. 1981. Isolation, enumeration, and maintenance of rumen anaerobic fungi in roll tubes. *Appl. Environ. Microbiol.* **42**:1119-1122.
- Kern, D. L., L. L. Slyter, E. C. Leffel, J. M. Weaver, and R. R. Oltjen. 1974. Ponies vs steers: microbial and chemical characteristics of intestinal ingesta. *J. Anim. Sci.* **38**:559-564.
- Leedle, J. A. Z., M. P. Bryant, and R. B. Hespell. 1982. Diurnal variations in bacterial numbers and fluid parameters in ruminal contents of animals fed low- or high-forage diets. *Appl. Environ. Microbiol.* **44**:402-412.
- Lewis, S. M., and B. A. Dehority. 1985. Microbiology and ration digestibility in the hindgut of the ovine. *Appl. Environ. Microbiol.* **50**:356-363.
- Mann, S. O., and E. R. Orskov. 1973. The effect of rumen and post-rumen feeding of carbohydrates on the caecal microflora of sheep. *J. Appl. Bacteriol.* **36**:475-484.
- McBee, R. H., J. L. Johnson, and M. P. Bryant. 1969. Ruminal microorganisms from elk. *J. Wildl. Manage.* **33**:181-186.
- Milne, A., M. K. Theodorou, M. G. C. Jordan, C. King-Spooner, and A. P. J. Trinci. 1989. Survival of anaerobic fungi in feces, in saliva, and in pure culture. *Exp. Mycol.* **13**:27-37.
- Nicoletti, J. M., C. L. Davis, R. B. Hespell, and J. A. Z. Leedle. 1983. Enumeration and presumptive identification of bacteria from the small intestine of sheep. *J. Dairy Sci.* **67**:1227-1235.
- Obispo, N. E. 1990. M.S. thesis. The Ohio State University, Columbus.
- Olumeyan, D. B., T. G. Nagaraja, G. W. Miller, R. A. Frey, and J. E. Boyer. 1986. Rumen microbial changes in cattle fed diets with or without salinomycin. *Appl. Environ. Microbiol.* **51**:340-345.
- Orpin, C. G., S. D. Mathiesen, Y. Greenwood, and A. S. Blix. 1985. Seasonal changes in the ruminal microflora of the high-arctic Svalbard reindeer (*Rangifer tarandus platyrhynchus*). *Appl. Environ. Microbiol.* **50**:144-151.
- Theodorou, M. K., M. Gill, C. King-Spooner, and D. E. Beaver. 1990. Enumeration of anaerobic chytridiomycetes as thallus-forming units: novel method for quantification of fibrolytic fungal populations from the digestive tract ecosystem. *Appl. Environ. Microbiol.* **56**:1073-1078.
- Thorley, C. M., M. E. Sharpe, and M. P. Bryant. 1968. Modification of the rumen bacterial flora by feeding cattle ground and pelleted roughage as determined with culture media with and without rumen fluid. *J. Dairy Sci.* **51**:1811-1816.
- Ulyatt, M. J., D. W. Dellow, C. S. W. Reid, and T. Bauchop. 1975. Structure and function of the large intestine of ruminants, p. 119-133. *In* I. W. McDonald and A. C. I. Warner (ed.), *Digestion and metabolism in the ruminant*. University of New England Publishing Unit, Armidale, New South Wales, Australia.
- van Gylswyk, N. O. 1970. A comparison of two techniques for counting cellulolytic rumen bacteria. *J. Gen. Microbiol.* **60**:191-197.

ERRATA

Bacterial and Fungal Numbers in Ruminant and Cecal Contents of the Blue Duiker (*Cephalophus monticola*)

BURK A. DEHORITY AND GABRIELLA A. VARGA

Department of Animal Science, Ohio Agricultural Research and Development Center, The Ohio State University, Wooster, Ohio 44691, and Department of Dairy and Animal Science, The Pennsylvania State University, University Park, Pennsylvania 16802

Volume 57, no. 2, p. 469, abstract, line 7: "high-forage diet" should read "high-concentrate diet"; "high-concentrate diet" should read "high-forage diet."

Lines 8 and 9: "high-forage and high-concentrate diets" should read "high-concentrate and high-forage diets."

Results, line 9: "high-forage diet" should read "high-concentrate diet."

Line 19: "high-concentrate diet" should read "high-forage diet."

Page 470, Table 3, column 1: "High forage" should read "High concentrate"; "High concentrate" should read "High forage."

Page 471, column 2, lines 38 and 39: "high-roughage diet" should read "high-concentrate diet"; "high-concentrate diet" should read "high-roughage diet."

Monohydroxylation of Phenol and 2,5-Dichlorophenol by Toluene Dioxygenase in *Pseudomonas putida* F1

J. C. SPAIN, G. J. ZYLSTRA, C. K. BLAKE, AND D. T. GIBSON

Air Force Engineering and Services Laboratory, Tyndall Air Force Base, Florida 32403, and Department of Microbiology, University of Iowa, Iowa City, Iowa 52242

Volume 55, no. 10, p. 2650, Table 2, boxhead over columns 2 to 4: "μmol" should read "nmol."