

NOTES

Assay for β -Glucuronidase in Species of the Genus *Escherichia* and Its Applications for Drinking-Water Analysis

EUGENE W. RICE,^{1*} MARTIN J. ALLEN,² DON J. BRENNER,³ AND STEPHEN C. EDBERG⁴

*U.S. Environmental Protection Agency, Cincinnati, Ohio 45268*¹; *American Water Works Association Research Foundation, Denver, Colorado 80235*²; *Centers for Disease Control, Atlanta, Georgia 30333*³; and *Yale University School of Medicine, New Haven, Connecticut 06510*⁴

Received 14 August 1990/Accepted 14 November 1990

Recently, *Escherichia* species other than *Escherichia coli* have been isolated from potable water. Environmental isolates as well as clinical isolates of *E. adecarboxylata*, *E. blattae*, *E. fergusonii*, *E. hermannii*, and *E. vulneris* were assayed for the enzyme β -glucuronidase by using EC MUG medium and the Colilert system. None of the isolates were positive for the enzyme by either method.

Since the beginning of this century *Escherichia coli* has been considered a specific bacterial indicator of fecal pollution of water and food. The U.S. Environmental Protection Agency recently issued major changes in the Safe Drinking Water Act Total Coliform Rule that places *E. coli* as a critical indicator of fecal pollution of drinking water (8). The total coliform group will remain the primary bacterial indicator. However, for each positive total coliform test, a fecal coliform or *E. coli* analysis must be performed for that sample. The U.S. Environmental Protection Agency is basing *E. coli* methods on β -glucuronidase (GUR) activity (9).

Kilian and Bulow (12) first reported the association of GUR with the identification of *E. coli*. This enzyme is limited to the genera *Escherichia* and *Shigella* and approximately 10% of *Salmonella* species. The association between GUR and *E. coli* has been used to develop a means of identifying this species from a variety of environmental and clinical sources (4, 5, 10, 13, 14).

During the past several years, a variety of coliform bacterial isolates from potable water have been submitted to our laboratories for definitive identification. These isolates were most often forwarded because they were positive for β -galactosidase activity, as measured by the *o*-nitrophenyl- β -D-galactopyranoside (ONPG) test, but were not readily identified as members of the coliform group. When identified by conventional biochemical means (6), approximately 50% were *Citrobacter diversus* and other indole-positive total coliforms, whereas the remaining isolates consisted of *Escherichia* species other than *E. coli*.

Escherichia species including *E. adecarboxylata*, *E. blattae*, *E. fergusonii*, *E. hermannii*, and *E. vulneris* have been isolated from a variety of clinical and environmental sources (2, 3, 6, 11, 14). The sources of the clinical isolates were predominantly infections of peripheral limbs secondary to trauma. These species are not routinely isolated from stool specimens. Their presence is not considered to necessarily be indicative of fecal contamination. In drinking-water analysis, it is therefore essential to differentiate these species from *E. coli*. Previous reports, based upon a relatively small

number of isolates, suggested that *Escherichia* species other than *E. coli* are negative in assays for GUR (10, 14). To determine whether negative reactions for GUR assays were specific for these species, we analyzed clinical and environmental isolates for GUR activity.

Forty-two *Escherichia* isolates from the Enteric Diseases Branch culture collection at the Centers for Disease Control were analyzed. When first received, the majority of these isolates did not yield a definitive bacterial identification and were designated as enteric groups 1, 10, and 11. Subsequently, on the basis of DNA relatedness studies, they were designated as separate species in the genus *Escherichia* (2, 3, 7). Once the species were established by genetic means, phenotypic biochemical tests were chosen to provide optimum separation among these *Escherichia* species.

Isolates from water supply systems were also included in the study (Table 1). Standard bacteriologic analyses of the water samples from which the isolates were recovered revealed no apparent recent fecal contamination, since *E. coli* was absent. All isolates were identified by conventional biochemical tests (6).

GUR assays were performed by using the 4-methylumbelliferyl- β -D-glucuronide (MUG) substrate. MUG at a concentration of 75 mg · liter⁻¹ in a mixture of salts, cofactors, and dispersant was used in the Colilert system (Access Analytical Systems, Branford, Conn.) (4, 5). EC medium containing MUG at 50 mg · liter⁻¹ was used in the fecal coliform test (10). All strains were streaked for isolation on plate count agar and incubated at 35°C for 24 h. An isolated colony was lightly touched with a sterile loop or wooden applicator stick and used as the inoculum for both procedures. Colilert tubes were hydrated with 10 ml of sterile distilled water and, after inoculation, were incubated at 35°C for 24 h. Lauryl tryptose broth (LTB) tubes (1) were inoculated and then incubated at 35°C for 24 to 48 h. After growth was observed, each culture was transferred to EC-MUG medium and incubated at 44.5°C for 24 h. Fluorescence was observed in both procedures by exposing the tubes to long-wave UV light (366 nm). Any fluorescence was considered positive. A recent environmental isolate of *E. coli* served as a positive control and routinely yielded a strong fluorescence. A *Klebsiella pneu-*

* Corresponding author.

TABLE 1. Distribution of *Escherichia* spp. isolated from drinking water

Species	No. of isolates	Location	Sample type
<i>E. hermannii</i>	1	Florida	New pipe
	1	Alberta	Hydrant, flushing
	3	Massachusetts	Distribution system
	2	Kentucky	Distribution system, dead-end location
	1	California	Holding cistern
	1	Washington	Holding tank
	1	Arizona	Distribution system, dead-end location
	1	Georgia	Holding reservoir
	3	Illinois	Holding reservoir
	<i>E. fergusonii</i>	1	Florida
1		Alberta	Distribution system
1		California	Distribution system, dead-end location
2		Connecticut	Distribution system
1		South Carolina	Distribution system
<i>E. vulneris</i>	2	Alberta	Hydrant, flushing
	1	Florida	New pipe
	1	Louisiana	Distribution system, dead-end location
<i>E. adecarboxylata</i>	1	Massachusetts	Distribution system

moniae isolate served as a negative control and was always GUR negative.

Fecal coliform tests were also performed in accordance with standard procedures (1). An isolated colony of each strain was transferred to LTB and incubated at 35°C for up to 48 h. After growth was observed, each culture was transferred to EC medium, incubated at 44.5°C for 24 h, and observed for growth and gas production. The results of fecal coliform testing and GUR assays are presented in Table 2.

The results from this study indicate that assays for GUR by using EC-MUG medium or the Colilert system are able to distinguish *E. coli* from other *Escherichia* species. These non-*E. coli* species also do not produce a positive fecal coliform test. The Colilert system provides more rapid results and requires only one incubation temperature, as opposed to the EC-MUG and fecal coliform procedures.

TABLE 2. Results of fecal coliform and GUR tests for *Escherichia* spp.

Species	No. of isolates	No. showing fecal coliform growth	No. showing fecal coliform gas	No. positive in GUR assay with:	
				EC-MUG	Colilert-MUG
CDC ^a clones					
<i>E. hermannii</i>	17	0	0	0	0
<i>E. fergusonii</i>	11	0	0	0	0
<i>E. vulneris</i>	13	1	0	0	0
<i>E. blattae</i>	1	0	0	0	0
Environmental isolates					
<i>E. hermannii</i>	14	0	0	0	0
<i>E. fergusonii</i>	6	0	0	0	0
<i>E. vulneris</i>	4	0	0	0	0
<i>E. adecarboxylata</i>	1	0	0	0	0

^a CDC, Centers for Disease Control.

These methods are in keeping with the requirements of the new Safe Drinking Water Act Total Coliform Rule (8). The use of these procedures will permit public health authorities to make decisions regarding acute violations and possible boil-water orders based on methods which have been shown to be specific for *E. coli* detection and are not subject to false-positive results.

REFERENCES

- American Public Health Association. 1989. Standard methods for the examination of water and wastewater, 17th ed. American Public Health Association, Washington, D.C.
- Brenner, D. J., B. R. Davis, A. G. Steigerwalt, C. F. Riddle, A. C. McWhorter, S. D. Allen, J. J. Farmer, Y. Saitoh, and G. R. Fanning. 1982. Atypical biogroups of *Escherichia coli* found in clinical specimens and description of *Escherichia hermannii* sp. nov. J. Clin. Microbiol. 15:703-713.
- Brenner, D. J., A. C. McWhorter, J. K. Leete Knutson, and A. G. Steigerwalt. 1982. *Escherichia vulneris*: a new specimen of *Enterobacteriaceae* associated with human wounds. J. Clin. Microbiol. 15:1133-1140.
- Edberg, S. C., M. J. Allen, D. B. Smith, and The National Collaborative Study. 1988. National field evaluation of a defined substrate method for the simultaneous enumeration of total coliforms and *Escherichia coli* from drinking water: comparison with the standard multiple-tube fermentation method. Appl. Environ. Microbiol. 54:1595-1601.
- Edberg, S. C., M. J. Allen, D. B. Smith, and The National Collaborative Study. 1989. National field evaluation of a defined substrate method for the simultaneous detection of total coliforms and *Escherichia coli* from drinking water: comparison with presence-absence techniques. Appl. Environ. Microbiol. 55:1003-1008.
- Farmer, J. J., B. R. Davis, F. W. Hickman-Brenner, A. McWhorter, G. P. Huntley-Carter, M. A. Asbury, C. Riddle, H. G. Wathen-Grady, C. Elias, G. R. Fanning, A. G. Steigerwalt, C. M. O'Hara, G. K. Morris, P. B. Smith, and D. J. Brenner. 1985. Biochemical identification of new species and biographs of *Enterobacteriaceae* isolated from clinical specimens. J. Clin. Microbiol. 21:46-76.
- Farmer, J. J., G. R. Fanning, B. R. Davis, C. M. O'Hara, C. Riddle, F. W. Hickman-Brenner, M. A. Asbury, V. A. Lowery, and D. J. Brenner. 1985. *Escherichia fergusonii* and *Enterobacteraylorae*, two new species of *Enterobacteriaceae* isolated from clinical specimens. J. Clin. Microbiol. 21:77-81.
- Federal Register. 1989. National primary drinking water regulations: total coliforms (including fecal coliforms and *E. coli*). Fed. Regist. 54:27544-30002.
- Federal Register. 1990. National primary drinking water regulations: analytical techniques; coliform bacteria; proposed rule. Fed. Regist. 55:22752-22756.
- Hartman, P. A. 1989. The MUG (glucuronidase) test for *Escherichia coli* in food and water, p. 290-308. In A. Balows, R. C. Tilton, and A. Turano (ed.), Rapid methods and automation in microbiology and immunology. Tipografia Queriniana, Brescia, Italy.
- Izard, D., J. Mergaert, F. Gavini, A. Beji, K. Kersters, J. De Ley, and H. Leclerc. 1985. Separation of *Escherichia adecarboxylata* from the "*Erwinia herbicola-Enterobacter agglomerans*" complex and from the other *Enterobacteriaceae* by nucleic acid and protein electrophoretic techniques. Ann. Inst. Pasteur Microbiol. 136B:151-168.
- Kilian, M., and P. Bulow. 1976. Rapid diagnosis of *Enterobacteriaceae*. I. Detection of bacterial glycosidases. Acta Pathol. Microbiol. Scand. Sect. B 84:245-251.
- Rice, E. W., M. J. Allen, and S. C. Edberg. 1990. Efficacy of β -glucuronidase assay for identification of *Escherichia coli* by the defined-substrate technology. Appl. Environ. Microbiol. 56:1203-1205.
- Robison, B. J. 1984. Evaluation of a fluorogenic assay for detection of *Escherichia coli* in foods. Appl. Environ. Microbiol. 48:285-288.