

Chorismate Mutase and 3-Deoxy-D-arabino-Heptulosonate 7-Phosphate Synthase of the Methylophilic Actinomycete *Amycolatopsis methanolica*

G. J. W. EUVERINK, G. I. HESSELS, C. FRANKE, AND L. DIJKHUIZEN*

Department of Microbiology, Groningen Biomolecular Sciences and Biotechnology Institute (GBB),
9751 NN Haren, The Netherlands

Received 18 April 1995/Accepted 29 August 1995

Chorismate mutase (CM) and 3-deoxy-D-arabino-heptulosonate 7-phosphate (DAHP) synthase (DS) are key regulatory enzymes in L-Phe and L-Tyr biosynthesis in *Amycolatopsis methanolica*. At least two CM proteins, CMIA and CMIB, are required for the single chorismate mutase activity in the wild type. Component CMIA (a homodimeric protein with 16-kDa subunits) was purified to homogeneity (2,717-fold) and kinetically characterized. The partially purified CMIB preparation obtained also contained the single DS (DSI) activity detectable in the wild type. The activities of CMIA and CMIB were inhibited by both L-Phe and L-Tyr. DSI activity was inhibited by L-Trp, L-Phe, and L-Tyr. A leaky L-Phe-requiring auxotroph, mutant strain GH141, grown under L-Phe limitation, possessed additional DS (DSII) and CM (CMII) activities. Synthesis of both CMII and DSII was repressed by L-Phe. An *ortho*-DL-fluorophenylalanine-resistant mutant of the wild type (strain oFPHE83) that had lost the sensitivity of DSII and CMII synthesis to L-Phe repression was isolated. DSII was partially purified (a 42-kDa protein); its activity was strongly inhibited by L-Tyr. CMII was purified to homogeneity (93.6 fold) and characterized as a homodimeric protein with 16-kDa subunits, completely insensitive to feedback inhibition by L-Phe and L-Tyr. The activity of CMII was activated by CMIB; the activity of CMII plus CMIB was again inhibited by L-Phe and L-Tyr. A tightly blocked L-Phe- plus L-Tyr-requiring derivative of mutant strain GH141, GH141-19, that had lost both CMIA and CMII activities was isolated. The above-described properties, and the N-terminal amino acid sequences, showed that CMIA and CMII are one and the same protein.

The aromatic amino acids L-Phe, L-Tyr and L-Trp are synthesized via a common pathway. Erythrose-4-phosphate and phosphoenolpyruvate are condensed into 3-deoxy-D-arabino-heptulosonate 7-phosphate (DAHP) by DAHP synthase (DS; EC 4.1.2.15). DAHP is converted via the shikimate pathway into chorismate, involving six enzyme steps (14). Chorismate mutase (CM; EC 5.4.99.5) synthesizes prephenate from chorismate; this synthesis is an important and committed step in L-Phe and L-Tyr biosynthesis and is widespread in nature (4). Chorismate is also converted into anthranilate (L-Trp biosynthesis), *p*-aminobenzoate (folic acid biosynthesis), *p*-hydroxybenzoate (ubiquinone biosynthesis), or isochorismate (menaquinone and enterobactin biosynthesis) (4).

Aromatic amino acid biosynthesis is generally controlled by feedback inhibition or repression at the level of DS and CM. These proteins are present either as monofunctional or bifunctional (iso)enzymes or as part of multienzyme complexes (25).

Current knowledge on the biochemistry and regulation of the pathways of primary metabolism in actinomycetes (gram-positive soil bacteria) is limited but considered to be important for further rational improvement of strains for overproducing aromatic amino acids and derived compounds (7, 29). Many secondary metabolites synthesized by actinomycetes are derived from the aromatic amino acids themselves or from intermediates in their biosyntheses (13, 34). Examples are the antibiotics rifamycin, vancomycin, and avoparcin, which are

produced by the industrial actinomycete strains *Amycolatopsis mediterranei*, *Amycolatopsis orientalis*, and *Amycolatopsis coloradensis*, respectively (16, 22, 25, 30).

We have initiated studies on glucose, methanol, and aromatic amino acid metabolism in the related methylophilic actinomycete *Amycolatopsis methanolica* (9, 16) and have purified and characterized several enzymes of glucose and quinolate metabolism, prephenate dehydratase, and the multiple aromatic aminotransferases present in this organism (1, 3, 17, 19). Previously, we have also shown that the single DS enzyme (DSI) detectable in wild-type *A. methanolica* has the unique property of being feedback inhibited by all three aromatic amino acids (12). In this paper we report a detailed biochemical analysis of the DS and CM enzymes of *A. methanolica*.

MATERIALS AND METHODS

Microorganisms and cultivation. The *A. methanolica* wild-type strain (NCIB 11946) (9, 28), the plasmid pMEA300 (41a)-deficient strain WV2 (42), and auxotrophic mutants derived from strain WV2 (this study) were used. The procedures followed for the cultivation in batch cultures, harvesting of cells, and measurements of growth have been described previously (11). Glucose (1 M) was heat sterilized, and amino acid supplements were filter sterilized.

Mutant isolation. Mutants blocked in aromatic amino acid biosynthesis were isolated following UV irradiation treatments (17, 18).

Isolation of L-Phe analog-resistant mutants. Mutants of the wild-type strain resistant to the toxic L-Phe analog *ortho*-fluoro-DL-phenylalanine (oFPhe, 27.3 mM) were isolated on 10 mM glucose mineral agar (1.5% [wt/vol]) plates containing the filter-sterilized analog. Fifteen agar plates were inoculated with approximately 5×10^7 cells each. After 2 weeks the spontaneous oFPhe-resistant colonies that had appeared were purified on homologous media.

Preparation of extracts and enzyme assays. Washed cell suspensions were disrupted in a French pressure cell at 1,000 MPa. Unbroken cells and debris were removed by centrifugation at $40,000 \times g$ for 30 min at 4°C. Following desalting through PD 10 Pharmacia columns, the supernatant, containing 10 to 20 mg of protein \cdot ml⁻¹, was used for enzyme assays.

* Corresponding author. Mailing address: Department of Microbiology, Groningen Biomolecular Sciences and Biotechnology Institute, Kerklaan 30, 9751 NN Haren, The Netherlands. Phone: 31.50.632153. Fax: 31.50.632154. Electronic mail address: L.DIJKHUIZEN@BIOL.RUG.NL.

CM was assayed by measuring the amount of prephenate formed after conversion to phenylpyruvate (15). The reaction mixture (100 μ l) contained 50 mM Tris-HCl (pH 7.5) and 2.0 mM chorismate and extract or protein, as indicated in the individual experiments. After 10 min, 10 μ l of 4.5 M HCl was added, and the reaction mixture was incubated for 15 min at 37°C. The phenylpyruvate formed was estimated by adding 890 μ l of a 1.58 M NaOH solution and measuring the A_{320} (ϵ_{320} [phenylpyruvate] = 17.5×10^3 M \cdot cm $^{-1}$). Endpoint measurements were indicative of initial reaction rates.

DS was assayed at 37°C by measuring the amount of DAHP formed from erythrose-4-phosphate and phosphoenolpyruvate (12, 27).

Purification of CMI from strain WV2. All chromatographic steps were carried out with a System Prep 10 liquid chromatography system (Pharmacia LKB Biotechnology Inc.).

(i) **Step 1. Glucose-grown cells** (25 g [wet weight]) were harvested in the late exponential phase of growth. Extracts were prepared as described above in 25 mM Tris-HCl (pH 7.5) containing 1 mM dithiothreitol (buffer A). DNase I (grade II, from bovine pancreas) and 1 mM MgCl₂ were added to the extract, and the mixture was incubated for 10 min.

(ii) **Step 2. Hydrophobic interaction chromatography.** The extract was adjusted to 1.1 M (NH₄)₂SO₄. Precipitated proteins were removed via centrifugation (15 min at 40,000 \times g). The resulting supernatant was applied to a column of butyl-Sepharose fast-flow (1.6 by 20 cm, 4°C) equilibrated in buffer A containing 1.1 M (NH₄)₂SO₄. Bound protein was eluted with a 400-ml decreasing linear gradient from 1.1 to 0.3 M (NH₄)₂SO₄ (flow rate, 4.0 ml \cdot min $^{-1}$; fractions, 4.0 ml).

(iii) **Step 3. Gel filtration chromatography.** The protein from step 2 was concentrated by slowly adding solid (NH₄)₂SO₄ to 50% saturation. The mixture was stirred for 30 min at 4°C and centrifuged for 15 min at 40,000 \times g. The pellet was dissolved in 3 ml of buffer A and applied to a Superdex 200 (XK 16/60) gel filtration column previously equilibrated in buffer A containing 0.15 M KCl (flow rate, 1 ml \cdot min $^{-1}$; fractions, 2 ml).

(iv) **Step 4. Anion-exchange chromatography.** The protein from step 3 was dialyzed against 50 mM Tris-HCl (pH 8.8) containing 1 mM dithiothreitol (buffer B) and was applied to a Mono Q (HR 5/5) anion-exchange column. Bound protein was eluted with a 30-ml linear increasing gradient from 0 to 0.5 M KCl in buffer B (flow rate, 1 ml \cdot min $^{-1}$; fractions, 0.5 ml).

Purification of CMII from mutant strain GH141. (i) **Step 1.** Glucose-grown cells (25 g [wet weight]) were harvested in the late exponential phase of growth, and an extract was prepared in buffer A as described above. DNase I (grade II, from bovine pancreas) and 1 mM MgCl₂ were added to the extract, and the mixture was incubated for 10 min.

(ii) **Step 2. (NH₄)₂SO₄ precipitation.** Solid (NH₄)₂SO₄ was slowly added to the extract to 35% saturation. The mixture was stirred for 30 min at 4°C and centrifuged for 15 min at 40,000 \times g. The resulting supernatant was adjusted to 50% saturation by adding solid (NH₄)₂SO₄. The mixture was stirred for 30 min at 4°C and centrifuged for 15 min at 40,000 \times g. The pellet was dissolved in 8 ml of buffer B and dialyzed against the same buffer.

(iii) **Step 3. Gel filtration chromatography.** The protein from step 2 was applied to a Superdex 200 (XK 16/60) gel filtration column previously equilibrated in buffer B containing 0.15 M KCl (flow rate, 1 ml \cdot min $^{-1}$; fractions, 2 ml).

(iv) **Step 4. Anion-exchange chromatography.** The protein from step 3 was dialyzed against buffer B and was applied to a Mono Q (HR 5/5) anion-exchange column. Bound protein was eluted with a 30-ml linear increasing gradient from 0 to 0.5 M KCl in buffer B (flow rate, 1 ml \cdot min $^{-1}$; fractions, 1 ml).

(v) **Step 5. Hydrophobic interaction chromatography.** The protein from step 4 was adjusted to 1.5 M (NH₄)₂SO₄ and applied to a phenyl-Superose (HR 5/5) column. Bound protein was eluted with a 20-ml linear decreasing gradient from 1.5 to 0 M (NH₄)₂SO₄ in buffer B (flow rate, 0.5 ml \cdot min $^{-1}$; fractions, 0.5 ml).

SDS-PAGE. Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) was performed as described by Laemmli and Favre (31), with the marker proteins from the Combithek calibration protein kit (Boehringer, Mannheim, Germany). Gels were stained with Coomassie brilliant blue R250.

Estimation of molecular mass. Molecular masses were estimated by using a Superdex 200 column (XK 16/60) and Bio-Rad (Richmond, Calif.) gel filtration standards.

Kinetic studies. Kinetic parameters were determined at 37°C in 50 mM Tris-HCl (pH 7.5) and were calculated with the use of Sigmaplot for Windows 2.0 (Jandell Scientific Software) by using curve fitting with the Michaelis-Menten equation and its derivatives for various types of inhibition. Substrate and effector concentrations were as shown in the individual experiments.

Automated amino acid sequence determination. The CMIa and CMII proteins were applied to a Pro-spin cartridge (Applied Biosystems, Warrington, United Kingdom) containing a polyvinylidene difluoride membrane. Sequencing was performed on an Applied Biosystems model 475A/120A automated gas-phase sequencer equipped with on-line high-pressure liquid chromatography (HPLC) for detection of phenylthiohydantoin amino acid derivatives (Eurosequence bv, Groningen, The Netherlands).

Analytical methods. Protein concentrations were determined with the Bio-Rad protein determination kit, with bovine serum albumin as the standard (5). Glucose concentrations were determined with the GOD-period kit from Boehringer. Amino acid concentrations were determined by HPLC analysis (19).

TABLE 1. Specific activities of DS and CM in extracts of glucose-grown cells of *A. methanolica* WV2 and mutants harvested in late exponential phase of growth

Strain	Sp act (U \cdot mg of protein $^{-1}$) ^a			
	CM		DS	
	Alone	With L-Phe + L-Tyr	Alone	With L-Trp
WV2 ^b	7	1	22	2
GH141	43	42	50	30
GH141 ^c	7	1	25	3
GH141-19 ^d	0	0	22	4
oFPHE83	33	13	188	114
oFPHE83 ^c	32	11	246	162

^a CM activity was determined without effectors and in the presence of 1 mM L-Phe plus 1 mM L-Tyr. DS activity was determined without effectors and in the presence of 1 mM L-Trp.

^b CM is also inhibited by 1 mM L-Phe (59%) and 1.0 mM L-Tyr (50%) separately. DAHP synthase is inhibited not only by L-Trp (90% at 1.0 mM) but also by L-Phe (74% at 1.0 mM) and L-Tyr (60% at 1.0 mM) separately (12).

^c Glucose-grown cells supplemented with 100 mg of L-Phe liter $^{-1}$.

^d Glucose-grown cells supplemented with 100 mg of L-Phe plus 100 mg of L-Tyr liter $^{-1}$.

Biochemicals. Chorismate was obtained as a barium salt from Sigma. Before use, barium ions were removed via precipitation with excess K₂SO₄. Chorismate was further purified on a Supelcosil LC-18-DB semi-prep (5- μ m particle diameter) column (20.0 cm by 10.0 mm [inside diameter]; Supelco, Inc., Bellefonte, Pa.) (8). All other chemicals were analytical grade and commercially available.

RESULTS

CM and DS in *A. methanolica*. CM and DS activities in extracts of wild-type *A. methanolica* WV2 were 7 and 22 mU \cdot mg of protein $^{-1}$, respectively. CM was inhibited by both L-Phe and L-Tyr, and DS was inhibited by all three aromatic amino acids (Table 1). The addition of L-Phe, L-Tyr, or L-Trp (100 mg \cdot l $^{-1}$ each), separately or in various combinations, to the growth medium of *A. methanolica* WV2 had no effect on the specific activities of CM and DS and feedback inhibition patterns.

Purification of CM from wild-type strain WV2. Purification of CM from *A. methanolica* WV2 turned out to be rather difficult. In a first attempt, extract was applied to a Q-Sepharose anion-exchange column. Bound protein was eluted with a linear gradient of 0 to 1 M KCl, but no activity was found. Hydrophobic interaction chromatography on butyl-Sepharose, however, yielded a single activity peak. Also, gel filtration of extracts revealed a single CM activity peak, coeluting with a single DS activity peak, corresponding to a molecular mass of 240 kDa (Fig. 1A). In addition, dialysis against 50 mM Tris-HCl (pH 7.5 or pH 8.8) with or without 1 mM EDTA, centrifugation steps (40,000 to 100,000 \times g, 2 h), and ammonium sulfate precipitation did not inactivate CM. Small diffusible effector molecules, e.g., metal ions or metabolites, thus are not required for CM activity. We subsequently observed that after Q-Sepharose anion-exchange chromatography, CM activity could be restored by mixing the flowthrough (component CMIa) with fractions eluting from the column at approximately 0.3 M KCl (component CMIb). DS activity coeluted with component CMIb. Gel filtration of CMIa and CMIb separately, followed by reconstitution of CM activity with the other component, showed that they eluted in fractions corresponding to molecular masses of 31 and 160 kDa, respectively. In this step DS also coeluted with component CMIb (160 kDa), instead of behaving as a protein with a size of 240 kDa (see above). Gel filtration of extracts (in the presence of up to 1 M




FIG. 1. Superdex 200 gel filtration elution profiles of DS and CM in extracts of cells of *A. methanolica* WV2 (A) (21.9 mg of protein) and mutant strain GH141 (B) (20.3 mg of protein) grown in glucose mineral medium. ○, DS activity; ●, CM activity; ---, A₂₈₀.

KCl at pH 7.5) and hydrophobic interaction chromatography did not affect the interaction between CMIa and CMIb, showing that their binding is relatively strong. Gel filtration of extracts at pH 8.5 and 0.15 M KCl did result in separation of the CMIa and CMIb components.

Component CMIa, a minor protein in *A. methanolica*, was purified (2,717-fold) to homogeneity in four steps, with an overall yield of 5.2% (Table 2). The purification protocol was designed in such a way that CM became separated into components CMIa and CMIb in the last step in the purification scheme only, eluting at 0.05 and 0.35 M KCl, respectively. In this way CM activity could be monitored more conveniently, and homogeneous preparations of CMIa were obtained more easily. CM and DS coeluted during purification; in the final step, DS coeluted with CMIb. The CMIb preparation was not subjected to further purification. SDS-PAGE of CMIa revealed a single band migrating at 16 kDa. In view of its native

molecular mass of 31 kDa, CMIa appears to be a homodimeric protein with 16-kDa subunits.

The identification of N-terminal amino acids of CMIa was hampered by the low signal-to-noise ratio. The following seven N-terminal amino acids were tentatively identified: X, X, Q, X, N, E, K, L, T, and P (X, not identified). A BLASTP (2) search of the available databases revealed no significant homology with any other proteins described.

Kinetic studies were carried out with 0.3 μg of protein of pure component CMIa and 20 μg of the CMIb preparation (Fig. 2; Table 2). Under these conditions, CMIa is clearly limiting for the overall CM activity (Fig. 2 inset). The K_m for chorismate was 2.0 ± 0.3 mM (mean \pm standard deviation). The calculated V_{max} value was 47 ± 2.9 U · mg of protein⁻¹. CM activity was competitively inhibited by L-Phe and/or L-Tyr. The K_i values for L-Phe and L-Tyr were 0.66 ± 0.02 mM and 0.52 ± 0.01 mM, respectively.

Characterization of L-Phe auxotrophic mutants. A leaky L-Phe auxotrophic mutant, strain GH141, which had lost 90% of the L-Phe aminotransferase activity was isolated previously (1). Cultivation of the strain in glucose mineral medium showed that the doubling time of strain GH141 had increased to 6 h, because growth had become limited by the rate of L-Phe biosynthesis. The doubling time of mutant strain GH141 in glucose mineral medium supplemented with L-Phe (100 mg · liter⁻¹) was comparable to that of the wild type (2.5 h). Extracts of mutant strain GH141 cells grown in glucose mineral medium displayed DS and CM activities at levels about 2.5 to 6 times higher than those of the wild type. In mutant strain GH141, CM activity was not feedback inhibited by either L-Phe or L-Tyr, and the sensitivity of DS to L-Trp was strongly reduced (Table 1). Fractionation of extracts of mutant strain GH141, grown on glucose mineral medium, on a Superdex 200 gel filtration column revealed additional CM (CMIi, 31 kDa) and DS (DSII, 42 kDa) activity peaks (Fig. 1B). The CMI and DS I enzymes in the wild type and in mutant strain GH141 possessed identical molecular masses and feedback inhibition sensitivities. Cells of strain GH141 grown in glucose mineral medium supplemented with 100 mg of L-Phe liter⁻¹ displayed wild-type levels of CM and DS activities and properties (Table 1) and completely lacked the CMIi and DSII activity peaks.

Strain GH141-19, a tightly blocked L-Phe- plus L-Tyr-requiring auxotrophic mutant, subsequently was derived from strain GH141. Strain GH141-19 had completely lost CM activity (Table 1). The addition of pure component CMIa to extracts of mutant GH141-19 restored CM activity to wild-type levels. This indicates that strain GH141-19 is deficient in component CMIa specifically.

Characterization of DSII. Further attempts to purify DSII after gel filtration chromatography failed because of a significant loss of activity, which occurred during anion-exchange or hydrophobic interaction chromatography. The addition of divalent cations to buffer solutions and mixing of the fractions

TABLE 2. Purification of CMI from glucose-grown cells of *A. methanolica* WV2

Step	Vol (ml)	Protein (mg)	Total activity (U) ^a	Sp act (U · mg ⁻¹)	Purification (fold)	Yield (%)
Extract	44	1,562	8.3	0.005	1.0	100.0
Butyl-Sepharose	38	47.9	2.0	0.042	7.9	24.2
Superdex 200	8	13.0	1.2	0.091	17.0	14.2
Mono Q ^b	1	0.03	0.4	14	2,717.0	5.2

^a One unit of activity is defined as 1 μmol of prephenate formed per min from chorismate.

^b Excess CMIb was added to assay the chorismate mutase activity of CMIa (see the text).




FIG. 2. Specific CM activity of component CMIa (0.3 μg of protein) plus excess of component CMIb (20 μg of protein) with increasing chorismate concentrations in the absence of an effector (●) or in the presence of 2.0 mM L-Phe (■), 2.0 mM L-Tyr (○), and 2.0 mM L-Phe plus 2.0 mM L-Tyr (□). (Inset) CM activity with increasing concentrations of CMIa in the presence of 20 μg of protein of preparation CMIb at a chorismate concentration of 2.0 mM. PPA, prephenate.

were not successful. Some characteristics of DSII therefore were determined after gel filtration. DSII remained active for at least 24 h at 4°C. The apparent K_m values for phosphoenolpyruvate (3.8 mM erythrose-4-phosphate) and erythrose-4-phosphate (3.0 mM phosphoenolpyruvate) were 0.40 ± 0.06 mM and 1.1 ± 0.2 mM, respectively. With 3.8 mM phosphoenolpyruvate and 1.7 mM erythrose-4-phosphate, DSII activity was inhibited by 1.0 mM L-Trp (17%) and 1.0 mM L-Tyr (83%) but not by L-Phe. L-Tyr inhibition was competitive with respect to the erythrose-4-phosphate concentration, and the K_i value for L-Tyr was 0.031 ± 0.006 mM. Noncompetitive inhibition by L-Tyr was found with respect to the phosphoenolpyruvate concentration, and the K_i value was 0.8 ± 0.2 mM. Intermediates in the biosynthesis of aromatic amino acids (shikimate, anthranilate, chorismate, and prephenate; 1 mM each) inhibited DSII activity by less than 15%.

Characterization of CMII. The CMII enzyme was much more abundant than CMIa in *A. methanolica*. CMII was purified (93.6-fold) to homogeneity from cells of mutant strain GH141, with an overall yield of 25% (Table 3). CMII bound to

a Mono Q anion-exchange column at pH 8.8, but not at pH 7.5, and eluted at approximately 0.05 M KCl. SDS-PAGE of CMII revealed a single band migrating at 16 kDa. In view of its native molecular mass of 31 kDa, CMII appears to be a homodimeric protein with 16-kDa subunits.

The following 24 N-terminal amino acids were identified: M, A, Q, T, N, E, K, A, T, P, X, E, T, S, G, E, P, V, A, S, A, X, E, and I (X, not identified). A BLASTP (2) search against the available databases revealed no homology with other CM enzymes described. Six of seven amino acids, however, were identical to tentatively identified N-terminal amino acids of component CMIa (see above).

A virtually linear relationship was observed between the chorismate concentration, in the range of 0 to 4 mM, and CMII activity (Fig. 3). L-Phe and L-Tyr (1 mM concentrations) did not inhibit CMII activity or modify this linear relationship (data not shown). CMII was activated by CMIb, resulting in Michaelis-Menten kinetics (Fig. 3 inset). The K_m value for chorismate was 2.2 ± 0.2 mM, and a V_{\max} of 47 ± 2.5 U \cdot mg of protein $^{-1}$ could be calculated. The activity of CMII plus

TABLE 3. Purification of CMII from glucose-grown cells of *A. methanolic* mutant strain GH141

Step	Vol (ml)	Protein (mg)	Total activity (U) ^a	Sp act (U · mg ⁻¹)	Purification (fold)	Yield (%)
Extract	49	882.0	47.5	0.05	1.0	100.0
35–50% (NH ₄) ₂ SO ₄	8	2.5	50.3	0.13	2.5	105.9
Superdex 200	24	4.6	21.6	0.25	4.6	45.5
Mono Q	4	69.5	17.7	3.75	69.5	37.2
Phenyl-Superose	2	2.3	11.7	5.05	93.6	24.6

^a One unit of activity is defined as 1 μ mol of prephenate formed per min from chorismate.

CMIIb was inhibited by L-Phe and L-Tyr (Fig. 3), albeit less strongly than that of CMIIa plus CMIIb (Fig. 2). CMII plus CMIIb activity was competitively inhibited by L-Phe and/or L-Tyr. The K_i values for L-Phe and L-Tyr were 4.5 ± 0.4 mM and 3.8 ± 0.5 mM, respectively.

Isolation and characterization of oFPHE83. De Boer et al. (10) provided evidence that the toxic L-Phe analog oFPhe blocks growth of *A. methanolic* in glucose mineral medium via

prephenate dehydratase or CM inhibition. In the present study we observed that oFPhe-resistant mutant strains of wild-type *A. methanolic* (NCIB 11946) could be isolated readily. After 2 weeks of incubation, numerous spontaneous oFPhe-resistant colonies were clearly visible against a background of tiny colonies. Initially, a total of 400 colonies resistant to oFPhe were selected. After repeated transfers, 122 oFPhe-resistant mutants still scored clearly positive. The CM enzymes of five




FIG. 3. Specific CM activity of CMII (1.2 μ g of protein) (▲) and CMII (0.23 μ g of protein) plus excess of component CMIIb (20 μ g of protein) with increasing chorismate concentrations in the absence of an effector (●) or in the presence of 2.0 mM L-Phe (■), 2.0 mM L-Tyr (○), and 2.0 mM L-Phe plus 2.0 mM L-Tyr (□). (Inset) CM activity with increasing concentrations of CMII in the presence of 20 μ g of protein of preparation CMIIb at a chorismate concentration of 2.0 mM.




FIG. 4. Regulation of aromatic amino acid biosynthesis in *A. methanolica*. FI, feedback inhibition; FA, feedback activation; FR, feedback repression; dashed lines, multiple enzyme steps; none¹, data from De Boer et al. (12).

oPhe-resistant strains with normal growth rates in glucose mineral medium were analyzed for their sensitivities towards feedback regulation. In four strains, CM and DS displayed wild-type levels of activities and degrees of sensitivity for feedback regulation. Mutant strain oPHE83 possessed (very) high levels of CM and DS activities (Table 1). Fractionation of extracts of mutant strain oPHE83 via gel filtration revealed the additional presence of DSII and CMII enzymes (data not shown). Unlike the situation in the wild type and in mutant strain GH141, the synthesis of the DSII and CMII enzymes was not repressed by L-Phe in mutant strain oPHE83 (Table 1). The CM Ia and CM II enzymes and the DS I and DS II enzymes in the wild-type mutant strain GH141 and in strain oPHE83 possessed otherwise identical molecular masses and feedback inhibition sensitivities.

DISCUSSION

The limited studies thus far carried out with gram-positive bacteria have revealed the presence of isoenzymes of CM in bacilli only (21, 32, 36). In *Bacillus subtilis* one of these CM isoenzymes constitutes a bifunctional protein with DS (35). A CM-DS enzyme complex has also been reported for *Brevibacterium flavum* (37, 40, 41). This paper is the first report of a complex of DS and CM in an actinomycete. CM activity in wild-type *A. methanolica* requires at least two proteins (CM Ia

and CM Ib), and in several purification steps coelution of DS (DS I) and CM occurred (this study). In the final step of the purification protocol for component CM Ia, DS I coeluted with CM Ib. Also, the molecular masses of DS I (168 kDa [12]) and CM Ib (160 kDa) are similar. These data thus suggest that CM Ib and DS I are identical. Considering the molecular masses of components CM Ia and CM Ib-DS I, two CM Ia dimers (31 kDa) are most likely associated with CM Ib-DS I in a 240-kDa complex (Fig. 1A).

CM activity in *A. methanolica* is feedback inhibited by L-Phe and L-Tyr, a characteristic that is shared with the enzyme from *A. mediterranei* (43). The *Corynebacterium glutamicum* (24) and *B. flavum* (40) CM enzymes were also inhibited by L-Phe and L-Tyr; this inhibition was released in the presence of L-Trp, a characteristic not shown by *A. methanolica* CM. The spore-forming members of the order *Actinomycetales* display CM activity that is inhibited by L-Tyr and/or L-Trp only (26, 38). The CM of *Streptomyces aureofaciens* contains at least three subunits with sizes of 14 kDa; the enzyme is insensitive to inhibition by aromatic amino acids (20). Also, the 75-kDa CM detected in extracts of *Streptomyces* sp. strain 3022a, a chloramphenicol producer, was not inhibited by aromatic amino acids (33).

Characterization of the leaky L-Phe aminotransferase mutant strain GH141 (1) revealed that synthesis of a second CM enzyme (CM II) and a second DS enzyme (DS II) became de-

repressed under L-Phe-limiting conditions. Following growth in the presence of L-Phe, only the constitutive DSI with L-Trp as the main negative effector and CMI with L-Phe and L-Tyr as the main negative effectors were present (Table 1). The analog-resistant mutant strain oFPHE83 most likely has lost L-Phe feedback repression sensitivity, resulting in enhanced DS and CM activities due to DSII and CMII derepression. L-Phe repression of CM (but not of DS) synthesis was also reported in *C. glutamicum* (23, 24). In *B. flavum*, both CM and DS syntheses are repressed by L-Tyr (37).

DSII is a 42-kDa protein that is strongly feedback inhibited by L-Tyr and less strongly inhibited by L-Trp. De Boer et al. (12) reported that the 41-kDa subunits of the tetrameric DSI protein from *A. methanolicus* may also display DS activity. It appears unlikely, however, that subunits of DSI are responsible for DSII activity: L-Tyr strongly inhibits DSII but not the DSI subunits (Table 1). Thus, *A. methanolicus* most likely possesses DSI and DSII isoenzymes. This also may explain why an extensive search for auxotrophic mutants of *A. methanolicus* yielded approximately 150 aromatic amino acid auxotrophs but not DS-negative mutants (18).

The low-molecular-mass DS protein (DSII) detected in *A. methanolicus* is unique in prokaryotes and has been reported for the yeast *Saccharomyces cerevisiae* (39- and 41-kDa proteins) only (6). The only other actinomycete DS that has been described thus far is a relatively large, oligomeric protein of *Streptomyces rimosus* (39).

The respective native and subunit molecular masses of CMIa and CMII are equal; also the K_m values for chorismate and the V_{max} values for CMIa plus CMIIb and CMII plus CMIIb are similar. These properties, and the N-terminal amino acid sequences, suggest that CMIa and CMII are identical proteins. This was confirmed by the isolation of the CM-deficient mutant strain GH141-19. A striking difference between CMIa and CMII, the activity of CMII alone, can be explained by the much higher concentrations of CMII present in the extracts. When CMIa and CMII were used at similar low concentrations, no activity was detectable; the addition of CMIIb restored activity in both cases. The differences in feedback inhibition sensitivity between the CM activity of CMIa plus CMIIb and CMII plus CMIIb for L-Phe and/or L-Tyr remain to be explained, however.

The N-terminal amino acids identified for the CMIa and CMII proteins did not show any significant similarity with other known proteins, suggesting that the *A. methanolicus* enzyme may belong to a new class of CM enzymes. Cloning of the gene encoding CMIa-CMII is currently in progress. In future work we will study the unique regulatory properties of this actinomycete CM in more detail.

In conclusion, L-Phe and L-Tyr biosynthesis in wild-type *A. methanolicus* is regulated via (i) L-Phe inhibition and L-Tyr activation of prephenate dehydratase activity, (ii) L-Phe and L-Tyr inhibition of CM activity, and (iii) repression of CMII-CMIa and DSII synthesis by L-Phe (Fig. 4). Under L-Phe-limiting conditions, derepression of both CMII-CMIa and DSII synthesis increases the flux of intermediates toward L-Phe and L-Tyr biosynthesis. L-Tyr inhibition of DSII will prevent the accumulation of L-Tyr under L-Phe limitation (Fig. 4). The relative in vivo contribution of these control mechanisms remains to be determined.

ACKNOWLEDGMENTS

This work was supported by grant no. GBI81.1510 from the Netherlands Technology Foundation (STW), which is subsidized by the Netherlands Organization for the Advancement of Pure Research (NWO).

REFERENCES

- Abou-Zeid, A. M., G. J. W. Euverink, G. I. Hessels, and L. Dijkhuizen. 1995. Biosynthesis of L-phenylalanine and L-tyrosine in the actinomycete *Amycolatopsis methanolicus*. *Appl. Environ. Microbiol.* **61**:1298–1302.
- Altschul, S. F., G. Warren, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment search tool. *J. Mol. Biol.* **215**:403–410.
- Alves, A. M. C. R., G. J. W. Euverink, H. J. Hektor, G. I. Hessels, J. Van der Vlag, J. W. Vrijbloed, D. H. A. Hondmann, J. Visser, and L. Dijkhuizen. 1994. Enzymes of glucose and methanol metabolism in the actinomycete *Amycolatopsis methanolicus*. *J. Bacteriol.* **176**:6827–6835.
- Bentley, R. 1990. The shikimate pathway—a metabolic tree with many branches. *Crit. Rev. Biochem. Mol. Biol.* **25**:307–384.
- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**:248–254.
- Braus, G. H. 1991. Aromatic amino acid biosynthesis in the yeast *Saccharomyces cerevisiae*—a model system for the regulation of a eukaryotic biosynthetic pathway. *Microbiol. Rev.* **55**:349–370.
- Bushell, M. E. 1989. The process physiology of secondary metabolite production, p. 95–120. In S. Baumberg, I. S. Hunter, and P. M. Rhodes (ed.), *Microbial products: new approaches*. Cambridge University Press, Cambridge.
- Connelly, J. A., and D. L. Siehl. 1987. Purification of chorismate, prephenate, and arogenate by HPLC. *Methods Enzymol.* **142**:422–431.
- De Boer, L., L. Dijkhuizen, G. Grobden, M. Goodfellow, E. Stackebrandt, J. H. Parlett, D. Whitehead, and D. Witt. 1990. *Amycolatopsis methanolicus* sp. nov., a facultatively methylotrophic actinomycete. *Int. J. Syst. Bacteriol.* **40**:194–204.
- De Boer, L., G. Grobden, J. W. Vrijbloed, and L. Dijkhuizen. 1990. Biosynthesis of aromatic amino acids in *Nocardia* sp. 239: effects of amino acid analogues on growth and regulatory enzymes. *Appl. Microbiol. Biotechnol.* **33**:183–189.
- De Boer, L., W. Harder, and L. Dijkhuizen. 1988. Phenylalanine and tyrosine metabolism in the facultative methylotroph *Nocardia* sp. 239. *Arch. Microbiol.* **149**:459–465.
- De Boer, L., J. W. Vrijbloed, G. Grobden, and L. Dijkhuizen. 1989. Regulation of aromatic amino acid biosynthesis in the ribulose monophosphate cycle methylotroph *Nocardia* sp. 239. *Arch. Microbiol.* **151**:319–325.
- Demain, A. L. 1983. New applications of microbial products. *Science* **219**:709–714.
- Dewick, P. M. 1989. The biosynthesis of shikimate metabolites. *Nat. Prod. Rep.* **6**:263–290.
- Dopheide, T. A. A., P. Crewther, and B. E. Davidson. 1972. Chorismate mutase-prephenate dehydratase from *Escherichia coli* K-12. II. Kinetic properties. *J. Biol. Chem.* **247**:4447–4452.
- Embley, T. M. 1991. The family *Pseudonocardiaceae*, p. 996–1027. In A. Balows, H. G. Trüper, M. Dworkin, W. Harder, and K.-H. Schleifer (ed.), *The prokaryotes*, vol. 1. Springer Verlag, New York.
- Euverink, G. J. W., G. I. Hessels, J. W. Vrijbloed, J. R. Coggins, and L. Dijkhuizen. 1992. Purification and characterization of a dual function 3-dehydroquinase dehydratase from *Amycolatopsis methanolicus*. *J. Gen. Microbiol.* **138**:2449–2457.
- Euverink, G. J. W., J. W. Vrijbloed, G. I. Hessels, and L. Dijkhuizen. Aromatic amino acid metabolism in *Amycolatopsis methanolicus*: isolation of auxotrophic mutants and feedback inhibition resistant mutants. Submitted for publication.
- Euverink, G. J. W., D. J. Wolters, and L. Dijkhuizen. 1995. Prephenate dehydratase of the actinomycete *Amycolatopsis methanolicus*: purification and characterization of wild type and deregulated mutant proteins. *Biochem. J.* **308**:313–320.
- Görisch, H. 1987. Chorismate mutase from *Streptomyces aureofaciens*. *Methods Enzymol.* **142**:464–472.
- Gray, J. V., B. Golinelli-Pimpaneau, and J. R. Knowles. 1990. Monofunctional chorismate mutase from *Bacillus subtilis*: purification of the protein, molecular cloning of the gene, and overexpression of the gene product in *Escherichia coli*. *Biochemistry* **29**:376–383.
- Gygax, D., O. Ghisalba, H. Treichler, and J. Nüesch. 1990. Study to the biosynthesis of the rifamycin-chromophore in *Nocardia mediterranei*. *J. Antibiot.* **43**:324–326.
- Hagino, H., and K. Nakayama. 1974. DAHP synthetase and its control in *Corynebacterium glutamicum*. *Agric. Biol. Chem.* **38**:2125–2134.
- Hagino, H., and K. Nakayama. 1975. Regulatory properties of chorismate mutase from *Corynebacterium glutamicum*. *Agric. Biol. Chem.* **39**:331–342.
- Haslam, E. 1993. Shikimic acid: metabolism and metabolites. John Wiley & Sons Ltd., Chichester, United Kingdom.
- Hund, H. K., B. Keller, and F. Lingens. 1987. Phenylalanine and tyrosine biosynthesis in sporeforming members of the order Actinomycetales. *Z. Naturforsch. Sect. C* **42**:387–393.
- Jensen, R. A., and E. W. Nester. 1966. Regulatory enzymes of aromatic amino acid biosynthesis in *Bacillus subtilis*. I. Purification and properties of 3-deoxy-D-arabino-heptulosonate 7-phosphate synthetase. *J. Biol. Chem.* **241**:3365–3372.

28. Kato, N., K. Tsuji, Y. Tani, and K. Ogata. 1974. A methanol utilizing actinomycete. *J. Ferment. Technol.* **52**:917–920.
29. Kell, D. B., K. Van Dam, and H. V. Westerhoff. 1989. Control analysis of microbial growth and productivity, p. 61–93. In S. Baumberg, I. S. Hunter, and P. M. Rhodes (ed.), *Microbial products: new approaches*. Cambridge University Press, Cambridge.
30. Labeda, D. P. 1995. *Amycolatopsis coloradensis* sp. nov., the avoparcin (LL-AV290)-producing strain. *Int. J. Syst. Bacteriol.* **45**:124–127.
31. Laemmli, U. K., and K. Favre. 1973. Maturation of the head of bacteriophage T4. I. DNA packaging events. *J. Mol. Biol.* **80**:575–599.
32. Lorence, J. H., and E. W. Nester. 1967. Multiple molecular forms of chorismate mutase in *Bacillus subtilis*. *Biochemistry* **6**:1541–1545.
33. Lowe, D. A., and D. W. S. Westlake. 1972. Regulation of chloramphenicol synthesis in *Streptomyces* sp. 3022a: branch-point enzymes of the shikimic acid pathway. *Can. J. Biochem.* **50**:1064–1073.
34. Martin, J. F., and P. Liras. 1982. Microbial fundamentals, p. 211–233. In H. J. Rehm and G. Reed (ed.), *Biotechnology*, vol. 1. Verlag-Chemie, Weinheim, Germany.
35. Nakatsukasa, W. M., and E. W. Nester. 1972. Regulation of aromatic amino acid biosynthesis in *Bacillus subtilis* 168. I. Evidence for and characterization of a trifunctional enzyme complex. *J. Biol. Chem.* **247**:5972–5979.
36. Nasser, D., G. Henderson, and E. W. Nester. 1969. Regulated enzymes of aromatic amino acid synthesis: control, isozymic nature, and aggregation in *Bacillus subtilis* and *Bacillus licheniformis*. *J. Bacteriol.* **98**:44–50.
37. Shio, I., and S. Sugimoto. 1979. Two components of chorismate mutase in *Brevibacterium flavum*. *J. Biochem.* **86**:17–25.
38. Speth, A. R., H. K. Hund, and F. Lingens. 1989. Terminal phenylalanine and tyrosine biosynthesis of *Microtetraspora glauca*. *Biol. Chem. Hoppe-Seyler* **370**:591–599.
39. Stuart, F., and I. S. Hunter. 1993. Purification and characterisation of 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase from *Streptomyces rimosus*. *Biochim. Biophys. Acta* **1161**:209–215.
40. Sugimoto, S., and I. Shio. 1980. Purification and properties of dissociable chorismate mutase from *Brevibacterium flavum*. *J. Biochem.* **88**:167–176.
41. Sugimoto, S., and I. Shio. 1980. Purification and properties of bifunctional 3-deoxy-D-arabino-heptulosonate 7-phosphate synthetase chorismate mutase component A from *Brevibacterium flavum*. *J. Biochem.* **87**:881–890.
- 41a. Vrijbloed, J. W., J. Madoñ, and L. Dijkhuizen. 1994. A plasmid from the methylotrophic actinomycete *Amycolatopsis methanolica* capable of site-specific integration. *J. Bacteriol.* **176**:7087–7090.
42. Vrijbloed, J. W., J. Madoñ, and L. Dijkhuizen. Transformation of the methylotrophic actinomycete *Amycolatopsis methanolica* with plasmid DNA. Stimulatory effect of a pMEA300-encoded gene. Plasmid, in press.
43. Xia, T. H., and J. S. Chiao. 1989. Regulation of the biosynthetic pathway of aromatic amino acids in *Nocardia mediterranei*. *Biochim. Biophys. Acta* **991**:6–11.