

Isolation and Characterization of Polygalacturonase Genes (*pecA* and *pecB*) from *Aspergillus flavus*†

M. P. WHITEHEAD,^{1‡} M. T. SHIEH,¹ T. E. CLEVELAND,² J. W. CARY,² AND R. A. DEAN^{1*}

Department of Plant Pathology and Physiology, Clemson University, Clemson, South Carolina 29634-0377,¹
and Southern Regional Research Laboratory, Agricultural Research Service,
U.S. Department of Agriculture, New Orleans, Louisiana 70179²

Received 8 May 1995/Accepted 9 July 1995

Two genes, *pecA* and *pecB*, encoding endopolygalacturonases were cloned from a highly aggressive strain of *Aspergillus flavus*. The *pecA* gene consisted of 1,228 bp encoding a protein of 363 amino acids with a predicted molecular mass of 37.6 kDa, interrupted by two introns of 58 and 81 bp in length. Accumulation of *pecA* mRNA in both pectin- or glucose-grown mycelia in the highly aggressive strain matched the activity profile of a pectinase previously identified as P2c. Transformants of a weakly aggressive strain containing a functional copy of the *pecA* gene produced P2c in vitro, confirming that *pecA* encodes P2c. The coding region of *pecB* was determined to be 1,217 bp in length interrupted by two introns of 65 and 54 bp in length. The predicted protein of 366 amino acids had an estimated molecular mass of 38 kDa. Transcripts of this gene accumulated in mycelia grown in medium containing pectin alone, never in mycelia grown in glucose-containing medium, for both highly and weakly aggressive strains. Thus, *pecB* encodes the activity previously identified as P1 or P3. *pecA* and *pecB* share a high degree of sequence identity with polygalacturonase genes from *Aspergillus parasiticus* and *Aspergillus oryzae*, further establishing the close relationships between members of the *A. flavus* group. Conservation of intron positions in these genes also indicates that they share a common ancestor with genes encoding endopolygalacturonases of *Aspergillus niger*.

The ability to degrade plant tissues is an important characteristic for many plant pathogenic and saprophytic organisms. Hence, microorganisms produce a battery of plant cell wall-degrading enzymes (19). Among these are endopolygalacturonases (EC 3.2.1.15), which degrade pectin, a major constituent of plant cell walls. Endopolygalacturonases have been proposed as being important in some plant pathogen interactions (4, 38). They are the first cell wall-degrading enzymes produced by fungal pathogens when cultured on isolated plant cell walls or during infection (3). Additionally, endopolygalacturonases are sometimes phytotoxic (31), which may induce the hypersensitive response in incompatible interactions (19). Endopolygalacturonases also release oligogalacturonides from cell walls which can result in the elicitation of plant defense responses (10).

Several fungal endopolygalacturonases have been cloned and characterized, including those from *Aspergillus niger*, *Aspergillus tubigenis*, *Cochliobolus carbonum*, *Fusarium moniliforme*, and *Sclerotinia sclerotiorum* (5-8, 46). Disruption of the endopolygalacturonase gene of *C. carbonum* had no effect on the pathogenicity to maize (41). Analyses of genes from *A. niger* have concentrated on their use in the food industry and for juice clarification. However, endopolygalacturonase genes have not been previously cloned from an opportunistic plant-pathogenic fungus, such as *Aspergillus flavus*, in which endo-

polygalacturonases are thought to be essential in aggressiveness towards infection of cotton bolls (4, 13).

A. flavus produces aflatoxins, the most carcinogenic naturally occurring compounds known, during infection of agronomically important crops such as corn, peanuts, and cotton. Two widely occurring strains of *A. flavus* can be isolated from naturally infected cotton bolls. Highly aggressive strains degrade the intercarpellary membrane that divides a cotton boll and are able to spread throughout the boll. Weakly aggressive strains cannot degrade this membrane, and thus their growth is restricted to individual locules. The only consistent difference detected between these two strain types is their pectinase production. Both produce three pectinase activities, two endopolygalacturonases (P1 and P3) and a pectin methyl esterase (PE). However, highly aggressive isolates produce an additional endopolygalacturonase (termed P2c) (13, 14). P2c activity accumulates in medium containing glucose or pectin, whereas the activity of the other enzymes is repressed by glucose and is only produced in pectin medium. The ability to produce endopolygalacturonases in the presence of glucose is rare in filamentous fungi (2). However, the ability to produce enzyme activity in the absence of an inducer may be a key element in the pathogenesis by some organisms.

The two strain types of *A. flavus* provide an excellent opportunity to analyze, at the molecular level, production of endopolygalacturonases in an opportunistic pathogen and to determine their role in pathogenesis. Further knowledge of the control and regulation of endopolygalacturonase also may prove useful to industrial production of this enzyme. We describe here the cloning and characterization of two endopolygalacturonase genes from a highly aggressive strain of *A. flavus*. We establish by DNA transformation that one of these genes, *pecA*, encodes the pectinase P2c.

* Corresponding author. Mailing address: Department of Plant Pathology and Physiology, 120 Long Hall, Clemson University, Box 340377, Clemson, SC 29634-0377. Phone: (803) 656-5737. Fax: (803) 656-0274. Electronic mail address: RDEAN@CLEMSON.EDU.

† This is journal paper no. 4032 of the South Carolina Agricultural Experiment Station, Clemson, SC 29634.

‡ Present address: School of Applied Sciences, University of Wolverhampton, Wolverhampton WV1 1SB, United Kingdom.

MATERIALS AND METHODS

Strains and culture conditions. *A. flavus* 70 and 12 (15) were maintained on 4% V-8 agar medium (pH 6.5) at 30°C. *Escherichia coli* DH5 α and XL1-blue were used for all bacterial manipulations.

A. flavus was grown in the liquid medium as described by Adye and Mateles (1) at 30°C and 200 rpm. The medium contained either pectin at 0.5% or other carbon sources at 1%. Mycelia were grown in glucose medium for 15 h before being shifted to glucose, pectin, pectin and glucose, or mannitol for 5 h prior to harvest for RNA extraction (17). All carbon sources were purchased from Sigma Chemical Company (St. Louis, Mo.).

DNA and RNA isolation and manipulation. Plasmid DNA was isolated by the alkaline lysis method. Restriction enzyme site mapping was conducted according to standard methods (30). *A. flavus* genomic DNA was isolated as described by Horng et al. (20). To isolate RNA, approximately 0.5 g of press dried mycelia was ground to a fine powder in liquid N₂. The powder was placed in oak ridge tubes containing liquid N₂, 3 ml of 8 M guanidinium HCl, and 3 ml of phenol-chloroform. The powder was thawed on ice and centrifuged at 3,000 rpm for 10 min in an SS34 rotor. The supernatant was placed over a 1.5-ml 5.7 M CsCl₂ cushion and centrifuged for 20 h at 26,000 rpm in an SW41 swinging bucket rotor. The resulting RNA pellet was resuspended in deionized water.

Digested genomic DNA was fractionated on a 0.8% TAE (Tris-acetic acid-EDTA) agarose gel, while total RNA was separated on a 1% formaldehyde agarose gel, before being transferred to a Hybond N nylon filter (Amersham Corp., Arlington Heights, Ill.), as described by Maniatis et al. (30). DNA labeling and Southern and Northern (RNA) hybridizations were performed according to the method of Maniatis et al. (30) in 6 \times SSC (1 \times SSC is 0.15 M NaCl plus 0.015 M sodium citrate)-5 \times Denhardt's solution-0.1% sodium dodecyl sulfate (SDS)-50 mM PO₄ buffer (pH 6.6). The temperature of the hybridizations and washes varied between 56 and 68°C for different experiments. The membranes were washed in 3 \times to 0.1 \times SSC-0.1% SDS, before being exposed to autoradiographic film.

DNA sequencing. Restriction enzyme fragments from pCFC9, pCFC12, and pCFC41 were subcloned into pBluescript KS+ or SK+. Both strands of the subclones were sequenced by the dideoxynucleotide chain termination method (40). Either a Sequenase version 2.0 (United States Biochemical, Cleveland, Ohio) with [α -³²S]dCTP or a Taq Dye Deoxy Terminator Cycle Sequencing (Applied Biosystem, Foster City, Calif.) kit was used to label the DNA. Reactions were analyzed on a standard sequencing gel (30) or an Applied Biosystems autosequencer, respectively. DNA and protein sequence analyses were performed by using the software package GeneWorks 2.2 for Macintosh (Intelligentics, Inc., Mountain View, Calif.).

Genomic library screening and phage manipulations. A genomic library of *A. flavus* 70 was constructed by cloning partially digested *Sau*3A genomic DNA into EMBL3. Phage clones were maintained in *E. coli* LE392. Screening of the genomic library was performed according to the method of Sambrook et al. (39). Hybridization of the *C. carbonum pgn-1* gene or *Aspergillus parasiticus pec-1* gene was done as described above at 56 or 60°C, respectively. Individual purified plaques were amplified, and DNA was isolated by using Magic Lambda Preps (Promega, Madison, Wis.) according to the manufacturer's instructions.

Construction of plasmid pCFC80. Plasmid pCFC80 was constructed by first cloning a 1.7-kb *Bam*HI-*Hind*III fragment of pCFC41 into pBluescript SK+. A 2.3-kb *Bam*HI fragment from pCFC12 was then inserted into the *Bam*HI site to create plasmid pCFC80, which contains the entire *pecA* gene.

Transformation. *niaD* mutants from strain 12 (weakly aggressive) were isolated from chlorate medium as described previously (44). Several mutants that were unable to produce P2c were defective in *niaD*. Polyethylene glycol-mediated transformation was adapted from the method of Unkels et al. (44). Protoplasts of one *niaD* mutant were cotransformed with plasmids pCFC80 and pSTA10 (containing the *niaD* gene from *A. niger*) (43). Following transformation, protoplasts were plated on selective medium (minimal medium containing 1.2 M sorbitol and 10 mM nitrate as the sole nitrogen source). Transformants were streaked for single colonies and further evaluated.

Pectinase activity assays. Transformants were grown in liquid medium containing glucose as the sole carbon source for 48 h. Culture filtrates were then collected and examined for P2c activity by cup-plate assay as described previously (13). Alternatively, culture filtrates were collected from selected transformants grown in liquid medium containing pectin. These culture filtrates were then concentrated by sucrose overlay and dialyzed against 1% glycine (14). The 10-fold-concentrated filtrates were then applied to isoelectric focusing gels (pH 3.5 to 9.5). Isoelectric focusing gels were assayed for pectinase activity by pectin-agarose overlay (14).

Nucleotide sequence accession numbers. The GenBank accession numbers of the nucleotide sequences of *pecA* and *pecB* (see Fig. 2 and 3) are U05015 and U05020, respectively.

RESULTS

Isolation of genes encoding endopolygalacturonases. A strategy of cloning by heterologous hybridization was employed. A 541-bp *Pst*I-*Sac*I fragment of the *C. carbonum pgn-1*




FIG. 1. Partial restriction map and localization of *pecB* in pCFC9 (A) and *pecA* in pCFC12 and pCFC41 (B). The large solid arrows indicate the positions of the genes and the direction of transcription. The sequencing strategy for each gene is outlined below the restriction map by small arrows. Ac, *Acc*I; Bg, *Bgl*I; B, *Bam*HI; E, *Eco*RI; H, *Hind*III; K, *Kpn*I; P, *Pst*I; S, *Sal*I; St, *Stu*I; V, *Eco*RV; X, *Xba*I.

endopolygalacturonase gene (46) was used to probe genomic *A. flavus* DNA isolated from strains 70 (highly aggressive) and 12 (weakly aggressive). This fragment consistently hybridized to an approximately 3.8-kb *Bam*HI fragment in both strains (data not shown) and was therefore hybridized to a genomic library of *A. flavus* 70 constructed in EMBL3. Strongly hybridizing plaques were isolated and analyzed in detail. A 3.8-kb *Bam*HI fragment, which hybridized to the *C. carbonum pgn-1* gene, was isolated and cloned into pBluescript KS+ to create pCFC9. pCFC9, in addition to hybridizing to a 3.8-kb *Bam*HI fragment, produced a weak hybridization signal at 2.5 kb when hybridized to *Bam*HI-digested *A. flavus* DNA (data not shown). pCFC9 was restriction mapped, and the region hybridizing to the *C. carbonum Pst*I-*Sac*I fragment was determined by Southern analysis (Fig. 1A). This putative endopolygalacturonase was termed *pecB*.

In order to isolate additional endopolygalacturonase genes, a cDNA clone of the endopolygalacturonase, *pec-1*, of *A. parasiticus* (9) was used to probe *A. flavus* 70 and 12 genomic DNA. The *pec-1* gene preferentially hybridized in 6 \times SSC at 60°C to 2.5- and 2.3-kb *Bam*HI fragments and only very slightly to the 3.8-kb *Bam*HI fragment which contains *pecB* (data not shown). The *pec-1* gene was therefore used to screen the genomic library of *A. flavus* 70. Strongly hybridizing plaques were isolated and analyzed further. *Bam*HI fragments, 2.5 and 2.3 kb, were isolated from one clone and subcloned into pBluescript KS+ to create pCFC41 and pCFC12, respectively. These were restriction mapped, and the regions of hybridization to *pec-1* were determined by Southern analysis (Fig. 1B). These two clones were assessed to contain one putative endopolygalacturonase gene, termed *pecA*.

Sequence analysis of *pecA* and *pecB*. The nucleotide sequence of *pecA* (Fig. 2) and *pecB* (Fig. 3) were determined by using the subcloning strategy outlined in Fig. 1. The coding



FIG. 2. Nucleotide sequence of region containing *A. flavus pecA* and deduced amino acid sequence of the encoded protein. The amino acid sequence is shown in single-letter code above the corresponding codons. Intron sequences are in lowercase letters. Potential N glycosylation sites are indicated by asterisks.

region of *pecA* consists of 1,228 bp (including introns) with the potential of encoding a protein of 363 amino acids with a molecular mass of 37,650 Da. Potential N glycosylation sites are found at Asn-162-Ile-Thr-164 and Asn-294-Gly-Ser-296. The optimum site for cleavage of the signal peptide is at Ala-16 according to the rules of von Heijne (45). The coding region of *pecA* is interrupted by two introns with lengths of 58 bp (intron 1) and 81 bp (intron 2) which are bounded by GTA--TAG and GTA--CAG, respectively. These conform to the consensus for the boundaries of fungal introns as determined by Gurr et al. (18). Additionally, the internal conserved sequence of filamentous fungal introns, PyGCTAAC, is present in both introns: TACTAAG (intron 1) and TGCTAAT (intron 2).

The coding region of the *pecB* gene is 1,217 bp (including introns) long interrupted by two introns with lengths of 65 bp (intron 1) and 54 bp (intron 2). Again, these contain the conserved splice sites of GTA--TAG and GTA--CAG characteristic of fungal introns. However, the internal intron recognition sequence is not so highly conserved, with sequences of AGT-TAAT and ATATTAC being observed for intron 1 and intron 2, respectively. The *pecB* gene encodes in a protein of 366 amino acids with a predicted molecular mass of 38,146 Da. A

signal sequence cleavage site is predicted at Ala-17 (45). One potential N glycosylation site is found at Asn-278-Ile-Thr-280.

No recognizable TATAA or CAAT motif was observed in the 5' region of *pecA*. However, sequences of TATACA and CAAAT were observed at -57 and -108, respectively, for *pecB*. Both genes obeyed the translation start rule of having a purine, normally an A, at the -3 position, to direct efficient ribosomal binding (26).

Comparisons of endopolygalacturonase sequences. The nucleic acid of the coding region and predicted amino acid sequences of *pecA* and *pecB* were compared with other fungal endopolygalacturonase sequences, by using the alignment algorithms of the GeneWorks analysis program (IntelliGenetics) (Fig. 4). The degree of nucleic acid identity ranged from 55 to 65% for the majority of genes analyzed, with some notable exceptions. The nucleotide identity between the *A. flavus pecA* gene and the *A. oryzae* gene encoding polygalacturonase (PG) and *A. parasiticus pec-1* gene was very high, 99 and 96%, respectively. Also, the regions 5' (240 bp) and 3' (180 bp) to the coding sequence of these genes demonstrated exceptional levels of identity with the *pecA* gene of *A. flavus*. *pecA* was 99% identical to the *A. oryzae* PG gene and 92% identical to *A.*




FIG. 3. Nucleotide sequence of region containing *A. flavus pecB* and deduced amino acid sequence of the encoded protein. The amino acid sequence is shown in single-letter code above the corresponding codons. Intron sequences are in lowercase letters. A potential N glycosylation site is indicated by an asterisk.

	<i>A. flavus pec A</i>	<i>A. oryzae</i>	<i>A. parasiticus</i>	<i>A. niger Pga II</i>	<i>A. tubigenis</i>	<i>A. niger Pga I</i>	<i>A. flavus pec B</i>	<i>S. sclerotiorum</i>	<i>C. carbonum</i>	<i>A. niger Pga C</i>	<i>F. moniliforme</i>
<i>A. flavus pec A</i>	99	96	65	64	67	67	63	67	59	55	
<i>A. oryzae</i>	98	96	65	64	67	67	63	67	59	55	
<i>A. parasiticus</i>	97	97	65	64	66	65	62	67	60	56	
<i>A. niger Pga II</i>	59	59	60	90	63	62	58	65	57	55	
<i>A. tubigenis</i>	61	61	62	94	64	63	57	64	55	55	
<i>A. niger Pga I</i>	61	60	61	58	58	76	61	65	60	55	
<i>A. flavus pec B</i>	58	57	58	57	58	67	60	67	59	51	
<i>S. sclerotiorum</i>	58	58	59	56	59	54	54	64	59	52	
<i>C. carbonum</i>	61	62	63	58	58	56	63	61	58	57	
<i>A. niger Pga C</i>	54	54	56	50	51	56	53	55	53	46	
<i>F. moniliforme</i>	46	46	45	41	41	39	42	43	47	38	

FIG. 4. Percentage identity of nucleotide and deduced amino acid sequences of 11 fungal endopolygalacturonase genes. The sequences were compared in all pairwise combinations from *A. oryzae* (9), *A. parasiticus* (9), *A. niger* (5-7), *S. sclerotiorum* (37), *C. carbonum* (46), *F. moniliforme* (8), and *A. flavus* (this study). N.A., nucleotides; A.A., amino acids.

parasiticus pec-1. High levels of nucleotide identity were also observed between the *A. flavus pecB* gene and the *A. niger pga-1* gene, levels much higher (76%) than that observed between the *A. niger* endopolygalacturonase genes (57 to 63%) or between the *pecA* and *pecB* genes of *A. flavus* (67%). The nucleic acid identity observed between the 5' regions of *pga-1* and *pecB* (62%) was unusually high compared with the degree of identity in the same region between *pecA* and *pecB* (34%).

Gross structural similarities between the fungal endopolygalacturonase genes were analyzed. Certain introns of the *A. niger*, *A. flavus*, and *A. parasiticus* genes are conserved in the same position (Fig. 5). The introns for the *A. oryzae* PG gene were identical in sequence and position to those of the *A. flavus pecA* gene. Thus, according to the endopolygalacturonase sequence data available, all the genes possess introns 1 and 2, except for *A. niger pga-II*, which lacks intron 1. The *A. niger pgaC* gene has a third intron not found in other genes.

Expression analysis of *pecA* and *pecB*. The *pecB* gene transcript was found to accumulate in the presence of pectin only when glucose was absent. A 1.2-kb transcript accumulated in

mycelia of strain 70 when grown in pectin medium but was completely absent when *A. flavus* was grown in either glucose or glucose and pectin (Fig. 6A). This pattern was observed for both strains 70 and 12.

The *pecA* gene product (1.3 kb) was found to accumulate in both pectin- and glucose-grown mycelia for *A. flavus* 70 (Fig. 6B). However, the level of transcript accumulation when cells were grown in pectin was increased compared with the glucose levels. *pecA* mRNA was not detected in *A. flavus* 12 grown in either pectin or glucose (data not shown).

Production of P2c by introducing active *pecA* into a weakly aggressive strain. To directly determine whether *pecA* encodes P2c, a plasmid containing the *pecA* gene (pCFC80) was introduced into strain 12 (weakly aggressive), and P2c production was examined. The *pecA* gene was introduced into *niaD* mutants of *A. flavus* 12 by cotransformation with a plasmid containing the *niaD* gene and a plasmid containing the *pecA* gene (pCFC80) through polyethylene glycol-mediated transformation. Transformants were then screened for their ability to produce P2c when grown in medium containing glucose. Pectinase activities were detected by the cup-plate assay (13). Several transformants potentially expressing P2c and several others not expressing P2c were further evaluated for pectinase activity by using a pectin agarose overlay of isoelectric focusing gels (15). While the recipient parental strain and negative (not expressing P2c) transformants showed only PE bands, both PE and P2c bands were detected in the positive (expressing P2c) transformants (Fig. 7). To check the genotypes of these transformants, genomic DNA from positive and negative transformants was digested with *HindIII* and hybridized with *pecA* (Fig. 8). As predicted from the restriction map of *pecA* (Fig. 1B), three fragments (1.1, 2.3, and 2.5 kb) were detected in strain 70 (Fig. 8A, lane 1). On the other hand, hybridization of *pecA* to *HindIII*-digested DNA from strain 12 detected two of the same three bands (1.1 and 2.5 kb), with the exception that the probe hybridized to a 2.9-kb band instead of a 2.3-kb band (Fig. 8A, lanes 2 and 3). This polymorphism was valuable in establishing the addition of the *pecA* gene from strain 70 to strain 12. All transformants tested expressing P2c contained both the 2.3- and 2.9-kb bands, while only the 2.9-kb band was detected in the negative transformants (Fig. 8B). The additional bands detected in two of the transformants expressing P2c suggest multiple integrations of the *pecA* gene (Fig. 8B, lanes 2 and 3).

<i>A. flavus pecA</i>	GTCTCAACGA	TGGCACCCT	216	Intron 1	275	GTCATCTTCT	CGGGTGAGAC
<i>A. parasiticus pec1</i>	GTCTCAACGA	TGGCACCCT	216	Intron 1	269	GTCATCTTCT	CGGGTGAGAC
<i>A. niger pgaII</i>	GTCTCACCAG	CGGTACCAAG	213	-----	214	GTCATCTTCC	AGGGCACCAC
<i>A. niger pgaI</i>	ATGCTGCTGA	TGGCTCCACC	228	Intron 1	281	ATCACCTTCG	AGGGCACCAC
<i>A. flavus pecB</i>	AGGCCAAGGA	CGGTGCCACC	225	Intron 1	291	ATCACCTTCG	AGGGCACCAC
<i>A. niger pgaC</i>	ACCTGAATGA	TGGAACCCAC	264	Intron 1	340	GTCATCTTCC	AGGGAGAAAC
<i>A. flavus pecA</i>	TCGCCATCAA	CTCTGGATCG	691	Intron 2	773	CACATCACTT	TCACCAACGG
<i>A. parasiticus pec1</i>	TCGCCATCAA	CTCTGGATCG	685	Intron 2	762	CATATCACTT	TCACCAACGG
<i>A. niger pgaII</i>	TTGCGGTAA	CTCTGGCGAG	630	Intron 1	683	AACATCTGGT	TCACCGGGCG
<i>A. niger pgaI</i>	TTGCCATCAA	CTCTGGCGAG	700	Intron 2	763	AGCATCTTCT	TCACCGGGCG
<i>A. flavus pecB</i>	TGGCCATCAA	CTCTGGTGAG	707	Intron 2	763	AACATCGAAT	TCACCGGGCG
<i>A. niger pgaC</i>	TTGCCATCAA	TTCTGGAGAG	765	Intron 2	822	AACATTTATT	TCAGTGCCAG
<i>A. flavus pecA</i>	GGTTGTCAAC	TCCGAGAATG	902	-----	903	GTGTCGGTAT	CAAGACCGTC
<i>A. parasiticus pec1</i>	GGTTGTCAAC	TCTCAGAACG	891	-----	892	GTGTCGGTAT	CAAGACCGTC
<i>A. niger pgaII</i>	CGTGAGCAAT	TCCGAAAACG	812	-----	813	CCGTCCGAAT	TAAGACCATC
<i>A. niger pgaI</i>	TGTCAGCAAC	TCCGCAACG	892	-----	893	GTGTCGGTAT	CAAGACCATC
<i>A. flavus pecB</i>	CGTCTGTCAC	TCCGACAACG	892	-----	893	GTATCCGTAT	CAAGACCATC
<i>A. niger pgaC</i>	TGTTCTCAAG	TCCGACAACG	951	Intron 3	1005	CAATCCGTAT	CAAGACCATC

FIG. 5. Alignment of intron positions for endopolygalacturonase genes from *Aspergillus* species. The sequences aligned are from *A. parasiticus* (9), *A. niger* (5-7), and *A. flavus* (this study). Where present, introns are located in the same position in the nucleotide sequence.




FIG. 6. Transcript accumulation profiles of *pecA* and *pecB* in *A. flavus* 70 grown on various carbon sources. A 3.8-kb *Bam*HI fragment containing the *pecB* gene (A) and a 2.5-kb *Bam*HI fragment containing part of the *pecA* gene (B) were radiolabelled and hybridized to total RNA (10 μ g per lane) isolated from *A. flavus* 70 grown on pectin (lane 1), pectin and glucose (lane 2), or glucose (lane 3).

DISCUSSION

Two genes, *pecA* and *pecB*, encoding endopolygalacturonases were cloned from a highly aggressive strain of *A. flavus*. One gene, *pecB*, was found in both weakly and highly aggressive strains of *A. flavus* and was expressed at apparently similar levels in both organisms. This gene probably encodes either the P1 or P3 activity, because the transcript accumulates only in mycelia grown in medium containing pectin alone. The transcript of *pecA* was found to accumulate in mycelia grown in pectin- and glucose-containing medium, although the level of expression in pectin-grown mycelia was elevated compared with the level in glucose-grown cells. Expression of *pecA* was not detected in the weakly aggressive strain under any conditions. A *pecA* homolog was detected in the weakly aggressive strain by Southern hybridization; however, no transcript was detected. The *Hind*III polymorphism (2.3 versus 2.9 kb) between the two strains indicates differences in DNA composition and may account for the lack of *pecA* expression in the weakly aggressive strain. These data are consistent with *pecA* encoding the pectinase P2c. Further evidence that *pecA* encodes P2c is provided by transformation studies. A pectinase with the same activity profile as P2c was detected in the weakly aggressive strain when the *pecA* gene from an aggressive strain was introduced.

The expression of *pecA* was increased by the addition of pectin; hence, this gene retains a mechanism for pectin induction as observed for the expression of *pecB*. The *pecB* transcript did not accumulate in mycelia grown on neutral carbon sources, such as mannitol (results not shown), indicating that only pectin will induce the expression of *pecB*. However, *pecA* is expressed at similar levels in cells grown in glucose and in mannitol. Thus, its promoter contains elements for constitutive expression in the absence of pectin and also elements that enhance transcription in the presence of pectin.

There are a number of possible explanations for this finding. The low-level constitutive production of extracellular enzymes by plant-pathogenic organisms in the absence of an inducer has been postulated to be important for infection (25, 32) and may enable some plant pathogens to be primed to infect their hosts, quickly degrading and overcoming the mechanical barriers to infection. Breakdown products of the structural components of the plant cell wall are rapidly produced and further act to induce expression of extracellular enzymes, facilitating the infection process. The isolation of *pecA* provides an excellent opportunity to study this mechanism in a plant-pathogenic organism.

The elucidation of the mechanisms regulating the expression of *pecA* also may be useful to the industrial production of




FIG. 7. Pectinase activities in culture filtrates of transformants grown in pectin medium. Activities were examined by pectin-agarose overlay of isoelectric focusing gels. Lanes: 1 and 2, culture filtrates from transformants (7-82-11 and 6-10-19) not expressing P2c; 3 to 5, culture filtrates from transformants (7-82-14, 9-10-14, and 9-10-3) expressing P2c; 6 and 7, culture filtrates of strain 70 and strain 12, respectively.

pectinases. *A. niger* strains are employed as the main industrial producer of pectinases for use in the food industry. Considerable effort has been placed on increasing the production of pectinases in culture. If the mechanism by which *pecA* produces pectinase in the presence of glucose is determined, it may be possible to genetically alter native *A. niger* pectinase genes or introduce the *A. flavus pecA* gene and increase the productivity of pectinase manufacturers.

The analysis of *pecA* and *pecB* revealed an extremely high degree of sequence identity between *A. flavus pecA*, the *A. oryzae* PG gene, and *A. parasiticus pec-1*. These organisms are all part of the *A. flavus* group of fungi, which exhibits a high degree of variation (42). The question of speciation within this group has been intensely debated and is unsettled (12, 16, 22-24, 27, 33-36).

An extremely high degree of nucleotide identity has been found in other genes in the *A. flavus* group of fungi. The *nor-1* genes of *A. flavus* and *A. parasiticus* (4a) as well as the *apa-2* gene of *A. parasiticus* and the *aglR* gene of *A. flavus* are approximately 95% identical (11). On the basis of DNA reassociation experiments, the degree of complementarity between *A. flavus* and *A. oryzae* is 100% (29). Perhaps it is not surprising that, as shown here, the *A. flavus pecA* and *A. oryzae* PG genes are essentially identical. These observations are consistent with arguments that species in the *A. flavus* group should be reduced to the subspecies level (29).

The conservation of intron positions between *A. niger*, *A. flavus*, and *A. parasiticus* endopolygalacturonase genes indicates that these organisms and endopolygalacturonase genes




FIG. 8. Hybridization of *pecA* to strain 12 transformants. (A) A 2.3-kb *Bam*HI fragment from pCFC12 was radiolabelled and hybridized to *Hind*III-digested strain 70 (lane 1), strain 12 (lane 2), and a *niaD* mutant of strain 12 (lane 3). (B) The probe described for panel A was hybridized to three transformants expressing P2c, 7-82-14, 9-10-3, and 9-10-14 (lanes 1 to 3, respectively) and two transformants not expressing P2c, 6-10-19 and 7-82-11 (lanes 4 and 5, respectively).

share a common ancestor in their phylogeny. This finding and the high level of nucleotide identity seen between the *A. flavus* *pecB* gene and the *A. niger* *pgal* gene suggest that these are analogous genes. Duplication events that caused a number of endopolygalacturonase genes to be present in the genome of a common ancestor may have occurred before the divergence of *A. niger* from *A. parasiticus* and *A. flavus*.

The cloning and characterization of *pecA* and *pecB* from *A. flavus* now makes it possible to determine the role that pectinases, P2c (encoded by *pecA*) in particular, play in the infection of cotton bolls. As shown here, we have isolated transformants from a weakly aggressive strain which contain the *pecA* gene and produce P2c activities in vitro. In addition, transformants are being constructed from highly aggressive strains which contain a disrupted and therefore inactive *pecA* gene. Studies of the abilities of those strains to infect cotton bolls should provide further information on the genetic basis for the role of P2c in fungal pathogenesis. Other experiments are under way to determine the control mechanisms for endopolygalacturonase expression. These results will enhance our understanding of the role of pectinases in plant-pathogen interactions and provide further information for the manipulation of pectinase production by industrial filamentous fungi.

ACKNOWLEDGMENTS

The work was supported by a USDA ARS-SRA grant (no. 58-4432-0-105).

Thanks are extended to E. I. Zehr, Vance Baird, and W. M. Dowler for critical review of the manuscript.

REFERENCES

- Adye, J., and R. I. Mateles. 1964. Incorporation of labelled compounds into aflatoxins. *Biochim. Biophys. Acta* **86**:418-420.
- Aguiar, G., and C. Huitron. 1990. Constitutive exo-pectinase produced by *Aspergillus* sp. CH-Y-1043 on different carbon source. *Biotechnol. Lett.* **12**:655-660.
- Anderson, A. J. 1978. Extracellular enzymes produced by *Colletotrichum lindemuthianum* and *Helminthosporium maydis* during growth on isolated bean and corn cell walls. *Phytopathology* **68**:1585-1589.
- Brown, R. L., T. E. Cleveland, P. J. Cotty, and J. E. Mellon. 1993. Spread of *Aspergillus flavus* in cotton bolls, decay of intercarpellary membranes and production of fungal pectinases. *Phytopathology* **82**:462-467.
- Brown-Jenco, C. S. Personal communication.
- Bussink, H. J. D., K. B. Brouwer, L. H. de Graaf, H. C. M. Kester, and J. Visser. 1991. Identification and characterization of a second polygalacturonase gene of *Aspergillus niger*. *Curr. Genet.* **20**:301-307.
- Bussink, H. J. D., F. P. Buxton, B. A. Fraaye, L. H. de Graaf, and J. Visser. 1992. The polygalacturonases of *Aspergillus niger* are encoded by a family of diverged genes. *Eur. J. Biochem.* **208**:83-90.
- Bussink, H. J. D., F. P. Buxton, and J. Visser. 1991. Expression and sequence comparison of the *Aspergillus niger* and *Aspergillus tubigenensis* genes encoding polygalacturonase II. *Curr. Genet.* **19**:467-474.
- Caprari, C., A. Richter, C. Bergmann, S. Lo Cicero, G. Salvi, F. Cervone, and G. de Lorenzo. 1993. Cloning and characterization of a gene encoding the endopolygalacturonase of *Fusarium moniliforme*. *Mycol. Res.* **97**:497-505.
- Cary, J. W., R. L. Brown, T. E. Cleveland, M. P. Whitehead, and R. A. Dean. 1995. Cloning and characterization of a novel polygalacturonase-encoding gene from *Aspergillus parasiticus*. *Gene* **153**:129-133.
- Cervone, F., G. De Lorenzo, L. Degra, and B. Salvi. 1987. Elicitation of necrosis in *Vigna unguiculata* Walp. by homogeneous *Aspergillus niger* endopolygalacturonase and by D-galacturonate oligomers. *Plant Physiol.* **85**:626-630.
- Chang, P. K., J. W. Cary, D. Bhatnager, T. E. Cleveland, J. W. Bennett, J. E. Linz, C. P. Woloshuk, and G. A. Payne. 1993. Cloning the *Aspergillus parasiticus* *apa-2* gene associated with the regulation of aflatoxin biosynthesis. *Appl. Environ. Microbiol.* **59**:3273-3279.
- Christensen, M. 1981. A synoptic key and evaluation of species in the *Aspergillus flavus* group. *Mycologia* **73**:1056-1084.
- Cleveland, T. E., and P. J. Cotty. 1991. Invasiveness of *Aspergillus flavus* isolates in wounded cotton bolls is associated with production of a specific fungal polygalacturonases. *Phytopathology* **81**:155-158.
- Cleveland, T. E., and S. P. McCormick. 1987. Identification of pectinases produced in cotton bolls infected with *Aspergillus flavus*. *Phytopathology* **77**:1498-1503.
- Cotty, P. J. 1989. Virulence and cultural characteristics of two *Aspergillus flavus* strains pathogenic on cotton. *Phytopathology* **79**:808-814.
- Cruickshank, R. H., and J. I. Pitt. 1990. Isozyme patterns in *Aspergillus flavus* and closely related species, p. 259-265. In R. A. Samson and J. I. Pitt (ed.), *Modern concepts in Penicillium and Aspergillus classification*. Plenum, New York.
- Dean, R. A., and W. E. Timberlake. 1989. Regulation of the *Aspergillus nidulans* pectate lyase gene (*pelA*). *Plant Cell* **1**:275-284.
- Gurr, S. J., S. E. Unkles, and J. R. Kinghorn. 1987. The structure and organization of nuclear genes of filamentous fungi, p. 93-139. In J. R. Kinghorn (ed.), *Gene structure in eukaryotic microbes*. IRL Press, Oxford.
- Hahn, M. G., P. Bucheli, F. Cervone, S. H. Doares, R. A. O'Neill, A. Darvill, and P. Albersheim. 1989. Roles of cell wall constituents in plant-pathogen interactions. *Plant-Microbe Interact.* **3**:131-181.
- Hong, J. S., P.-K. Chang, J. J. Pestka, and J. E. Linz. 1990. Development of a homologous transformation system for *Aspergillus parasiticus* with the gene encoding nitrate reductase. *Mol. Gen. Genet.* **224**:294-296.
- Kitamoto, N., T. Kimura, Y. Kito, K. Ohmiya, and N. Tsukagoshi. 1993. Structural features of a polygalacturonase gene cloned from *Aspergillus oryzae* KBN616. *FEMS Microbiol. Lett.* **111**:37-42.
- Klich, M. A., and E. J. Mullaney. 1987. DNA restriction fragment polymorphism as a tool for rapid differentiation of *Aspergillus flavus* from *Aspergillus oryzae*. *Exp. Mycol.* **11**:170-175.
- Klich, M. A., and J. I. Pitt. 1985. The theory and practice of distinguishing species of the *Aspergillus flavus* group, p. 211-220. In R. A. Samson and J. I. Pitt (ed.), *Advances in Penicillium and Aspergillus systematics*. Plenum, New York.
- Klich, M. A., and J. I. Pitt. 1988. Differentiation of *Aspergillus flavus* from *Aspergillus parasiticus* and other closely related species. *Trans. Br. Mycol. Soc.* **91**:99-108.
- Koller, W., C. R. Allan, and P. E. Kolattukudy. 1982. Role of cutinase and cell wall degrading enzymes in infection of *Pisum sativum* by *Fusarium solani* f. sp. *pisi*. *Physiol. Plant Pathol.* **20**:47-60.
- Kozak, M. 1991. Structural features in eukaryotic mRNAs that modulate the initiation of translation. *Biol. Chem.* **266**:19867-19870.
- Kozakiewicz, A. 1982. The identity and typification of *Aspergillus parasiticus*. *Mycotaxon* **15**:293-305.
- Kurtzman, C. P. 1985. Molecular taxonomy of the fungi, p. 35-63. In J. W. Bennett and L. L. Lasure (ed.), *Gene manipulations in fungi*. Academic Press, Inc., New York.
- Kurtzman, C. P., M. J. Smiley, C. J. Robnett, and D. T. Wicklow. 1986. DNA relatedness among wild and domesticated species in the *Aspergillus flavus* group. *Mycologia* **78**:955-959.
- Maniatis, T., E. F. Fritsch, and J. Sambrook. 1982. *Molecular cloning: a laboratory manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Marinelli, F., S. Di Gregorio, and V. Nuti-Ronchi. 1991. Phytoalexin production and cell death in elicited carrot suspension cultures. *Plant Sci.* **77**:261-266.
- Mendgen, K., and H. Deising. 1993. Infection structures of fungal plant pathogens—a cytological and physiological evaluation. *New Phytol.* **124**:193-213.
- Moody, S. F., and B. M. Tyler. 1990. Restriction enzyme analysis of the mitochondrial DNA of the *Aspergillus flavus* group: *A. flavus*, *A. parasiticus*, and *A. nomius*. *Appl. Environ. Microbiol.* **56**:2441-2452.
- Moody, S. F., and B. M. Tyler. 1990. Use of nuclear restriction fragment polymorphisms to analyze the diversity of the *Aspergillus flavus* group: *A. flavus*, *A. parasiticus*, and *A. nomius*. *Appl. Environ. Microbiol.* **56**:2453-2461.
- Murakami, H. 1971. Classification of the koji mold. *J. Gen. Appl. Microbiol.* **17**:281-309.
- Raper, K. B., and D. I. Fennell. 1965. *The genus Aspergillus*. Williams & Wilkins, Baltimore.
- Riou, C., G. Freyssinet, and M. Fevre. 1992. Purification and characterization of extracellular pectinolytic enzymes produced by *Sclerotinia sclerotiorum*. *Appl. Environ. Microbiol.* **58**:578-583.
- Rodríguez-Palenzuela, P., T. J. Burr, and A. Collmer. 1990. Polygalacturonase is a virulence factor in *Agrobacterium tumefaciens* biovar 3. *J. Bacteriol.* **173**:6547-6552.
- Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Sanger, F., S. Nicklen, and A. R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**:5463-5467.
- Scott-Craig, J. S., D. G. Panaccione, G. Cervone, and J. D. Walton. 1990. Endopolygalacturonases are not required for pathogenicity of *Cochliobolus carbonum* on maize. *Plant Cell* **2**:1191-1200.
- Thom, C., and K. B. Raper. 1945. *A manual of the Aspergilli*. Williams & Wilkins, Baltimore.

43. **Unkels, S. E., E. I. Campbell, D. Carrez, C. Grieve, R. Contreras, W. Fiers, C. A. M. J. J. van den Hondel, and J. R. Kinghorn.** 1989. Transformation of *Aspergillus niger* with the homologous nitrate reductase gene. *Gene* **78**:157–166.
44. **Unkels, S. E., E. I. Campbell, Y. M. J. T. de Ruiter-Jacobs, M. Broekhuijsen, A. Macro, D. Carrez, R. Contreras, C. A. M. J. J. van den Hondel, and J. R. Kinghorn.** 1989. The development of a homologous transformation system for *Aspergillus oryzae* based on the nitrate assimilation pathway: a convenient and general selection system for filamentous fungal transformation. *Mol. Gen. Genet.* **218**:99–104.
45. **von Heijne, G.** 1986. A new method for predicting signal sequence cleavage sites. *Nucleic Acids Res.* **14**:4683–4691.
46. **Walton, J. D., and F. Cervone.** 1990. Endopolygalacturonase from the maize pathogen *Cochliobolus carbonum*. *Physiol. Mol. Plant Pathol.* **36**:351–359.