

## The Enterococcal Surface Protein, Esp, Is Involved in *Enterococcus faecalis* Biofilm Formation

ALEJANDRO TOLEDO-ARANA,<sup>1</sup> JAIONE VALLE,<sup>1</sup> CRISTINA SOLANO,<sup>1</sup> MARÍA JESÚS ARRIZUBIETA,<sup>1</sup>  
CARME CUCARELLA,<sup>2</sup> MARTA LAMATA,<sup>3</sup> BEATRIZ AMORENA,<sup>1</sup> JOSÉ LEIVA,<sup>3</sup>  
JOSÉ RAFAEL PENADÉS,<sup>2</sup> AND IÑIGO LASA<sup>1\*</sup>

*Instituto de Agrobiotecnología y Recursos Naturales and Departamento de Producción Agraria, Universidad Pública de Navarra-Consejo Superior de Investigaciones Científicas, Campus de Arrosadía, 31006 Pamplona,<sup>1</sup> Unit of Biochemistry, Department of Basic Biomedical Sciences, Cardenal Herrera-CEU University, 46113 Moncada,<sup>2</sup> and Department of Microbiology, University Clinics, 31008 Pamplona,<sup>3</sup> Spain*

Received 24 April 2001/Accepted 11 July 2001

**The enterococcal surface protein, Esp, is a high-molecular-weight surface protein of unknown function whose frequency is significantly increased among infection-derived *Enterococcus faecalis* isolates. In this work, a global structural similarity was found between Bap, a biofilm-associated protein of *Staphylococcus aureus*, and Esp. Analysis of the relationship between the presence of the Esp-encoding gene (*esp*) and the biofilm formation capacity in *E. faecalis* demonstrated that the presence of the *esp* gene is highly associated ( $P < 0.0001$ ) with the capacity of *E. faecalis* to form a biofilm on a polystyrene surface, since 93.5% of the *E. faecalis esp*-positive isolates were capable of forming a biofilm. Moreover, none of the *E. faecalis esp*-deficient isolates were biofilm producers. Depending on the *E. faecalis* isolate, insertional mutagenesis of *esp* caused either a complete loss of the biofilm formation phenotype or no apparent phenotypic defect. Complementation studies revealed that Esp expression in an *E. faecalis esp*-deficient strain promoted primary attachment and biofilm formation on polystyrene and polyvinyl chloride plastic from urine collection bags. Together, these results demonstrate that (i) biofilm formation capacity is widespread among clinical *E. faecalis* isolates, (ii) the biofilm formation capacity is restricted to the *E. faecalis* strains harboring *esp*, and (iii) Esp promotes primary attachment and biofilm formation of *E. faecalis* on abiotic surfaces.**

*Enterococcus faecalis* is a saprophytic commensal that inhabits the oral cavity and gastrointestinal flora of humans and animals, although it can behave as an opportunistic pathogen causing severe urinary tract infections, surgical wound infections, bacteremia, and bacterial endocarditis (21, 32, 47). Over the past 2 decades, *E. faecalis* has become responsible for up to 12% of nosocomial infections, with mortality rates for bloodstream infections ranging from 20 to 68% depending on the patient population (11). The increased incidence of *E. faecalis* infection has been related to the innate resistance of this microorganism to many commonly used antimicrobial agents and to its ability to become resistant to most, and in some cases to all, of the presently available antibiotics, either by mutation or by incorporation of foreign genetic material (6, 24). However, antibiotic resistance alone does not explain the prevalence of *E. faecalis* in enterococcal nosocomial infections, since *Enterococcus faecium*, a species less susceptible to commonly used antimicrobial agents, is responsible for only 20% of hospital-acquired enterococcal infections whereas *E. faecalis* is responsible for most of the remaining infections (33). This observation strongly supports the existence of additional virulence properties that may facilitate or enhance virulence of the *E. faecalis* isolates associated with infections.

Although knowledge on the pathogenic factors of *E. faecalis* is still limited, several virulence molecules associated preferentially with infection-derived *E. faecalis* strains have been described; these include cytolysin (16, 25, 26), aggregation substance (7, 28, 35), extracellular superoxide (22, 23), surface carbohydrates (17, 24), and surface proteins, such as Ace (34), EfaA (30), and Esp (43). Among these molecular species, Esp is the only one whose role in virulence has not been defined. Esp is a large surface protein of 1,873 amino acids with an N-terminal domain (amino acids 50 to 743) without significant similarity to other proteins in the database. The central core region (amino acids 744 to 1665) consists of a series of two distinct tandem repeat units encoded by nearly identical DNA sequence repeats and shows global structural similarity to C alpha and Rib proteins of group B streptococci. The C-terminal domain (amino acids 1666 to 1873) contains a membrane-spanning hydrophobic region and includes a slight variation of the LPXTGX motif found in most wall-associated surface proteins of gram-positive bacteria. It is presently hypothesized that the N-terminal region of Esp might participate in interactions with the host and that the central repeat region might serve to retract the protein from the surface, hiding the protein from the immune system.

In previous work aimed at identifying new factors involved in *Staphylococcus aureus* biofilm formation, we reported a high-molecular-weight cell wall-associated protein of 2,276 amino acids named Bap (9). The corresponding *bap* gene was found among staphylococcal species isolated from

\* Corresponding author. Mailing address: Instituto de Agrobiotecnología y Recursos Naturales, Universidad Pública de Navarra, Campus de Arrosadía, Pamplona-31006, Spain. Phone: 34 948 24 28 34. Fax: 34 948 23 21 91. E-mail: ilasa@unavarra.es.

TABLE 1. Strains, oligonucleotides, and plasmids used in this study

<i>E. faecalis</i> strain, oligonucleotide, or plasmid	Relevant characteristics	Position (nucleotides) <sup>a</sup>	Strand	Source or reference
<i>E. faecalis</i> strains				
54	<i>esp</i> <sup>+</sup> , strong biofilm forming			CUN <sup>b</sup>
54M	54 <i>esp</i> ::pBT2 (Cm <sup>r</sup> )			This study
1	<i>esp</i> <sup>+</sup> , strong biofilm forming			CUN
1M	1 <i>esp</i> ::pBT2 (Cm <sup>r</sup> )			This study
11279	<i>esp</i> <sup>+</sup> , medium biofilm forming			CUN
11279M	11279 <i>esp</i> ::pBT2 (Cm <sup>r</sup> )			This study
11262	<i>esp</i> <sup>+</sup> , weak biofilm forming			CUN
11262M	11262 <i>esp</i> ::pBT2 (Cm <sup>r</sup> )			This study
14377	<i>esp</i> <sup>+</sup> , adherence <sup>++</sup> , the <i>esp</i> gene contains one A repeat and four C repeats			This study
23	<i>esp</i> <sup>-</sup> , non-biofilm forming (Cm <sup>s</sup> )			CUN
23C	23 containing pTA2 (Cm <sup>r</sup> )			This study
Oligonucleotides				
<i>esp</i> 11	TTGCTAATGCTAGTCCACGACC	1217–1238	+	43
<i>esp</i> 12	GCGTCAACACTTGCATTGCCGAA	2171–2149	–	43
<i>esp</i> 46	TTACCAAGATGGTTCTGTAGGCAC	2256–2279	+	43
<i>esp</i> 47	CCAAGTATACTTAGCATCTTTTGG	3169–3192	–	43
<i>esp</i> 2	CAGATGGATCATCTGATGAAGT	3254–3275	+	43
<i>esp</i> 5	GTAACGTTACTGTTACATCTGC	5338–5359	–	43
<i>esp</i> 3S	CGCCTTGGTATGCTAAC	1004–1020	+	This study
<i>esp</i> 4S	GGTAAGCTTACGCCGT	2435–2420	–	This study
<i>esp</i> 2S	GGGTCGACACTTCTATTCATCCTCT	5707–5691	–	This study
<i>pbac</i> S <sup>c</sup>	GCTTGCCATCAAAAATAAACTACATGGGTATAAATAGCAAT GAAATGCATTTCAAAAATATTTTGAGGAGAATTTAGT ATGTTTGGAAAA		+	This study
Plasmids				
pCU1	<i>E. coli</i> - <i>S. aureus</i> shuttle vector			3
pBT2	<i>E. coli</i> - <i>S. aureus</i> shuttle vector with a thermosensitive origin of replication for <i>S. aureus</i>			5
pTA1	Vector for recombination experiments, 0.95-kb PCR fragment of <i>esp</i> subcloned in pBT2			This study
pTA2	Vector for complementation experiments; a 4.3-kb PCR fragment containing <i>esp</i> from <i>E. faecalis</i> 14377 fused with the enterococcal constitutive <i>bacA</i> promoter cloned in pCU1			This study

<sup>a</sup> Nucleotide sequence positions refer to the *esp* sequence in the GenBank database under accession no. AF034779.

<sup>b</sup> CUN, University Clinics of Navarra.

<sup>c</sup> Regions -35 and -10 and the ribosome binding site from the *bacA* promoter according to Fujimoto et al. (15) are underlined. The ATG codon for the first amino acid of Esp is also underlined.

bovine mastitis but not among the human *S. aureus* isolates tested. Interestingly, all of the *S. aureus* isolates harboring Bap were strong biofilm producers, and transposon-insertional mutagenesis of *bap* caused the loss of the biofilm formation phenotype. Bap showed global organizational similarities to an outer membrane protein-like protein of *Pseudomonas putida* involved in adhesion to seeds (12), to a proline/threonine-rich protein of unknown function of *Salmonella enterica* serovar Typhi, and to the Esp protein of *E. faecalis*.

Taking into account the global structural similarity between Bap and Esp, in this work we analyzed the association between the ability of *E. faecalis* isolates to produce a biofilm and the presence of *esp*. Furthermore, we performed insertional mutagenesis of *esp* in different *E. faecalis* isolates and analyzed the effects produced in different steps of the biofilm formation process at the macroscopic and microscopic levels. A role as a putative virulence factor involved in bacterial biofilm formation was assigned to Esp.

## MATERIALS AND METHODS

**Bacterial strains, plasmids, and culture conditions.** A total of 200 enterococcal isolates were used: 152 *E. faecalis* isolates, 39 *E. faecium* isolates, 7 *E. avium* isolates, 1 *E. gallinarum* isolate, and 1 *E. durans* isolate. They were isolated from patients during 1999 and 2000 at the Microbiology Department of the University Clinics of Navarra, Pamplona, Spain. Bacterial species were identified with API STREP (bioMérieux, Marcy l'Etoile, France) or GPI-Vitek (bioMérieux).

*Escherichia coli* XL1-Blue cells were grown in Luria-Bertani broth or on Luria-Bertani agar (Pronadisa, Madrid, Spain) with appropriate antibiotics. Enterococci were grown in brain heart infusion (BHI) broth, BHI agar, or Trypticase soy broth (TSB) supplemented with glucose (0.25 or 0.5%, wt/vol), as required. Antibiotics were used at the following concentrations: chloramphenicol, 20 µg/ml; and ampicillin, 100 µg/ml.

**DNA manipulations.** Routine DNA manipulations were performed using standard procedures (4, 41) unless otherwise stated. Plasmid DNA from *E. coli* was purified with a Quantum Prep Plasmid Miniprep (Bio-Rad) or Midiprep (QIAGEN) kit. Plasmids were transformed into *E. faecalis* by electroporation using a previously described method (48). Enterococcal transformations were enhanced by inducing chloramphenicol acetyltransferase translation with a subinhibitory concentration of chloramphenicol (0.2 µg/ml) for 2 h after electroporation. Restriction enzymes were purchased from Boehringer Mannheim and used according to the manufacturer's instructions. Oligonucleotides were obtained from Life Technologies (Table 1).

For Southern hybridization, chromosomal DNA was purified as previously described (31), digested with *EcoRI* and *PstI*, and analyzed by agarose gel electrophoresis. DNA fragments were transferred by alkaline capillary blotting onto nylon membranes (Hybond-N; Amersham Life Science) using standard methods (4). A 950-bp PCR fragment obtained with primers *esp11* and *esp12* (Table 1) was used as a DNA probe. Labeling of the probe and DNA hybridization were performed according to the protocol supplied with the PCR-DIG DNA-labeling and chemiluminescent detection kit (Boehringer Mannheim).

**Epidemiology study on the presence of the *esp* gene and repeat number variation.** Primers *esp11* and *esp12* (Table 1) were used to amplify a 950-bp fragment within the N-terminal region of *esp* and to detect its presence in DNAs from the different enterococcal isolates. To assess the repeat number variation among the *esp*-positive enterococcal isolates, primers *esp46* and *esp47* (Table 1) were used for amplification across the A repeat region, whereas primers *esp2* and *esp5* (Table 1) were used for amplification across the C repeat region. According to the sequences of the *esp* repeat regions described by Shankar et al. (43), the number of A repeats was calculated as follows:  $nA = (f_{aPCR} - 182)/252$ , where  $nA$  is the number of A repeats and  $f_{aPCR}$  is the size of the DNA fragment after PCR amplification using primers *esp46* and *esp47* (Table 1). Similarly, the number of C repeats was calculated as follows:  $nC = (f_{cPCR} - 384)/246$ , where  $nC$  is the number of C repeats and  $f_{cPCR}$  is the size of the DNA fragment after PCR amplification using primers *esp2* and *esp5* (Table 1).

**Disruption of *esp*.** For disruption of *esp* in three *esp*-positive biofilm-forming isolates (Table 1), a 950-bp PCR fragment within the N-terminal region of the *esp* gene amplified with primers *esp11* and *esp12* was cloned into the pGEM-T Easy vector (Promega, Madison, Wis.). The *esp* fragment was then cloned into the *EcoRI* site of the shuttle vector pBT2, and the resulting plasmid (pTA1) was transformed into *E. faecalis* by electroporation. After electroporation, bacterial strains were incubated for 24 to 48 h at 30°C on BHI agar with chloramphenicol. Ten milliliters of BHI broth-chloramphenicol was subsequently inoculated with a single colony which had been previously resuspended in 100  $\mu$ l of BHI broth, and the culture was incubated for 24 h at 43.5°C without shaking. Tenfold serial dilutions of this culture in sterile BHI broth were plated on BHI agar-chloramphenicol and incubated for 24 h at 43.5°C. After overnight incubation, colonies were analyzed for disruption of the *esp* gene by colony PCR with primers *esp3S* and *esp4S* (Table 1), and the results were confirmed by Southern blot analysis.

**Biofilm assay on polystyrene plates and adherence to polyvinyl chloride (PVC) plastic.** The ability of the enterococcal strains to form a biofilm on an abiotic surface was quantified essentially as described elsewhere (37). Briefly, *E. faecalis* strains were grown overnight in TSB with 0.25% glucose at 37°C. The culture was diluted 1:40 in TSB-0.25% glucose, and 200  $\mu$ l of this cell suspension was used to inoculate sterile 96-well polystyrene microtiter plates (Iwaki, Tokyo, Japan). After 24 h at 37°C, wells were gently washed three times with 200  $\mu$ l of phosphate-buffered saline (PBS), dried in an inverted position, and stained with 1% crystal violet for 15 min. The wells were rinsed again, and the crystal violet was solubilized in 200  $\mu$ l of ethanol-acetone (80:20, vol/vol). The optical density at 595 nm ( $OD_{595}$ ) was determined using a microplate reader (Multiskan EX; Labsystems). Each assay was performed in triplicate and repeated three times.

Bacterial adherence to PVC plastic was studied using a phase-contrast microscope (magnification,  $\times 1,000$ ; Nikon Optiphot) as described elsewhere (39), with the following modifications. *E. faecalis* strains were grown overnight in TSB-0.25% glucose at 37°C. Subsequently, the culture was diluted 1:40 in TSB-0.25% glucose, and 200  $\mu$ l of this cell suspension was inoculated into the wells of 96-well microtiter dishes containing sterile PVC disks from urine collection bags and incubated for 24 h at 37°C.

To compare biofilm formation ability and adherence to PVC of wild-type, recombinant, and complemented strains, bacteria were cultured overnight in TSB-0.25% glucose, supplemented with chloramphenicol (20  $\mu$ g/ml) when appropriate, and subcultured in TSB-0.25% glucose using microtiter dishes.

**Primary adherence assay.** Early adherence of *E. faecalis* to a polystyrene surface was determined as previously described (14), with the following modifications. *E. faecalis* strains were grown in TSB-0.5% glucose, supplemented with chloramphenicol (20  $\mu$ g/ml) when appropriate, overnight at 37°C. Cultures were then adjusted with TSB-0.5% glucose to an  $OD_{578}$  of 0.1. Ten milliliters of each suspension was added to two polystyrene petri dishes. After incubation for 2 h at 37°C, petri dishes were washed three times with PBS. Cells were fixed with Bouin solution (Sigma) and Gram stained. Adherent bacterial cells were observed by oil immersion microscopy, and the mean count was determined in five microscopic fields. Each experiment was repeated three times.

**Analysis of cell surface expression of *Esp*.** *E. faecalis* strains were grown overnight in TSB-0.25% glucose, supplemented with chloramphenicol (20  $\mu$ g/ml) when appropriate, at 37°C. The culture was diluted 1:40 in TSB-0.25% glucose, and 200  $\mu$ l of this cell suspension was used to inoculate sterile 96-well

polystyrene microtiter plates (Iwaki). After 24 h at 37°C, wells were gently washed three times with 200  $\mu$ l of PBS containing 0.1% Tween 20. Wells were blocked with 5% bovine serum albumin at 37°C for 1 h prior to a 2-h incubation with anti-*Esp* serum diluted 1:5,000 in PBS containing 0.1% Tween 20. Bound antibodies were detected with a peroxidase-conjugated goat anti-rabbit immunoglobulin G antibody (Jackson ImmunoResearch Laboratories, Inc., Bar Harbor, Maine) diluted 1:2,500.

**Cell surface hydrophobicity.** The cell surface hydrophobicities of *E. faecalis* strains were verified as previously described (40), with the following modifications. Cells were grown overnight in TSB-0.25% glucose, supplemented with chloramphenicol (20  $\mu$ g/ml) when appropriate, at 37°C. Three hundred microliters of the test hydrocarbon (*n*-hexadecane; Merck, Darmstadt, Germany) was added to round-bottom test tubes containing 3 ml of washed cells which had been suspended in PUM buffer (22.2 g of  $K_2HPO_4 \cdot 3H_2O$ , 7.26 g of  $KH_2PO_4$ , 1.8 g of urea, 0.2 g of  $MgSO_4 \cdot 7H_2O$ , and distilled water to 1,000 ml [pH 7.1]) (39) to an  $OD_{470}$  of 1.0. Following a 10-min preincubation at 37°C, tubes were shaken for 30 s. The aqueous phase was carefully removed with a Pasteur pipette, and light absorbance was determined at 470 nm, using a Milton Roy 20D model spectrophotometer. The percentage of bacterial adhesion to hydrocarbon was calculated as follows:  $[1 - (OD_i/OD_f)] \times 100$ , where  $OD_i$  and  $OD_f$  are the ODs of cells resuspended in PUM buffer determined at the beginning and the end of the experiment, respectively.

**Complementation studies.** The *esp* gene was amplified with high-fidelity thermostable DNA polymerase (Expand Long Template PCR System; Roche) from *E. faecalis* 14377 because the *esp* gene of this strain contains only one A repeat and four C repeats with primers *pbacS* and *esp2S* (Table 1). The *pbacS* primer includes the -35, -10, and ribosome binding sites of the constitutive promoter of the *bacA* gene of *E. faecalis* (15, 45). The chimeric *pbac-esp* gene was cloned into the plasmid pCU1, and the resulting plasmid, pTA2, was transformed by electroporation into *E. faecalis* strain 23. Stable expression of *Esp* was analyzed by enzyme-linked immunosorbent assay (ELISA) as described above.

In order to compare the biofilm formation capacities of the wild-type strain and the complemented strain, plasmid pCU1 was introduced into the wild-type strain to render it chloramphenicol resistant, and the assays were performed in the presence of chloramphenicol (20  $\mu$ g/ml).

**Statistical analysis.** A nonparametric Kendall rank correlation analysis was used to assess the association between the presence of *esp* and biofilm formation. For analysis of primary adherence and biofilm formation, a two-tailed Student's *t* test was applied.

## RESULTS

### Relationship between biofilm formation and presence of *esp*.

We searched eubacterial genome databases for sequence homologies to the *S. aureus* Bap protein using the gapped BLASTP program (1) and found that Bap and *Esp* share 33% sequence identity and 50% sequence similarity in an overall alignment. Individual pairwise alignments between the two proteins revealed that the N-terminal domain of Bap (region B, amino acids 361 to 819) shows 33% identity with the 694-residue N-terminal domain of *Esp*. Furthermore, the C repeat region of Bap, which accounts for 52% of the protein, also shows 33% identity with the C repeat region of *Esp* (Fig. 1).

Because of this remarkable structural similarity and previous findings on the involvement of Bap in biofilm formation, we studied biofilm formation on polystyrene microtiter plates by 200 infection-derived enterococcal strains. The adherence values ( $OD_{595}$ ) obtained allowed the classification of isolates into four groups: non-biofilm forming ( $OD_{595} \leq 1$ ), weak biofilm forming ( $1 < OD_{595} \leq 2$ ), medium biofilm forming ( $2 < OD_{595} \leq 3$ ), and strong biofilm forming ( $OD_{595} > 3$ ) (Table 2). Simultaneously, the presence of the *esp* gene in all of the isolates was analyzed by PCR. The results indicate that biofilm formation ability is highly and significantly associated with the presence of *esp* (Kendall rank correlation, 0.845;  $P < 0.0001$ ). The biofilm formation ability was restricted to *E. faecalis* isolates harboring *esp*, and *esp* was detected neither in any of the *E.*




FIG. 1. Structural similarity between Bap and Esp. Shown is the percentage of identity between different regions of these proteins. The Esp structure is derived from strain MMH594 (43). The Bap structure is derived from the *S. aureus* bovine strain V329 (9). The signal peptide (SP), membrane anchor (MA), region D (D), and A and C repeats regions are shown. The N-terminal domain of Esp (residues 49 to 743) and region B of Bap (residues 361 to 819) share 33% sequence identity. The C repeat regions of Esp (residues 1064 to 1648) and Bap (residues 819 to 2148) share 33% sequence identity.

*faecium*, *E. avium*, or *E. gallinarum* isolates tested, as previously described (43), nor in any of the *E. faecalis* isolates unable to produce a biofilm. The majority (93.5%) of the *E. faecalis* *esp*-positive isolates were able to form biofilm, being classified as strong, medium, or weak biofilm producers (Table 2).

**Disruption of the *esp* gene in *E. faecalis* strains with different abilities to form a biofilm.** To further analyze the possible role of the Esp protein in *E. faecalis* biofilm formation, disruption of the *esp* gene was performed in strong (strain 54), medium (strain 11279), and weak (strain 11262) biofilm-forming strains. For this purpose, a 950-bp PCR fragment corresponding to an internal N-terminal region of *esp* was cloned into the shuttle vector pBT2, producing the pTA1 plasmid. The natural biofilm-forming strains were transformed with pTA1, and transformants were grown at a nonpermissive replicative temperature. Recombinants (Fig. 2A) were confirmed by PCR using primers *esp*3S and *esp*4S (data not shown) and Southern blotting of *Pst*I- and *Eco*RI-digested chromosomal DNA (Fig. 2B).

ELISA analysis of the expression of the Esp protein in parental strains and isogenic mutants showed that parental strains expressed this surface protein, whereas Esp could not be detected in any of the mutants (Fig. 2C). The fact that six Esp-positive strains were not able to produce a biofilm encour-

aged us to determine the presence of Esp in these strains. The results showed that five out of six of these strains did not produce ELISA-detectable Esp, suggesting that these strains either did not produce Esp under our experimental conditions or produced an Esp variant which could not be recognized by the polyclonal antiserum. Accordingly, Esp could not be detected in any of the natural *esp*-deficient strains (data not shown).

To establish a relationship between the expression of the Esp protein and the ability to form a biofilm, parental and mutant strains were also tested for their ability to form a 24-h biofilm on polystyrene microtiter plates. Surprisingly, disruption of the *esp* gene in two strong biofilm-forming strains (isolates 54 and 1) did not lead to a significant decrease in this ability. In contrast, the *esp* mutant strains of the medium (11279M) or weak (11262M) biofilm producers hardly adhered to the surface, being unable to form a biofilm and exhibiting behavior indistinguishable from that of the natural *esp*-deficient strains (Fig. 2D and E).

**Cell surface hydrophobicities of parental and mutant strains.** Figure 3 illustrates the hydrophobic nature of the cell surfaces of parental strains and isogenic *esp* mutants. Parental *esp*-positive strains always showed similar (strain 54) or higher affinity for *n*-hexadecane compared with mutant strains. Remarkably, a similar decrease in hydrophobicity was observed when the hydrophobicity of a *bap*-positive *S. aureus* strain (V329) was compared with that of its *bap*-negative mutant (M556) (data not shown).

**Variation in number of A and C repeats of *esp* in relation to biofilm formation capacity.** In order to determine whether the number of A or C repeats present in Esp was related to the intensity of the biofilm formation ability, we calculated the number of repeats in each *E. faecalis* strain. In agreement with previously reported results (43), the number of A repeat units ranged from 1 to 3 whereas the number of C repeat units ranged from 4 to 10, with a majority of the isolates displaying 7 C repeat units. No association was found between the number of A or C repeat units within the *esp* gene and the biofilm formation capacity in the *E. faecalis* *esp*-positive isolates (data not shown).

TABLE 2. Distribution of enterococcal clinical isolates according to the capacity for biofilm formation on polystyrene microtiter plates and the presence or absence of *esp*

<i>esp</i> gene status	Bacteria	No. of isolates with the following biofilm formation capacity <sup>a</sup> :				Total no. of isolates
		-	+	++	+++	
Positive	<i>E. faecalis</i>	6	26	18	43	93
Negative	<i>E. faecalis</i>	59	0	0	0	59
	Other enterococci	48	0	0	0	48
Total		113	26	18	43	200

<sup>a</sup> -, non-biofilm forming (OD<sub>595</sub> ≤ 1); +, weak biofilm forming (1 < OD<sub>595</sub> ≤ 2); ++, medium biofilm forming (2 < OD<sub>595</sub> ≤ 3); +++, strong biofilm forming (OD<sub>595</sub> > 3).




FIG. 2. Biofilm formation phenotypes of *E. faecalis* *esp* mutants. The relationship between the expression of the Esp protein and the ability to form a biofilm in wild-type *E. faecalis* strains (strains 54, 11279, and 11262), their corresponding *esp* mutant strains (strains 54M, 11279M, and 11262M), and a natural *esp*-deficient strain (strain 23) is shown. (A) Schematic representation of the predicted structure after plasmid integration into *esp*. (B) Southern blot analysis after digestion of chromosomal DNA with *Pst*I and *Eco*RI. (C) Esp production assessed by ELISA. (D) Biofilm phenotype on wells of a polystyrene microtiter plate. (E) Quantification of biofilm formation. Error bars indicate standard deviations.




FIG. 3. Differences in cell surface hydrophobicities of wild-type *E. faecalis* strains in relation to their corresponding *esp* mutants as measured by affinity to *n*-hexadecane. Assays were performed in triplicate. Mean values and standard deviations are shown. BATH, bacterial adhesion to hydrocarbon.

**Involvement of Esp in primary attachment and biofilm formation on PVC plastic.** Early adherence of the wild-type *E. faecalis* 11279 and its isogenic *esp* mutant 11279M is illustrated in Fig. 4A. Strain 11279 adhered to polystyrene much more efficiently than mutant 11279M. In addition, direct observation of PVC plastic tabs from a urine collection bag using phase-contrast microscopy showed that the parental strain produced a layer of cells covering the PVC surface (Fig. 4B). In contrast, very few cells of the *esp* mutant were attached to PVC plastic. These results on early adherence are consistent with those described above on biofilm formation in polystyrene microtiter plates.

**Esp allows primary attachment to an abiotic surface.** Complementation tests were performed to determine whether overexpression of Esp was able to confer biofilm formation ability to an *E. faecalis* *esp*-negative strain. Since the nucleotide sequence upstream of the ATG codon of the *esp* gene (accession number AF034779) is very short, we were uncertain whether the promoter of *esp* was included in this sequence. To solve this problem, *esp* was overexpressed under the control of a constitutive promoter in *E. faecalis*, as described in Materials and Methods. The expression of the Esp protein in the complemented strain 23C measured by ELISA is shown in Fig. 5A. The complemented strain showed an enhanced capacity to form a biofilm on a polystyrene surface (Fig. 5B). However, the complemented strain was always classified into the group of weak biofilm-forming strains ( $1 < OD_{595} \leq 2$ ). The number of cells bound to polystyrene was quantified in order to determine whether the enhanced capacity for biofilm formation in the complemented strain was due to a modification of primary adherence to polystyrene. It was observed that the number of attached cells in the complemented strain was at least 20-fold higher than that in the wild-type strain ( $P < 0.0001$ ), demonstrating that Esp highly enhances primary binding to polystyrene (Fig. 5C). In addition, we assessed the ability of 23C to attach to PVC plastic from urinary bags using phase-contrast




FIG. 4. Comparisons at different steps of the biofilm formation process of a wild-type strain (11279) and its *esp*-deficient mutant (11279M). (A) Primary attachment assay. Significant differences ( $P < 0.0001$ ) between 11279 and 11279M were detected. Adherent bacterial cell mean values and standard deviations are shown. (B) Attachment to PVC plastic. The capacity of strains 11279 and 11279M to form a 24-h biofilm on PVC plastic was observed by phase-contrast microscopy (magnification,  $\times 1,000$ ).

microscopy. The complemented strain 23C showed multiple groups of cells adhered to the PVC plastic. In contrast, very few individual cells were attached to the plastic in the case of the wild-type strain (data not shown).




FIG. 5. Complementation studies. (A) Esp production assessed by ELISA in *E. faecalis* 23, a wild-type *esp*-deficient strain, and in this strain complemented with pTA2 (23C), (B) Biofilm formation capacities of 23 and 23C on polystyrene plates. Assays were performed in triplicate; mean values and standard deviations are shown. (C) Capacity of these strains for primary attachment in polystyrene plates. The number of attached cells of the complemented strain 23C was significantly higher ( $P < 0.0001$ ) than that of the wild-type strain. Assays were performed in triplicate; adherent bacterial cell mean values and standard deviations are shown.

## DISCUSSION

Like other gram-positive microorganisms, enterococci are able to produce biofilms on abiotic surfaces (27), increasing their high innate resistance to antibiotics (13), yet the factors controlling enterococcal biofilm formation and maintenance remain unknown (36). The initial step in the colonization of catheters and biofilm formation is bacterial adherence to the biomaterial (20). To date, mutagenesis studies on the initial attachment of gram-positive bacteria to abiotic surfaces have been performed mainly with *Staphylococcus epidermidis* (19, 42, 46) and have shown that a variety of bacterial surface proteins are involved in the process. Among these, the best characterized protein is AtlE, a cell surface-localized autolysin which has also vitronectin-binding activity and therefore is involved in binding to both biotic and abiotic surfaces (19). Since not all *S. epidermidis* strains harboring the *atlE* gene produce a biofilm in vitro, the presence of *atlE* appears to be necessary but insufficient to induce primary attachment to abiotic surfaces.

The *esp* gene has been identified in a random sequencing project on the genome of a clinical strain of *E. faecalis* (43). It has been preferentially detected in infection-derived *E. faecalis* strains but not in other less pathogenic enterococcal species, strongly suggesting a role of Esp in the pathogenesis of *E. faecalis* (43). Further supporting this hypothesis are the results obtained in this study on the highly significant association ( $P <$

0.0001) between the presence of the *esp* gene and the ability to produce a biofilm *in vitro*. Eighty-seven out of 93 *E. faecalis* strains harboring *esp* were able to produce a biofilm *in vitro*, whereas none of the 59 *esp*-deficient strains tested were able to produce a biofilm. Furthermore, Esp expression determined by ELISA revealed that five out of the six *esp*-positive, biofilm-negative *E. faecalis* isolates analyzed did not produce Esp, strongly suggesting that the absence of biofilm formation in these strains is associated with the lack of Esp expression. These results strongly suggest that *esp* is involved in the biofilm formation process of *E. faecalis*.

Insertional mutagenesis of *esp* has shown no significant effect on biofilm formation in the strongest biofilm-producing *E. faecalis* strain analyzed (strain 54). This could be attributed to additional surface adhesins that might mediate the initial attachment to the abiotic surface in the absence of *esp*. On the other hand, the finding that none of the *esp*-defective *E. faecalis* strains was able to produce a biofilm suggests the existence of a genetic association between the presence of *esp* and the presence of these adhesins. A striking common feature between *esp* and *bap* genes, both of which are associated with abiotic surface attachment, is that they are flanked by a sequence similar to that of the transposase IS905 in the case of *esp* (24) and IS431 in the case of *bap* (our unpublished results). Based on this observation and in agreement with other authors (10, 44) it is tempting to speculate that *esp* could be part of a pathogenicity island (PAI). It is known that adhesins, which mediate the capacity of bacteria to attach to specific eukaryotic receptor molecules, are major virulence factors encoded by PAIs (18). The possibility that additional adhesins might flank *esp* and *bap* and constitute a PAI is under study.

The high degree of conservation of the nucleotide sequences of A and C repeats present in both *esp* (43) and *bap* (9) strongly suggests that these repeats could have an important role in the function of this protein. Although our data reveal that the number of repeats is not related to the amount of biofilm produced, we found, in agreement with a previous report (43), that none of the *esp*-positive strains exhibited a complete loss of either A or C repeats, strongly suggesting that both regions are important for the functionality of the protein. These A and C repeats are not the only structural feature shared by Esp and Bap proteins. Analysis of their amino acid sequence reveals the presence of dimerization domains and calcium binding motifs in both of them. We are presently evaluating the contribution of these domains to the function of Esp and Bap.

Esp exhibits characteristics of surface protein receptors designated microbial surface components recognizing adhesive matrix molecules (38) that adhere to components of the host to initiate colonization. Many of these proteins have a modular design and contain a number of tandem repeat domains that probably arise from a series of recombination and/or duplication events. Although we have no evidence for the presence of domains for binding to host factors in Esp, we cannot exclude for Esp under *in vivo* conditions a direct ligand-binding activity to the extracellular matrix or an indirect role modulating ligand-binding activity of other molecules. If this is the case, Esp-mediated adherence of *E. faecalis* to plastic biomaterials would not be the primary role of this protein in the host-bacterium interaction. It has been proposed that the presence

of Esp could increase cell surface hydrophobicity and facilitate hydrophobic interactions (43). In contrast with previous results (27) where it was observed that adherence of *E. faecalis* to urinary catheters was not related to bacterial hydrophobicity, our results demonstrate that the presence of Esp in the cell surface increases hydrophobicity, adherence to abiotic surfaces, and biofilm formation. This apparent discrepancy could be at least partially explained if the *E. faecalis* clinical strains analyzed were *esp* deficient, which would likely result in a low hydrophobicity.

Routinely, antibiotic susceptibility is determined in clinical laboratories using the broth microdilution susceptibility test. However, it is well established that antimicrobials directed to planktonic cells may not be efficient against biofilm cells (8, 29). In *E. faecalis*, improvement of the selection of effective antimicrobial agents against recalcitrant infections is urgently needed. Taking into account the strong correlation between the presence of *esp* and the ability to produce a biofilm, it may be possible to screen for putative biofilm-forming *E. faecalis* strains by testing for the presence of the *esp* gene by PCR or for its product by ELISA. The genetic (presence of *esp*) rather than phenotypic (adherence or biofilm formation) nature of this screening is advantageous in that it allows preliminary identification of strains which are highly adherent and are thus good candidates for antibiotic susceptibility testing in biofilms (2).

It has been estimated that over 65% of nosocomial infections in the developed world are derived from biofilm-related infections. This represents a warning signal, since in the near future the use of medical implants is likely to increase. In this context, bacterial molecules involved in attachment mediated by host proteins and others, such as Esp, involved in adherence to abiotic surfaces and biofilm formation could become promising therapeutic targets in control programs for eradicating persistent enterococcal infections associated with the presence of biofilms.

#### ACKNOWLEDGMENTS

Alejandro Toledo-Arana and Jaione Valle contributed equally to this work.

We express our gratitude to V. Shankar for providing the antibodies against Esp, to F. Götz for plasmid pCU1, and to R. Brückner for plasmid pBT2.

This work was supported by grant BIO99-0285 from the Comisión Interministerial de Ciencia y Tecnología and grants from the Cardenal Herrera-CEU University and from the Departamento de Educación y Cultura del Gobierno de Navarra. Alejandro Toledo-Arana is a predoctoral Mutis program fellow from Agencia Española de Cooperación Internacional (AECI), Spain. Jaione Valle is a predoctoral fellow from the Ministerio de Ciencia y Tecnología (FPI), Spain. C. Cuarellá is a predoctoral fellow from the Cardenal Herrera-CEU University.

#### REFERENCES

1. Altschul, S. F., T. L. Madden, A. A. Schaffer, J. Zhang, Z. Zhang, W. Miller, and D. J. Lipman. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**:3389–3402.
2. Amorena, B., E. Gracia, M. Monzón, J. Leiva, C. Oteiza, M. Pérez, J. Alabart, and J. Hernandez-Yago. 1999. Antibiotic susceptibility of *Staphylococcus aureus* in biofilms developed *in vitro*. *J. Antimicrob. Chemother.* **44**:43–55.
3. Augustin, J., R. Rosenstein, B. Wieland, U. Schneider, N. Schnell, G. Engelke, K. D. Entian, and F. Götz. 1992. Genetic analysis of epidermin biosynthetic genes and epidermin-negative mutants of *Staphylococcus epidermidis*. *Eur. J. Biochem.* **204**:1149–1154.
4. Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A.

- Smith, and K. Struhl (ed.). 1990. Current protocols in molecular biology. John Wiley & Sons, New York, N.Y.
5. Bruckner, R. 1997. Gene replacement in *Staphylococcus carnosus* and *Staphylococcus xylosum*. FEMS Microbiol. Lett. **151**:1–8.
  6. Cetinkaya, Y., P. Falk, and C. G. Mayhall. 2000. Vancomycin-resistant enterococci. Clin. Microbiol. Rev. **13**:686–707.
  7. Chow, J. W., L. A. Thal, M. B. Perri, J. A. Vazquez, S. M. Donabedian, D. B. Clewell, and M. J. Zervos. 1993. Plasmid-associated hemolysin and aggregation substance production contribute to virulence in experimental enterococcal endocarditis. Antimicrob. Agents Chemother. **37**:2474–2477.
  8. Costerton, J. W., P. S. Stewart, and E. P. Greenberg. 1999. Bacterial biofilms: a common cause of persistent infections. Science **284**:1318–1322.
  9. Cucarella, C., C. Solano, J. Valle, B. Amorena, I. I. Lasa, and J. R. Penades. 2001. Bap, a *Staphylococcus aureus* surface protein involved in biofilm formation. J. Bacteriol. **183**:2888–2896.
  10. Eaton, T. J., and M. J. Gasson. 2001. Molecular screening of *Enterococcus* virulence determinants and potential for genetic exchange between food and medical isolates. Appl. Environ. Microbiol. **67**:1628–1635.
  11. Edmond, M. B., J. F. Ober, J. D. Dawson, D. L. Weinbaum, and R. P. Wenzel. 1996. Vancomycin-resistant enterococcal bacteremia: natural history and attributable mortality. Clin. Infect. Dis. **23**:1234–1239.
  12. Espinosa-Urgel, M., A. Salido, and J. L. Ramos. 2000. Genetic analysis of functions involved in adhesion of *Pseudomonas putida* to seeds. J. Bacteriol. **182**:2363–2369.
  13. Foley, L., and P. Gilbert. 1997. In-vitro studies of the activity of glycopeptide combinations against *Enterococcus faecalis* biofilms. J. Antimicrob. Chemother. **40**:667–672.
  14. Fournier, B., and D. C. Hooper. 2000. A new two-component regulatory system involved in adhesion, autolysis, and extracellular proteolytic activity of *Staphylococcus aureus*. J. Bacteriol. **182**:3955–3964.
  15. Fujimoto, S., and Y. Ike. 2001. pAM401-based shuttle vectors that enable overexpression of promoterless genes and one-step purification of tag fusion proteins directly from *Enterococcus faecalis*. Appl. Environ. Microbiol. **67**:1262–1267.
  16. Gilmore, M. S., R. A. Segarra, M. C. Booth, C. P. Bogie, L. R. Hall, and D. B. Clewell. 1994. Genetic structure of the *Enterococcus faecalis* plasmid pAD1-encoded cytolytic toxin system and its relationship to lantibiotic determinants. J. Bacteriol. **176**:7335–7344.
  17. Guzman, C. A., C. Pruzzo, G. LiPira, and L. Calegari. 1989. Role of adherence in pathogenesis of *Enterococcus faecalis* urinary tract infection and endocarditis. Infect. Immun. **57**:1834–1838.
  18. Hacker, J., and J. B. Kaper. 2000. Pathogenicity islands and the evolution of microbes. Annu. Rev. Microbiol. **54**:641–679.
  19. Heilmann, C., M. Hussain, G. Peters, and F. Gotz. 1997. Evidence for autolysin-mediated primary attachment of *Staphylococcus epidermidis* to a polystyrene surface. Mol. Microbiol. **24**:1013–1024.
  20. Heilmann, C., O. Schweitzer, C. Gerke, N. Vanittanakom, D. Mack, and F. Gotz. 1996. Molecular basis of intercellular adhesion in the biofilm-forming *Staphylococcus epidermidis*. Mol. Microbiol. **20**:1083–1091.
  21. Hunt, C. P. 1998. The emergence of enterococci as a cause of nosocomial infection. Br. J. Biomed. Sci. **55**:149–156.
  22. Huycke, M. M., and M. S. Gilmore. 1997. In vivo survival of *Enterococcus faecalis* is enhanced by extracellular superoxide production. Adv. Exp. Med. Biol. **418**:781–784.
  23. Huycke, M. M., W. A. Joyce, and M. F. Wack. 1996. Augmented production of extracellular superoxide production by blood isolates of *Enterococcus faecalis*. J. Infect. Dis. **173**:743–746.
  24. Huycke, M. M., D. F. Sahn, and M. S. Gilmore. 1998. Multiple-drug resistant enterococci: the nature of the problem and an agenda for the future. Emerg. Infect. Dis. **4**:239–249.
  25. Ike, Y., D. B. Clewell, R. A. Segarra, and M. S. Gilmore. 1990. Genetic analysis of the pAD1 hemolysin/bacteriocin determinant in *Enterococcus faecalis*: Tn917 insertional mutagenesis and cloning. J. Bacteriol. **172**:155–163.
  26. Ike, Y., H. Hashimoto, and D. B. Clewell. 1987. High incidence of hemolysin production by *Enterococcus (Streptococcus) faecalis* strains associated with human parenteral infections. J. Clin. Microbiol. **25**:1524–1528.
  27. Joyanes, P., A. Pascual, L. Martinez-Martinez, A. Hevia, and E. J. Perea. 2000. In vitro adherence of *Enterococcus faecalis* and *Enterococcus faecium* to urinary catheters. Eur. J. Clin. Microbiol. Infect. Dis. **19**:124–127.
  28. Kreft, B., R. Marre, U. Schramm, and R. Wirth. 1992. Aggregation substance of *Enterococcus faecalis* mediates adhesion to cultured renal tubular cells. Infect. Immun. **60**:25–30.
  29. Lewis, K. 2001. Riddle of biofilm resistance. Antimicrob. Agents Chemother. **45**:999–1007.
  30. Lowe, A. M., P. A. Lambert, and A. W. Smith. 1995. Cloning of an *Enterococcus faecalis* endocarditis antigen: homology with adhesins from some oral streptococci. Infect. Immun. **63**:703–706.
  31. Mengaud, J., S. Dramsi, E. Gouin, J. A. Vazquez-Boland, G. Milon, and P. Cossart. 1991. Pleiotropic control of *Listeria monocytogenes* virulence factors by a gene which is autoregulated. Mol. Microbiol. **5**:2273–2283.
  32. Moellering, R. C., Jr. 1992. Emergence of Enterococcus as a significant pathogen. Clin. Infect. Dis. **14**:1173–1176.
  33. Mundy, L. M., D. F. Sahn, and M. Gilmore. 2000. Relationships between enterococcal virulence and antimicrobial resistance. Clin. Microbiol. Rev. **13**:513–522.
  34. Nallapareddy, S. R., X. Qin, G. M. Weinstock, M. Hook, and B. E. Murray. 2000. *Enterococcus faecalis* adhesin, Ace, mediates attachment to extracellular matrix proteins collagen type IV and laminin as well as collagen type I. Infect. Immun. **68**:5218–5224.
  35. Olmsted, S. B., G. M. Dunny, S. L. Erlandsen, and C. L. Wells. 1994. A plasmid-encoded surface protein on *Enterococcus faecalis* augments its internalization by cultured intestinal epithelial cells. J. Infect. Dis. **170**:1549–1556.
  36. O'Toole, G., H. B. Kaplan, and R. Kolter. 2000. Biofilm formation as microbial development. Annu. Rev. Microbiol. **54**:49–79.
  37. O'Toole, G. A., and R. Kolter. 1998. Initiation of biofilm formation in *Pseudomonas fluorescens* WCS365 proceeds via multiple, convergent signaling pathways: a genetic analysis. Mol. Microbiol. **28**:449–461.
  38. Patti, J. M., and M. Hook. 1994. Microbial adhesins recognizing extracellular matrix macromolecules. Curr. Opin. Cell Biol. **6**:752–758.
  39. Pratt, L. A., and R. Kolter. 1998. Genetic analysis of *Escherichia coli* biofilm formation: roles of flagella, motility, chemotaxis and type I pili. Mol. Microbiol. **30**:285–293.
  40. Rosenberg, M., D. Gutnick, and E. Rosenberg. 1980. Adherence of bacteria to hydrocarbons: a simple method for measuring cell-surface hydrophobicity. FEMS Microbiol. Lett. **9**:29–33.
  41. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
  42. Schumacher-Perdreau, F., C. Heilmann, G. Peters, F. Gotz, and G. Pulverer. 1994. Comparative analysis of a biofilm-forming *Staphylococcus epidermidis* strain and its adhesion-positive, accumulation-negative mutant M7. FEMS Microbiol. Lett. **117**:71–78.
  43. Shankar, V., A. S. Baghdayan, M. M. Huycke, G. Lindahl, and M. S. Gilmore. 1999. Infection-derived *Enterococcus faecalis* strains are enriched in *esp*, a gene encoding a novel surface protein. Infect. Immun. **67**:193–200.
  44. Shankar, V., and M. S. Gilmore. 1997. Characterization of the *Enterococcus faecalis* alpha C protein homolog. Evidence for the expression of alternate forms in commensal and infection derived isolates. Adv. Exp. Med. Biol. **418**:1045–1048.
  45. Tomita, H., S. Fujimoto, K. Tanimoto, and Y. Ike. 1997. Cloning and genetic and sequence analyses of the bacteriocin 21 determinant encoded on the *Enterococcus faecalis* pheromone-responsive conjugative plasmid pPD1. J. Bacteriol. **179**:7843–7855.
  46. Veenstra, G. J., F. F. Cremers, H. van Dijk, and A. Fleer. 1996. Ultrastructural organization and regulation of a biomaterial adhesin of *Staphylococcus epidermidis*. J. Bacteriol. **178**:537–541.
  47. Woodford, N. 1998. Glycopeptide-resistant enterococci: a decade of experience. J. Med. Microbiol. **47**:849–862.
  48. Xu, Y., K. V. Singh, X. Qin, B. E. Murray, and G. M. Weinstock. 2000. Analysis of a gene cluster of *Enterococcus faecalis* involved in polysaccharide biosynthesis. Infect. Immun. **68**:815–823.