

Determination of Oligopeptide Diversity within a Natural Population of *Microcystis* spp. (Cyanobacteria) by Typing Single Colonies by Matrix-Assisted Laser Desorption Ionization–Time of Flight Mass Spectrometry

JUTTA FASTNER,^{1*} MARCEL ERHARD,² AND HANS VON DÖHREN¹

Biotechnology Center and Max Vollmer Institute, Technical University Berlin, 10587 Berlin,¹ and AnagnosTec GmbH, 14943 Luckenwalde,² Germany

Received 23 May 2001/Accepted 28 August 2001

Besides the most prominent peptide toxin, microcystin, the cyanobacteria *Microcystis* spp. have been shown to produce a large variety of other bioactive oligopeptides. We investigated for the first time the oligopeptide diversity within a natural *Microcystis* population by analyzing single colonies directly with matrix-assisted laser desorption ionization–time of flight mass spectrometry (MALDI-TOF MS). The results demonstrate a high diversity of known cyanobacterial peptides such as microcystins, anabaenopeptins, microginins, aeruginosins, and cyanopeptolins, but also many unknown substances in the *Microcystis* colonies. Oligopeptide patterns were mostly related to specific *Microcystis* taxa. *Microcystis aeruginosa* (Kütz.) Kütz. colonies contained mainly microcystins, occasionally accompanied by aeruginosins. In contrast, microcystins were not detected in *Microcystis ichthyoblabe* Kütz.; instead, colonies of this species contained anabaenopeptins and/or microginins or unknown peptides. Within a third group, *Microcystis wesenbergii* (Kom.) Kom. in Kondr., chiefly a cyanopeptolin and an unknown peptide were found. Similar patterns, however, were also found in colonies which could not be identified to species level. The significance of oligopeptides as a chemotaxonomic tool within the genus *Microcystis* is discussed. It could be demonstrated that the typing of single colonies by MALDI-TOF MS may be a valuable tool for ecological studies of the genus *Microcystis* as well as in early warning of toxic cyanobacterial blooms.

Freshwater and marine cyanobacteria are known to produce a variety of bioactive compounds, among them potent hepatotoxins and neurotoxins (for an overview, see reference 45). Many of the toxic species of cyanobacteria tend to massive proliferation in eutrophicated water bodies and thus have been the cause for considerable hazards for animal and human health (3, 23). One of the most widespread bloom-forming cyanobacteria is the genus *Microcystis*, a well-known producer of the hepatotoxic peptide microcystin (45). Microcystins are a group of closely related cyclic heptapeptides sharing the common structure cyclo(D-Ala-L-X-D-MeAsp-L-Z-Adda-D-Glu-Mdha), in which MeAsp is D-erythro- β -methylaspartic acid, Mdha is N-methyldehydroalanine, Adda is 2S,3S,8S,9S-3-amino-9-methoxy-2,6,8-trimethyl-10-phenyldeca-4E,6E-dienoic acid, and X and Z are variable L-amino acids, e.g., microcystin-LR (MC-LR) contains leucine (L) and arginine (R) (5). So far, more than 60 derivatives of microcystins have been identified, varying largely by the degree of methylation, peptide sequence, and toxicity (for an overview, see reference 45).

The hepatotoxicity of microcystins is based on their inhibition of protein phosphatases 1 and 2A in combination with transport into hepatocytes via the bile acid carrier, leading to acute liver failure due to the disruption of hepatocyte cytoskeletal components (11, 26). The widespread occurrence and

acute toxicity of microcystins and their tumor-promoting properties imply the need for identification and prediction of toxic blooms (23).

The traditional botanical code describes the genus *Microcystis* as a coccal, unicellular cyanobacterium that grows as mucilaginous colonies of irregularly arranged cells (under natural conditions, while strain cultures usually grow as single cells). According to this tradition, morphological criteria such as size of the individual cells, colony morphology, and mucilage characteristics are used for species delimitation within *Microcystis* (i.e., morphospecies) (20, 21). Microcystin-producing strains as well as strains that do not synthesize microcystin have been reported for all species within the genus *Microcystis*. However, whereas most field samples and strains of *Microcystis aeruginosa* and *Microcystis viridis* studied to date were found to contain microcystins (17, 47, 49–51), strains of *M. wesenbergii*, *M. novaceckii*, and *M. ichthyoblabe* have only sporadically been reported to contain microcystins (34, 38, 49).

Beside microcystins, various other linear and cyclic oligopeptides such as aeruginosins, anabaenopeptilides, cyanopeptolins, anabaenopeptins, and microginins are found within the genus *Microcystis* (31). Similar to microcystins, these peptides possess unusual amino acids like 3-amino-6-hydroxy-2-piperidone (Ahp) in cyanopeptolins or 2-carboxy-6-hydroxyoctahydroindol (Choi) in aeruginosin-type molecules, and numerous structural variants also exist within these groups (14, 29, 31). These peptides show diverse bioactivities, frequently protease inhibition (31).

The presence of D-amino acids, unusual amino acids, as well

* Corresponding author. Mailing address: Federal Environmental Agency, Corrensplatz 1, 14195 Berlin, Germany. Phone: 49 30 8903 1390. Fax: 49 30 8903 1830. E-mail: jutta.fastner@uba.de.

TABLE 1. Colony characteristics and cell size of *Microcystis* species isolated from Lake Wannsee in 1999

Species	Colony characteristics	Mean cell diam (μm) ± SD	Reference
<i>M. aeruginosa</i> (Kützing) Kützing	More or less firm, elongated and lobate, usually with distinct holes, net-like clathrate; often more or less dense epiphytic populations of <i>Pseudanabaena muscicola</i> or <i>Nitzschia</i> spp.	4.92 ± 0.13	21
<i>M. ichthyoblabe</i> Kützing	Colonies more or less irregular-spherical, soft; distribution of cells in colonies is irregular or sponge like; margin of colony irregular	3.39 ± 0.18	21
<i>M. wesenbergii</i> (Komárek) Komárek in Kondrateva	Elongated, lobate cells in one layer, with visible margin of mucilage	5.29 ± 0.24	21

as their small size suggests that the cyanobacterial oligopeptides mentioned above are synthesized nonribosomally by multifunctional enzyme complexes, generally termed peptide synthetases, a pathway studied intensively in other bacteria and fungi (1, 19). The nonribosomal synthesis of microcystins in the axenic strain *Microcystis* sp. strain PCC 7806 and of anabaenopeptilides in *Anabaena* sp. strain 90 was recently demonstrated by site-directed mutagenesis and sequencing (6, 42, 46). Nonribosomal peptide synthetase genes have so far been detected in all strains of the genus *Microcystis*, but genes encoding for the so-called microcystin synthetase are usually detected only in toxic (i.e., microcystin-containing) *Microcystis* spp. (7, 35). This corresponds to the observation of oligopeptides in all *Microcystis* strains investigated to date showing various combinations of microcystins and/or other oligopeptides such as aeruginosins, cyanopeptolins, or anabaenopeptins (8, 27, 31).

The cooccurrence of both microcystins and other oligopeptides such as anabaenopeptins and cyanopeptolins in natural *Microcystis* populations was recently demonstrated (10, 14, 36). It is well known that the species and genotype composition in natural *Microcystis* populations is heterogeneous, and both microcystin- and non-microcystin-containing strains have been isolated from the same sample (41, 48, 52). Rohrlack et al. (41) isolated 13 *Microcystis* strains from Lake Wannsee (Berlin, Germany) in 1995 which produced either microcystins or anabaenopeptins (T. Rohrlack, M. Erhard, and M. Henning, unpublished data). Furthermore, isolated strains may show both a different qualitative and quantitative microcystin pattern than the original population (41, 48). These results suggest a considerable diversity of genotypes with different oligopeptide patterns in natural *Microcystis* populations.

Our study aimed to investigate the inter- and intraspecific oligopeptide diversity in a natural population of the genus *Microcystis*. Since isolation of strains from natural populations is likely to be selective, we recorded the oligopeptide pattern directly in single *Microcystis* colonies selected from natural populations using matrix-assisted laser desorption/ionization-time of flight mass spectrometry (MALDI-TOF MS).

MATERIALS AND METHODS

Sampling and isolation. *Microcystis* spp. were harvested biweekly in hypertrophic Lake Wannsee, Berlin, Germany, from June to October 1999 with a plankton net (40-μm mesh size). The samples were stored in the cool and dark until isolation of *Microcystis* colonies the same day. The colonies were isolated by serial dilution with tap water and micromanipulation techniques using an inverted microscope at a magnification of ×160 to ×400. Isolated colonies were

washed by transferring them to several drops of water until all other organisms were removed.

Epiphytic cyanobacteria and algae sticking in the mucilage of *Microcystis aeruginosa* could not be detached (Table 1). Cell size and morphological characteristics were recorded for each colony and species were determined according to Komárek and Anagnostidis (21) (Table 1). Additionally, aliquots of the concentrated net samples were taken and either frozen at -20°C and lyophilized or fixed with formaldehyde solution and stored in the dark for detailed cell size determination. The mean cell diameter of the species was determined by measuring the diameters of 50 cells (10 cells per colony) every sampling day.

Extraction and preparation of *Microcystis* colonies, lyophilized strains, and field samples for MALDI-TOF MS analysis. Single colonies were directly transferred onto a stainless steel template, and immediately 1 μl of matrix (10 mg of 2,5-dihydroxybenzoic acid per ml in water-acetonitrile [1:1] with 0.03% trifluoroacetic acid) was added. The extraction of the oligopeptides from the cells was achieved by the solvent fraction of the matrix. An immediate change of the colony color from green to brownish yellow after addition of the matrix was observed due to the degradation of chlorophyll *a* by the acidic solution.

Lyophilized *Microcystis* field samples, the axenic *Microcystis* strains PCC 7806 and PCC 7813, and the unialgal *Microcystis* strains HUB 5-2-4, HUB 5-3, and HUB 063 were extracted with acetonitrile-ethanol-water (1:1:1) with 0.03% trifluoroacetic acid. Then 1 μl of the extract was prepared for MALDI-TOF MS analysis as described for the single colonies.

MALDI-TOF MS analysis. Positive ion mass spectra were recorded from each colony and the lyophilized field samples and strains using a MALDI-TOF mass spectrometer (Voyager DE-PRO; PerSeptive BioSystems, Framingham, Mass.) equipped with a reflectron. For desorption of the components, a nitrogen laser beam (λ = 337 nm) was focused on the template. The acceleration voltage was set at 20 kV. All measurements were carried out in the delayed extraction mode, allowing the determination of monoisotopic mass values (*m/z*; mass-to-charge ratio). Analyses were performed in the positive-ion mode, giving mainly singly protonated molecular ions ([M+H]⁺). Chlorophyll *a* degradation products pheophytin *a* and pheophorbide *a* with mass values of *m/z* 871.57 and 593.27 [M+H]⁺, respectively, were used for internal calibration. A low mass gate of 500 Da improved measurement by filtering out the most intensive matrix ions.

After determination of monoisotopic mass values, post-source decay (PSD) measurements for recording fragment ions were performed directly from the same colony on the template. The precursor ions were selected with a time ion selector having a mass window of 10 mass units. The operating voltages of the reflectron were reduced stepwise to record 12 spectral segments sequentially.

PSD spectra of the most prominent peptides were recorded several times over the entire sampling period from single colonies and additionally from the lyophilized *Microcystis* samples.

RESULTS

Species determination. The *Microcystis* population in Lake Wannsee in 1999 consisted of several species which were present during the entire investigation period. The majority of the isolated colonies belonged to one of three species with distinct colonial features and cell sizes (Table 1): *M. aeruginosa* (Kützing) Kützing, *M. ichthyoblabe* Kützing, and *M. wesenbergii* (Komárek) Komárek in Kondrateva. The other colonies isolated could not be unequivocally determined to species level




FIG. 1. Positive-ion MALDI-TOF mass spectra (m/z range 550 to 1,100 Da) of *Microcystis* strains PCC 7806 (axenic) (A), PCC 7813 (axenic) (B), HUB 5-3 (unialgal) (C), HUB 5-2-4 (unialgal) (D), and HUB 063 (unialgal) (E). For structure assignments, see Table 2.

and thus were grouped as *Microcystis* spp. About half of these colonies exhibited colonial characteristics similar to those of *Microcystis ichthyoblabe* but had larger cell sizes than the main phenotype isolated from this species. The other colonies showed diverse variations in cell size and colonial characteristics, often similar to features described for *M. flos-aquae* or *M. novaceckii* (Komárek) Compère. The sizes of the colonies isolated ranged between 0.2 and 4 mm, as measured at their longest dimension.

Oligopeptides identified by MALDI-TOF MS. Positive-ion mass spectra were recorded from the *Microcystis* strains, from 258 *Microcystis* colonies, and from the entire *Microcystis* population each time colonies were isolated (Fig. 1 and 2). In the mass range of m/z 500 to 1,100 Da, numerous structural variants of microcystins, anabaenopeptins, microginins, aeruginosins, and one cyanopeptolin were identified by means of characteristic fragment ions obtained by PSD measurements (Table 2). Most of these components are well known, and data about their structures and PSD data have been published previously (9, 10, 12) (Table 2). As an example, the PSD spectrum of microcystin-LR with characteristic fragment ions leading to

an unambiguous structure assignment is given in Fig. 3 (32). Fragment analysis assigned peptides with mass values of m/z 726, 728, 740, 742, 749, and 751 $[M+H]^+$ as microginin variants previously identified by means of amino acid analysis and PSD measurements (Table 2) (U. Neumann, J. Weckesser, and M. Erhard, unpublished data). The compound with a molecular mass of m/z 603 $[M+H]^+$ is probably a new variant of an aeruginosin-type peptide, as suggested by the fragment ion of m/z 140, indicating the presence of the unusual amino acid Choi, which is unique to aeruginosin-type molecules (8, 28, 29).

In addition, many unknown substances were detected in the colonies, of which those with mass values of m/z 619, 635, 771, 787, 804, 846, 1,007, 1,009, 1,015, and 1,021 were the most abundant.

Inter- and intraspecific oligopeptide diversity. Mass spectra from the whole *Microcystis* population in Lake Wannsee in 1999 showed a complex mixture of different microcystins, microginins, anabaenopeptins, and unknown components (Fig. 2A). In contrast, isolated *Microcystis* strains usually have a less diverse peptide pattern. As shown in Fig. 1, the axenic *Microcystis* strains PCC 7806 and PCC 7813 contain largely MC-LR and $[Asp^3]$ MC-LR and either cyanopeptolin D or the unknown peptide of m/z 603 $[M+H]^+$ (Fig. 1A and B; Table 2) (27). Microcystins are the most abundant oligopeptides in the unialgal strain HUB 5-2-4, while in HUB 5-3 an unknown peptide of m/z 804 $[M+H]^+$ and in HUB 063 anabaenopeptin B and F are found (Fig. 1C to E).

Similar to the clonal strains, the peptide patterns in the single *Microcystis* colonies were usually less complex (Fig. 2B to D). Comparison of the peptide composition in the total *Microcystis* population with that of single colonies suggests that the population is dominated by colonies with the specific oligopeptide patterns shown in Fig. 2B to D.

The oligopeptide patterns in colonies of *M. aeruginosa*, *M. ichthyoblabe*, and *M. wesenbergii* revealed pronounced differences (Fig. 2B to D and Fig. 4). In all but 2 of 111 *M. aeruginosa* colonies, microcystins were the chief oligopeptides detected (Fig. 2B, Fig. 4). The microcystin profiles within this species were rather homogeneous, with MC-RR, MC-YR, and MC-LR being codominant in most colonies. Similar microcystin compositions were found both by MALDI-TOF MS (Fig. 2A) and by high-pressure liquid chromatography (HPLC) analysis (data not shown) of the whole population in Lake Wannsee. MC-RR was detected in 79%, MC-YR in 89%, and MC-LR in 94% of all *M. aeruginosa* colonies. Minor microcystins, as indicated by low peak intensities and small amounts in HPLC analysis, found in *M. aeruginosa* were $[Dha^7]$ MC-RR, $[Dha^7]$ MC-LR, $[H_4]$ MC-YR, and MC-WR. They were detected in 13 to 37% of all colonies. In some colonies of *M. aeruginosa*, only $[Dha^7]$ MC-RR and $[Dha^7]$ MC-LR were found. In addition to microcystins, aeruginosins were present in some colonies, and in four colonies anabaenopeptins could be detected (Fig. 4). Unknown components were detected in many *M. aeruginosa* colonies, occasionally with molecular weights corresponding to those of known microcystin variants. However, signal intensity was too low to obtain reliable PSD spectra.

In order to control for the homogeneity of the colonies, eight large colonies of *M. aeruginosa* were divided into two to four parts, and each part was analyzed separately. No differ-




FIG. 2. Positive-ion MALDI-TOF mass spectra (m/z range, 600 to 1,100 Da) of the entire *Microcystis* population (A) and of colonies of *M. aeruginosa* (B), *M. ichthyoblabe* (C), and *Microcystis* sp. (D) in Lake Wannsee on 24 August 1999. For structure assignments, see Table 2.

ence in the microcystin pattern was found between the single parts (data not shown). Furthermore, no relationship between the presence of epiphytic cyanobacteria and algae in the *M. aeruginosa* colonies and their peptide pattern was found.

By contrast, microcystins were never detected in colonies of *M. ichthyoblabe* (Fig. 2C, Fig. 4). The oligopeptide pattern within this group was more diverse: they contained either mainly anabaenopeptins or both microginins and anabaenopeptins or various unknown peptides. Anabaenopeptins B and F and oscillamide Y were the most prominent anabaenopeptins, while anabaenopeptins I and A were less abundant. Major microginins were FR5 and FR3, while all other variants of this substance class were less frequent. Aeruginosins were also detected in the colonies of *M. ichthyoblabe* (Fig. 4).

M. wesenbergii colonies also did not contain microcystins. The colonies of this species show very similar patterns, with 11 of 14 colonies containing cyanopeptolin-S together with an unknown peptide of m/z 635 $[M+H]^+$ (Fig. 4).

Oligopeptide patterns of *Microcystis* colonies not identified to species level (*Microcystis* spp.) mainly fall into three clusters,

with peptide patterns similar to those observed for *M. aeruginosa* and *M. ichthyoblabe* (Fig. 5). The colonies contained either microcystins, microginin and/or anabaenopeptin, or chiefly unknown peptides (Fig. 2D, Fig. 5). Aeruginosins were detected in all three clusters.

DISCUSSION

MALDI-TOF MS analysis of the *Microcystis* populations from Lake Wannsee in 1995 to 1999 showed a complex and persisting mixture of microcystins, other oligopeptides, and unknown components (8, 10) (Fig. 2A). The cooccurrence of microcystins and cyanopeptolins in other *Microcystis* sp.-dominated field samples was reported previously (15, 36). By typing single *Microcystis* colonies, we could show for the first time that the actual peptide diversity in a natural population of this genus is substantially higher. Many of the substances detected belong to well-known groups of cyanobacterial peptides like microcystins, anabaenopeptins, microginins, cyanopeptolins, and aeruginosins, of which many have been discovered in *Microcystis* spp. (31). In addition, numerous unknown compo-

TABLE 2. Oligopeptides detected in *Microcystis* colonies from Lake Wannsee in 1999 using MALDI-TOF MS (results from positive-ion mass spectra and PSD measurements)

Substance class	[M+H] ⁺	Assignment	Reference(s)
Microcystin	981	[Dha ⁷] MC-LR	32, 12
	995	MC-LR	32, 12
	1,024	[Dha ⁷] MC-RR	18, 12
	1,038	MC-RR	32, 12
	1,045	MC-YR	32, 12
	1,049	[H ₄] MC-YR	33, 12
	1,068	MC-WR	32, 12
Anabaenopeptin	821	Anabaenopeptin I	30
	837	Anabaenopeptin B	13, 10
	844	Anabaenopeptin A	13
	851	Anabaenopeptin F	44, 9
	858	Oscillamide Y	43
Microginin	726	Microginin FR5	PSD fragmentation; Neumann et al., unpublished data
	728	Microginin FR3	See microginin FR5
	740	Microginin FR6	See microginin FR5
	742	Microginin FR2	See microginin FR5
	749	Microginin FR10	See microginin FR5
	751	Microginin FR9	See microginin FR5
Aeruginosin	561	Aeruginosamide	25
	603	Aeruginosin-type	PSD fragmentation
	609	Microcin SF608	2
	653 ^a	Aeruginosin 102A	28, 8
Cyanopeptolin	846 ^b /828 ^c	Cyanopeptolin-S	14, 8
	957 ^b /939 ^c	Cyanopeptolin D	27

^a [M - SO₃ + H]⁺.

^b [M - SO₃ + H]⁺.

^c [M - SO₃ - H₂O + H]⁺.

nents have been detected in the colonies. However, the origin of these unknown components has yet to be investigated, since besides the observed epiphytic cyanobacteria and algae, heterotrophic bacteria are also known to be present in *Microcystis* colonies (4).

Usually more than one type of oligopeptide was detected in the *Microcystis* colonies from Lake Wannsee in 1999. With respect to known peptides, combinations of anabaenopeptins, microginins, and aeruginosins were observed, while microcystins were found along with aeruginosins. This correlates to the detection of aeruginosins as well as cyanopeptolins in both toxic and nontoxic *Microcystis* culture strains (Fig. 1) (6, 31). Anabaenopeptins and microginins were usually not detected together with microcystins with the exception of four colonies

containing both microcystins and anabaenopeptins. Microcystins and anabaenopeptins were also never found simultaneously in more than 20 *Microcystis* strains investigated to date (M. Erhard, H. von Döhren, P. Jungblut, E. Dittmann, T. Börner, M. Henning, L. Rouhiainen, and K. Sivonen, Abstracts of the 4th European Workshop on the Molecular Biology of Cyanobacteria, p. 29, 1999). However, it must be considered that isolation of strains is selective and may pick up only those genotypes which are favored by the cultivation conditions. On the other hand, although one colony can be regarded as a clone and homogeneity was found for all of the divided colonies, a potential bias of conducting studies with selected colonies may be the contamination of a colony with cells of other clones.

Our data revealed a relationship between oligopeptide patterns and certain *Microcystis* taxa in Lake Wannsee. Microcystins were chiefly found in *M. aeruginosa*, while colonies of *M. ichthyoblabe* and *M. wesenbergii* did not contain microcystins but did contain anabaenopeptins, microginins, cyanopeptolins, or unknown peptides. Essentially the same results were found for the presence of the microcystin synthetase genes in single *Microcystis* colonies from Lake Wannsee in 2000 (24); microcystin synthetase genes were detected in 73% of *M. aeruginosa* colonies but in only 16% of colonies assigned to *M. ichthyoblabe* and in 0% of *M. wesenbergii* colonies. The low oligopeptide diversity within *M. aeruginosa* suggests that the species-related peptide patterns observed may have been caused by the dominance of only certain genotypes within the *Microcystis* species in Lake Wannsee in 1999. Restriction fragment length polymorphism of the *mcyB* gene indicates the presence of five different genotypes among *M. aeruginosa* colonies with similar microcystin profiles in Lake Wannsee in 2000 (24). The occur-




FIG. 3. PSD spectrum of microcystin-LR: m/z 995 [M + H]⁺, 861 [M - 134 (Adda side chain) + H]⁺, 599 [Arg-Adda-Glu + H]⁺, 375 [C₁₁H₁₄O-Glu-Mdha]⁺, 286 [Arg-MeAsp + H]⁺, 213 [Glu-Mdha + H]⁺, 155 [Mdha-Ala + H]⁺, 135 [PhCH₂CH(OCH₃)⁺, and 70 [Leu - CO + H]⁺.




FIG. 4. Molecular masses (m/z) oligopeptides and unknown components in the range of m/z 500 to 1,100 Da detected in colonies of *M. aeruginosa*, *M. ichthyoblabe*, and *M. wesenbergii*. Symbols for substance classes: open diamonds, aeruginosin; open circles, microginin; open triangles, anabaenopeptin; open squares, microcystin; solid diamonds, cyanopeptolin; small solid rectangles, unknown components. For structure assignments of compounds, see Table 2.

rence of various combinations of several types of oligopeptides in *M. ichthyoblabe* implies a higher genotype diversity within this species in Lake Wannsee in 1999.

M. aeruginosa is worldwide the species most often associated with toxic water blooms and microcystin-producing strains (38, 47, 50), while the majority of *M. wesenbergii* and *M. ichthyoblabe* strains reported in the literature did not produce microcystin, although microcystin-producing strains are occasionally described (34, 38, 49). Japanese strains of a *M. aeruginosa*

S-type (i.e., small cell size), which were later classified as *M. ichthyoblabe* (51), also rarely contained microcystins (50). Rohrlack et al. (41) isolated 13 *Microcystis* strains from Lake Wannsee in 1995, of which the strains having larger cells contained microcystins, while those with smaller cells produced only anabaenopeptins (T. Rohrlack, M. Erhard, and M. Henning, unpublished data). Though these data support some relationship between morphospecies and microcystin production, we also detected the peptide patterns typical for *M.*




FIG. 5. Molecular masses (m/z) of oligopeptides and unknown components in the range of m/z 500 to 1,100 Da detected in colonies of *Microcystis* spp. Symbols for substance classes: open diamonds, aeruginosin; open circles, microginin; open triangles, anabaenopeptin; open squares, microcystin; solid diamonds, cyanopeptolin; small solid rectangles, unknown components. For structure assignments of compounds, see Table 2.

aeruginosa and *M. ichthyoblabe* in colonies with colonial characteristics different from these species (these were grouped as *Microcystis* spp.).

However, colony morphology and cell size, traditionally used in taxonomic differentiation of the genus *Microcystis* (20, 21), may be questionable criteria for species distinction; both parameters have been found to be variable in laboratory cultures and in the field (22, 39, 40). In culture particularly some strains of *M. aeruginosa*, *M. ichthyoblabe*, and *M. novaceckii* have developed colony morphologies similar to each other (39). Recent studies investigating the taxonomy of the genus *Microcystis* using genetic criteria in comparison to morphological traits and microcystin production show contradictory results. 16S rRNA analysis revealed no differences between different toxic and nontoxic strains of *M. aeruginosa*, *M. wesenbergii*, and *M. viridis* (34). In contrast, sequence data for the 16S to 23S internally transcribed spacer lead to three clusters, of which cluster I contained both toxic and nontoxic strains of *M. aeruginosa*, *M. novaceckii*, and *M. ichthyoblabe*, cluster II only toxic strains of mainly *M. viridis*, and cluster III only nontoxic strains of mainly *M. wesenbergii* (38). Similarly, allozyme divergence studied by Kato et al. (16) characterized *M. wesenbergii* and *M. viridis* as well-established species. In contrast to Otsuka et al. (38), allozyme divergence (16) revealed a separation of Japanese *M. aeruginosa* L-type strains (equivalent to *M. aeruginosa*) from strains of *M. aeruginosa* type S (included in *M. ichthyo-*

blabe [51]), which corresponds to our observation of a different peptide pattern in these species. More data are needed to determine whether or not the oligopeptide pattern may be used as a chemotaxonomical feature to clarify the taxonomic uncertainties within the genus *Microcystis*. These should include systematic studies on the distribution of oligopeptides in combination with morphological and molecular characterization of more strains and original colonies from different regions.

Our study demonstrates the unequivocal identification of microcystins and other oligopeptides in single *Microcystis* colonies by employing MALDI-TOF MS. It is thus possible to directly identify the toxic species and genotypes in natural *Microcystis* populations without time-consuming and probably selective isolation procedures. The typing of single *Microcystis* colonies may be a valuable tool in early warning of toxic bloom formation, since it enables rapid detection of whether or not a population contains microcystin-producing genotypes in an early phase of population growth. Furthermore, the succession of toxic and nontoxic species may be followed and the influence of biotic and abiotic factors on genotype succession assessed. In Lake Wannsee in 1999, *M. aeruginosa* colonies showed a persisting composition of MC-RR, MC-YR, and MC-LR over the entire sampling period from June to October. Similar quantitative relations of microcystins were determined in the whole *Microcystis* population by both HPLC and MALDI-TOF MS during this time span. This indicates that this type of *M. aeruginosa* colony determined the overall microcystin pattern in 1999.

Hypotheses about possible functions of microcystins often focused on the toxicity of microcystins. Although microcystins have been shown to affect diverse aquatic organisms (for an overview, see reference 45), they do not necessarily seem to be produced as a defense mechanism against zooplankton grazing. Speculations about an inter- or intracellular function of microcystins raise the question about substances playing a similar role in genotypes without microcystins (37). This is supported by our observation of various oligopeptides in the *Microcystis* population investigated. The coexistence of genotypes producing either mainly microcystins or other oligopeptides throughout the investigation period suggests that a comprehensive understanding of their possible functions and ecological benefits requires studying oligopeptides as a group rather than focusing on microcystins.

ACKNOWLEDGMENTS

We thank Frank Grützner and Adam Antebi and his group (Max Plank Institute for Molecular Genetics, Berlin, Germany) for kindly providing micromanipulation facilities. *Microcystis* strains were kindly provided by Rosemarie Rippka (Institute Pasteur, Paris, France) and Manfred Henning (Humboldt University, Berlin, Germany).

This work was financially supported by funds from the EU (ENV4-CT98-802).

REFERENCES

- Arment, A. R., and W. W. Carmichael. 1996. Evidence that microcystin is a thiotemplate product. *J. Phycol.* **32**:591–597.
- Banker, R., and S. Carmeli. 1999. Inhibitors of serine proteases from a waterbloom of the cyanobacterium *Microcystis* sp. *Tetrahedron* **55**:10835–10844.
- Bell, S. G., and G. A. Codd. 1994. Cyanobacterial toxins and human health. *Rev. Med. Microbiol.* **5**:256–264.
- Brunberg, A. K. 1999. Contribution of bacteria in the mucilage of *Microcystis*

- spp. (cyanobacteria) to benthic and pelagic bacterial production in a hypertrophic lake. *FEMS Microbiol. Lett.* **29**:13–22.
5. Carmichael, W. W., V. Beasley, D. L. Bunner, J. N. Eloff, I. Falconer, P. Gorham, K. I. Harada, T. Krishnamurthy, Y. Min-Juan, R. E. Moore, K. Rinehart, M. Runnegar, O. M. Skulberg, and M. Watanabe. 1988. Naming cyclic heptapeptide toxins of cyanobacteria (blue-green algae). *Toxicon* **26**: 971–973.
 6. Dittmann, E., B. A. Neilan, M. Erhard, H. von Döhren, and T. Börner. 1997. Insertional mutagenesis of a peptide synthetase gene that is responsible for hepatotoxin production in the cyanobacterium *Microcystis aeruginosa* PCC 7806. *Mol. Microbiol.* **26**:779–787.
 7. Dittmann, E., G. Christiansen, B. A. Neilan, J. Fastner, R. Rippka, and T. Börner. 1999. Peptide synthetases genes occur in various species of cyanobacteria, p. 615–621. In G. A. Peschek, W. Loeffelhardt, and G. Schmetterer (ed.), *The phototrophic prokaryotes*. Kluwer Academic/Plenum Publishing Corp., New York, N.Y.
 8. Erhard, M. 1999. Ph.D. thesis. Technische Universität Berlin, Berlin, Germany.
 9. Erhard, M., H. von Döhren, and P. Jungblut. 1997. Rapid typing and elucidation of new secondary metabolites of intact cyanobacteria using MALDI-TOF mass spectrometry. *Nat. Biotechnol.* **15**:906–909.
 10. Erhard, M., H. von Döhren, and P. Jungblut. 1999. Rapid identification of the new anabaenopeptin G from *Planktothrix agardhii* HUB 011 using matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *Rapid Commun. Mass Spectrom.* **13**:337–343.
 11. Falconer, I. R. 1998. Algal toxins and human health, p. 53–82. In J. Hrubec (ed.), *The handbook of environmental chemistry*, vol. 5, part C: quality and treatment of drinking water II. Springer-Verlag, Berlin, Germany.
 12. Fastner, J., M. Erhard, W. W. Carmichael, F. Sun, K. L. Rinehart, H. Rönické, and I. Chorus. 1999. Characterization and diversity of microcystins in natural blooms and strains of the genera *Microcystis* and *Planktothrix* from German freshwaters. *Arch. Hydrobiol.* **145**:147–163.
 13. Harada, K.-I., K. Fujii, T. Shimada, M. Suzuki, H. Sano, K. Adachi, and W. W. Carmichael. 1995. Two cyclic peptides, anabaenopeptins, a third group of bioactive compounds from the cyanobacterium *Anabaena flos-aquae* NRC 525-167. *Tetrahedron Lett.* **36**:1511–1514.
 14. Jacobi, C., L. Oberer, C. Quiquerez, W. A. König, and J. Weckesser. 1995. Cyanopeptolin S, a sulfate containing depsipeptide from a water-bloom of *Microcystis aeruginosa*. *FEMS Microbiol. Lett.* **129**:129–134.
 15. Jacobi, C., K. L. Rinehart, G. A. Codd, I. Carmienke and J. Weckesser. 1996. Occurrence of toxic water blooms containing microcystins in a German lake over a three year period. *J. Syst. Appl. Microbiol.* **19**:249–254.
 16. Kato, T., M. F. Watanabe, and M. Watanabe. 1991. Allozyme divergence in *Microcystis* (Cyanophyceae) and its taxonomic inference. *Algol. Studies* **64**: 129–140.
 17. Kaya, K., and M. M. Watanabe. 1990. Microcystin composition of an axenic clonal strain of *Microcystis viridis* and *Microcystis viridis*-containing waterblooms in Japanese freshwaters. *J. Appl. Phycol.* **2**:173–178.
 18. Kiviranta, J., M. Namikoshi, K. Sivonen, W. R. Evans, W. W. Carmichael, and K. L. Rinehart. 1992. Structure determination and toxicity of a new microcystin from *Microcystis aeruginosa* strain 205. *Toxicon* **30**:1093–1098.
 19. Kleinkauf, H., and H. von Döhren. 1996. A nonribosomal system of peptide biosynthesis. *Eur. J. Biochem.* **236**:335–351.
 20. Komárek, J. 1991. A review of water-bloom-forming *Microcystis* species, with regard to populations from Japan. *Algol. Studies* **64**:115–127.
 21. Komárek, J., and K. Anagnostidis. 1999. Cyanoprokaryota 1, Teil: Chroococcales. In H. Ettl, G. Gärtner, H. Heynig, and D. Mollenhauer (ed.), *Süßwasserflora von Mitteleuropa Band 19/1*. Gustav Fischer Verlag, Stuttgart, Germany.
 22. Krüger, G. H. J., J. N. Eloff, and J. A. Pretorius. 1991. Morphological changes in toxic and non-toxic *Microcystis* isolates at different irradiance levels. *J. Phycol.* **17**:52–56.
 23. Kuiper-Goodman, T., I. Falconer, and J. Fitzgerald. 1999. Human health aspects, p 113–153. In I. Chorus and J. Bartram (ed.), *Toxic cyanobacteria in water*. E & FN Spon, London, England.
 24. Kurmayer, R., E. Dittmann, J. Fastner, and I. Chorus. Diversity of microcystin genes within a population of the toxic cyanobacterium *Microcystis* spp. in Lake Wannsee. *Microb. Ecol.*, in press.
 25. Lawton, L. A., L. A. Morris, and M. Jaspers. 1999. A bioactive modified peptide, aeruginosamide, isolated from the cyanobacterium *Microcystis aeruginosa*. *J. Org. Chem.* **64**:5329–5332.
 26. MacKintosh, C., K. A. Beattie, S. Klump, P. Cohen, and G. A. Codd. 1990. Cyanobacterial microcystin-LR is a potent and specific inhibitor of protein phosphatases 1 and 2a from both mammals and higher plants. *FEBS Lett.* **264**:187–192.
 27. Martin, C., L. Oberer, T. Ino, W. A. König, M. Busch, and J. Weckesser. 1993. Cyanopeptolins, new depsipeptides from the cyanobacterium *Microcystis* PCC 7806. *J. Antibiot.* **46**:1550–1556.
 28. Matsuda, H., T. Okino, M. Murakami, and K. Yamaguchi. 1997. Aeruginosins 102-A and B, new thrombin inhibitors from the cyanobacterium *Microcystis viridis* (NIES-102). *Tetrahedron* **52**:14501–14506.
 29. Murakami, M., K. Ishida, T. Okino, Y. Okita, H. Matsuda, and K. Yamaguchi. 1995. Aeruginosin 98-A and-B, trypsin inhibitors from the blue-green alga *Microcystis aeruginosa* (NIES-98). *Tetrahedron Lett.* **36**:2758–2788.
 30. Murakami, M., S. Suzuki, Y. Itou, S. Kodani, and K. Ishida. 2000. New anabaenopeptins, potent carboxypeptidase-A inhibitors from the cyanobacterium *Aphanizomenon flos-aquae*. *J. Nat. Prod.* **63**:1280–1282.
 31. Namikoshi, M., and K. L. Rinehart. 1996. Bioactive compounds produced by cyanobacteria. *J. Ind. Microbiol.* **17**:373–384.
 32. Namikoshi, M., K. L. Rinehart, R. Sakai, R. R. Stotts, A. M. Dahlem, V. R. Beasley, W. W. Carmichael, and W. R. Evans. 1992. Identification of 12 hepatotoxins from a Homer Lake bloom of the cyanobacteria *Microcystis aeruginosa*, *Microcystis viridis*, and *Microcystis wesenbergii*: nine new microcystins. *J. Org. Chem.* **57**:866–872.
 33. Namikoshi, M., B. W. Choi, F. Sun, K. L. Rinehart, W. W. Carmichael, W. R. Evans, and V. R. Beasley. 1995. Seven more microcystins from Homer Lake cells: application of the general method structure assignment of peptides containing α,β -dehydroamino acid unit(s). *J. Org. Chem.* **60**:3671–3679.
 34. Neilan, B. A., D. Jacobs, T. del Dot, L. L. Blackall, P. R. Hawkins, P. T. Cox, and A. E. Goodman. 1997. rRNA sequences and evolutionary relationship among toxic and nontoxic cyanobacteria of the genus *Microcystis*. *Int. J. Syst. Bacteriol.* **47**:693–697.
 35. Neilan, B. A., E. Dittmann, L. Rouhiainen, R. A. Bass, V. Schaub, K. Sivonen, and T. Börner. 1999. Nonribosomal peptide synthesis and toxicogenicity of cyanobacteria. *J. Bacteriol.* **181**:4089–4097.
 36. Neumann, U., V. Campos, S. Cantarero, H. Urrutia, R. Heinze, J. Weckesser, and M. Erhard. 2000. Co-occurrence of non-toxic (cyanopeptolin) and toxic (microcystin) peptides in a bloom of *Microcystis* sp. from a Chilean lake. *Syst. Appl. Microbiol.* **23**:191–197.
 37. Orr, P. T., and G. J. Jones. 1998. Relationship between microcystin production and cell division rates in nitrogen-limited *Microcystis aeruginosa* cultures. *Limnol. Oceanogr.* **43**:1604–1614.
 38. Otsuka, S., S. Suda, R. Li, M. Watanabe, H. Oyaizu, S. Matsumoto, and M. M. Watanabe. 1999. Phylogenetic relationships between toxic and nontoxic strains of the genus *Microcystis* based on 16S to 23S internal transcribed spacer sequences. *FEMS Microbiol. Lett.* **172**:15–21.
 39. Otsuka, S., S. Suda, R. Li, S. Matsumoto, and M. M. Watanabe. 2000. Morphological variability of colonies of *Microcystis* morphospecies in culture. *J. Gen. Appl. Microbiol.* **46**:39–50.
 40. Reynolds, C. S., G. H. M. Jaworski, H. A. Cmiech, and G. F. Leedale. 1981. On the annual cycle of the blue-green alga *Microcystis aeruginosa* Kütz. *Emend. Elenkin. Phil. Trans. R. Soc. London B* **293**:419–477.
 41. Rohrlack, T., M. Henning, and J.-G. Kohl. 2001. Isolation and characterization of colonyforming *Microcystis aeruginosa* strains, p. 152–158. In I. Chorus (ed.), *Cyanotoxins—occurrence, causes, consequences*. Springer Verlag, Heidelberg, Germany.
 42. Rouhiainen, L., L. Paulin, S. Suomalainen, H. Hyytiäinen, W. Buikema, R. Hasselkorn, and K. Sivonen. 2000. Genes encoding syntheses of cyclic depsipeptides, anabaenopeptilides, in *Anabaena* strain 90. *Mol. Microbiol.* **37**: 156–167.
 43. Sano, T., and K. Kaya. 1995. Oscillamide Y, a chymotrypsin inhibitor from toxic *Oscillatoria agardhii*. *Tetrahedron Lett.* **36**:5933–5936.
 44. Shin, H. J., H. Matsuda, M. Murakami, and K. Yamaguchi. 1997. Anabaenopeptins E and F, two new cyclic peptides from the cyanobacterium *Oscillatoria agardhii* (NIES-204). *J. Nat. Prod.* **60**:139–141.
 45. Sivonen, K., and G. Jones. 1999. Cyanobacterial toxins, p. 41–111. In I. Chorus and J. Bartram (ed.), *Toxic cyanobacteria in water*. E & FN Spon, London, England.
 46. Tillet, D., E. Dittmann, M. Erhard, H. von Döhren, T. Börner, and B. A. Neilan. 2000. Structural organisation of the microcystin biosynthesis in *Microcystis aeruginosa* PCC7806: an integrated peptide-polyketide synthetase system. *Chem. Biol.* **7**:753–764.
 47. Vasconcelos, V. M., K. Sivonen, W. R. Evans, W. W. Carmichael, and M. Namikoshi. 1995. Isolation and characterization of microcystins (heptapeptide hepatotoxins) from Portuguese strains of *Microcystis aeruginosa* Kütz. *emend. Elenkin. Arch. Hydrobiol.* **134**:295–305.
 48. Vezic, C., L. Brient, K. Sivonen, G. Bertru, J.-C. Lefevre, and M. SalkinojaSalonen. 1998. Variation of microcystin content of cyanobacterial blooms and isolated strains in Lake Grand-Lieu (France). *Microb. Ecol.* **35**:126–135.
 49. Watanabe, M. F., S. Oishi, K.-I. Harada, K. Matsuura, H. Kawai, and M. Suzuki. 1988. Toxins contained in *Microcystis* species of cyanobacteria (blue-green algae). *Toxicon* **26**:1017–1025.
 50. Watanabe, M. F., M. Watanabe, T. Kato, K.-I. Harada, and M. Suzuki. 1991. Composition of cyclic peptide toxins among strains of *Microcystis aeruginosa* (blue-green algae, cyanobacteria). *Bot. Mag. Tokyo* **104**:49–57.
 51. Watanabe, M. 1996. Isolation, cultivation and classification of bloom-forming *Microcystis* in Japan, p. 13–35. In M. F. Watanabe, K.-I. Harada, W. W. Carmichael, and H. Fujiki (ed.), *Toxic Microcystis*. CRC Press, Boca Raton, Fla.
 52. Welker, M., S. Hoeg, and C. Steinberg. 1999. Hepatotoxic cyanobacteria in the shallow lake Müggelsee. *Hydrobiologia* **408/409**:263–268.