

## Natural Transformation of *Pseudomonas fluorescens* and *Agrobacterium tumefaciens* in Soil

SANDRINE DEMANÈCHE, ELISABETH KAY, FRANÇOIS GOURBIÈRE,  
AND PASCAL SIMONET\*

Laboratoire d'Écologie Microbienne, UMR 5557, Université Lyon I,  
69622 Villeurbanne Cedex, France

Received 12 October 2000/Accepted 7 March 2001

**Little information is available concerning the occurrence of natural transformation of bacteria in soil, the frequency of such events, and the actual role of this process on bacterial evolution. This is because few bacteria are known to possess the genes required to develop competence and because the tested bacteria are unable to reach this physiological state in situ. In this study we found that two soil bacteria, *Agrobacterium tumefaciens* and *Pseudomonas fluorescens*, can undergo transformation in soil microcosms without any specific physical or chemical treatment. Moreover, *P. fluorescens* produced transformants in both sterile and nonsterile soil microcosms but failed to do so in the various in vitro conditions we tested. *A. tumefaciens* could be transformed in vitro and in sterile soil samples. These results indicate that the number of transformable bacteria could be higher than previously thought and that these bacteria could find the conditions necessary for uptake of extracellular DNA in soil.**

According to gene sequence analyses and experimental data, gene transfer by natural transformation among microorganisms is involved in bacterial evolution and is likely to occur at present, providing bacteria the flexibility to rapidly adapt to changing environmental conditions (22, 26). However, such transfers remain very difficult to detect under natural conditions, particularly in soil, indicating they occur at very low frequencies (25, 27, 29). According to PCR data, large amounts of extracellular DNA are readily found in most soils and persist for months, indicating that the turnover of naturally released extracellular DNA would be quite slow (10, 19, 23, 24, 31). Extracellular DNA is thought to persist, because soil particles, particularly clay minerals, absorb macromolecules, a process which would not totally inhibit the transforming activity of DNA (7).

The main limitations to natural transformation in soil would be related to the recipient bacteria. Relatively few bacteria have been shown to carry the genes required to develop a natural state of competence; those described belong to phylogenetically distant taxa, including gram-positive and proteobacteria species (16). However, their number could be significantly higher considering that isolates are not systematically tested for this property and that more than 99% of all soil bacteria remain uncultured in vitro. Another limitation to transformation is related to the development under natural conditions of the competence state permitting active uptake of DNA. In soil, bacteria tend to live in a state of dormancy due to prevailing oligotrophic conditions, which would not be particularly favorable for the development of competence (16, 17). For instance, *Acinetobacter* sp. strain BD413, which exhibits high frequencies of transformation in vitro, is unable to develop a competence state when grown in soil. Moreover, this

physiological state is lost very rapidly in situ when in vitro-prepared competent cells are inoculated into soil (18).

In this study, we found that *Agrobacterium tumefaciens* and *Pseudomonas fluorescens*, two bacteria present in most soils and used as models for numerous molecular, physiological, and ecological studies, can be transformed without any physical or chemical treatment. Interestingly, the various conditions we tested did not permit *P. fluorescens* to reach the competence state in vitro, while transformants were detected in situ.

### MATERIALS AND METHODS

#### Bacterial strains, culture media, growth conditions, and cell enumeration.

Bacterial strains and plasmids used in this study are listed in Table 1. *Escherichia coli*, *A. tumefaciens*, and *P. fluorescens* strains were grown overnight at 37°C for *E. coli* and 28°C for the two other bacteria in liquid Luria-Bertani (LB) medium (containing, per liter, 10 g of Bacto Tryptone, 5 g of yeast extract, and 5 g of NaCl) supplemented with the appropriate antibiotics according to the concentrations given in Table 1. Bacterium counts were determined as the number of CFU on agar-solidified LB medium incubated overnight or for 48 h when targeting *E. coli* or the other soil bacteria, respectively.

**Inoculum preparation.** Bacterium inocula were prepared by diluting an overnight culture 20-fold into fresh medium and incubating it for an additional 2-h period (to an optical density of 0.6). These cultures were then centrifuged for 10 min at 6,500 × g before resuspension in the appropriate volume of sterile water to provide a 100-fold bacterial concentration. Using pure sterile water rather than a minimal salt medium avoided modification of competence development and natural transformation.

**Soil microcosms.** Microcosms consisted of 50-ml Falcon tubes in which were placed 30 g of air-dried and sieved (2-mm pore size) samples of a sandy loam soil (sand, 50%; silt, 41%; clay, 9%; organic matter, 40.6 g kg of dry soil<sup>-1</sup>; pH 5.6) collected at La Côte Saint-André (Isère, France). Sterile soil conditions were obtained by gamma-irradiating the microcosms at a dose of 25 kGy from a <sup>60</sup>Co source (Conservatome, Dagneux, France). Moisture was adjusted to 3 g of water for each microcosm according to the following protocol: 2.5 g of pure sterile water was added initially to the 30 g of dried soil, which was left to drain naturally for 2 days at room temperature. During the next 3 days, soils were mixed manually from time to time to homogenize moisture. Soils were then seeded with 15 µg of DNA in 0.25 ml of water and 0.25 ml of the concentrated recipient strain in water to provide a soil microcosm with 0.5 µg of DNA and 5 × 10<sup>9</sup> bacteria of dry soil<sup>-1</sup>.

In some experiments, pure plasmid DNA solutions were replaced by donor bacterial suspensions. Donor and recipient bacteria were inoculated at the same

\* Corresponding author. Mailing address: Laboratoire d'Écologie Microbienne, UMR 5557, Université Lyon I, 43 bd du 11 Novembre 1918, 69622 Villeurbanne cedex, France. Phone: 33 4 72 44 82 89. Fax: 33 4 72 43 12 23. E-mail: simonet@biomserv.univ-lyon1.fr.

TABLE 1. Bacterial strains and plasmids used in this study

Strain or plasmid	Resistance phenotype <sup>a</sup>	Use (reference)
<i>Escherichia coli</i> DH10B	Wild type	Donor cell
<i>Pseudomonas fluorescens</i> AK15	Wild type	Donor cell (32)
<i>Escherichia coli</i> 1504	Tet <sup>r</sup> (15) Spc <sup>r</sup> (60)	Recipient cell
<i>Pseudomonas fluorescens</i> LP59JG	Rif <sup>r</sup> (50) Gm <sup>r</sup> (10)	Recipient cell (11)
<i>Agrobacterium tumefaciens</i> GM19023	Rif <sup>r</sup> (100) Sm <sup>r</sup> (500)	Recipient cell
pKT230 (11.9 kb)	Kan <sup>r</sup> (50) Sm <sup>r</sup> (100)	Transforming DNA (1)
pPZP111 (8.9 kb)	Kan <sup>r</sup> (25) Cm <sup>r</sup> (25)	Transforming DNA (13)

<sup>a</sup> Numbers in parentheses are antibiotic concentrations (in micrograms per milliliter).

population level, with the final concentration remaining at  $5 \times 10^9$  bacteria g of dry soil<sup>-1</sup>. Control experiments were systematically conducted with pure water to replace each of the two biological inocula.

**Enumeration.** Following a 3-day incubation at 28°C to stabilize the population level (11), soil microcosms were treated with DNase I to degrade any persistent extracellular DNA (incubation for 3 h at 28°C with 1,000 U of DNase I in 90 ml of LB medium). Appropriate dilutions of the soil suspensions were spread on selective (Table 1) and LB media to enumerate transformants and recipient cells, respectively. Colonies were counted following a 2-day incubation at the appropriate temperature. Repetitions were performed as follows. For every experimental combination, six independent microcosms were treated separately, and each one was plated on five petri dishes. Thus, a total of 30 replicates were counted to detect transformants. Moreover, this entire procedure was repeated at least once subsequently.

Recipient cells were selected through their specific resistance to antibiotics (Table 1). The presence of plasmids in the growing colonies was confirmed by using a plasmid extraction kit (Qiagen Inc., Chatsworth, California), according to manufacturer instructions.

**Identification of strains.** The PCR amplification method included the use of 1 µl of bacterial cultures as template and a hot start procedure (3 min at 95°C) to release DNA, to avoid initial mispriming and to enhance primer specificity. Primers used, pA and pH (9), were complementary to part of the 16S ribosomal gene amplifying 1.6-kb-long PCR products. Restriction enzymes were from Boehringer Mannheim (Meylan, France) and used according to manufacturer instructions before fragments were separated on agarose gels.

**Recovering of transformants from soil.** We put 1, 10, 100, or 1,000 CFU of the plasmid-containing *Agrobacterium* or *Pseudomonas* into the sterile soil seeded with the stabilized plasmidless bacteria in the same way as described previously, followed by a 3-h incubation at 28°C in 90 ml of LB medium. Appropriate dilutions of the soil suspensions were spread on selective (Table 1) and LB media to enumerate plasmid-containing strains and plasmidless cells, respectively. Colonies were counted following a 2-day incubation at 28°C. Three independent microcosms were treated separately, and each was repeated subsequently. Each one was plated on five petri dishes.

**In vitro transformation protocols.** Plasmids were introduced into *E. coli* and *P. fluorescens* strains according to standard electroporation techniques (8). The development of a natural competence state in *P. fluorescens* was tested by mixing 20 µl of 100-fold-concentrated recipient cells from overnight cultures with 0.5 µg of DNA (20 µl of a 25-µg ml<sup>-1</sup> concentrated plasmid solution) or 20 µl of 100-fold-concentrated donor cells and deposited onto a GTTP filter (Millipore, Bedford, Massachusetts) on each of the solid media described below. After drying, the Petri dishes were incubated at 28°C for 24 h before bacteria were resuspended in 2 ml of pure sterile water and plated out on the same media under the appropriate selective conditions. Liquid conditions were also tested by mixing 40 µl of recipient cells from overnight cultures with 950 µl of each of the liquid culture media and 10 µl of a 100-µg ml<sup>-1</sup> concentrated plasmid solution. The tubes were incubated for 24 h at 28°C with shaking before the bacteria were plated out on media with appropriate antibiotics. Additional tests included a second 10-µl inoculation of a 100-µg ml<sup>-1</sup> concentrated plasmid solution 12 h after the first one and incubation was continued for another 12-h period. The various media we tested included (i) the standard P medium (60 mM lactic acid, 11 mM KH<sub>2</sub>PO<sub>4</sub>, 95 mM Na<sub>2</sub>HPO<sub>4</sub>, 0.81 mM MgSO<sub>4</sub>, 37 mM NH<sub>4</sub>Cl, 68 µM CaCl<sub>2</sub>, 1.8 µM FeSO<sub>4</sub>, and 1 ml of trace element liter<sup>-1</sup>) (21) and derived media, each with a specific deficiency obtained by decreasing the concentration of the corresponding nutrient as follows. To obtain a carbon-limited culture, only 20

mM lactic acid was added. For the nitrogen limitation, only 3 mM NH<sub>4</sub>Cl was added together with 60 mM lactic acid. The potassium limitation was obtained by adding only 50 µM KH<sub>2</sub>PO<sub>4</sub> and 30 mM Na<sub>2</sub>HPO<sub>4</sub> together with 60 mM lactic acid (20); (ii) the MM medium [containing, per liter, KH<sub>2</sub>PO<sub>4</sub>, 3.4 g; (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.5 g; MgSO<sub>4</sub> · 7H<sub>2</sub>O, 0.05 g; FeSO<sub>4</sub> · 7H<sub>2</sub>O, 0.125 mg; thiamine, 0.25 mg; glucose, 2 g; glycerol, 4 ml]; (iii) two soil-based media: a rough soil medium consisting of 50% (wt vol<sup>-1</sup>) ground soil in sterile distilled water used as a liquid medium or solidified by adding 15 g of agar liter<sup>-1</sup> and a mineral soil medium in which the previous soil suspension in water was centrifuged twice for 15 min at 4,500 × g before the supernatant was filtered (pore size, 0.2 µm; Millipore). The filtered solution was used directly as a liquid medium or supplemented with 15 g of agar liter<sup>-1</sup> to be used as a solid medium. Five repetitions were carried out for each sample, repeated at least twice subsequently. Controls consisted of filters with donor or recipient strains alone.

**Statistical analysis of data.** Assuming that each petri dish received roughly the same number of bacteria, each petri dish was considered to be a sampling unit, and the presence or absence of any transformant was considered to be the variable. In other words, the result was considered negative only if no transformants appeared. We calculated the probability that the in vitro results were due to a sampling effect and compared this to results of soil experiments. We assumed as a null hypothesis that the frequency of transformation and the probability of observing negative plates were the same in soil and in vitro experiments. We estimated the probability of obtaining negative plates in soil experiments and calculated the probability of observing no positive plates in the in vitro experiments under this hypothesis. We used the zero term of binomial distribution as in the most-probable-number method of Cochran (6), but using only one dilution:

$$P = (n_{s1}/N_s)^{N_v} \quad (1)$$

where  $n_{s1}$  is the number of negative plates in the soil experiment,  $N_s$  is the total number of plates in the soil experiment, and  $N_v$  is the total number of plates in the in vitro experiment.

For example, if we have 20 positive plates among 100 soil replicates and no positive plate among 15 in vitro replicates, the probability that this last observation results from a sampling effect is  $(80/100)^{15}$ , or 0.035. Given that a 95% confidence interval is acceptable ( $P < 0.05$ ), we can thus assume that the two sets of data were not significantly the same.

## RESULTS AND DISCUSSION

### Transformation of *A. tumefaciens* and *P. fluorescens* in soil.

Transformation experiments were conducted directly in soil microcosms by inoculating recipient bacteria and DNA. Detection of clones exhibiting the appropriate antibiotic resistance at frequencies significantly higher than the detection limit (number of recipient cells<sup>-1</sup>) and possessing the plasmid would indicate the ability of the tested bacteria to be naturally transformed. Because the plasmids we used were lacking the *tra* function and could not transfer autonomously, they could not be mobilized and could not form cointegrates with additional conjugative plasmids; the only way to transfer the marker genes to the recipient strain was via transformation. We did not detect mutants in the various DNA-less and bacterium-less controls. This would indicate that the spontaneous mutation rate remained below the detection limit. We tested the quantity of transformants (1 to 1,000 CFU) that we could recover in sterile soils containing  $5 \times 10^9$  plasmidless bacteria under the same conditions as those in transformation experiments. The method we used allowed the detection of  $18 \pm 15$  CFU g of dry soil<sup>-1</sup> when 10 CFU was seeded in the 30-g soil microcosms thanks to the enrichment step.

In sterile soil, the total population of *A. tumefaciens* stabilized at  $5.5 \times 10^8 \pm 2.4 \times 10^7$  cells g of dry soil<sup>-1</sup>. When the soils were inoculated with a plasmid solution at 0.5 µg of DNA g of dry soil<sup>-1</sup>, transformants exhibiting resistance to specific antibiotics and carrying plasmid pPZP111 (Fig. 1) were detected at frequencies reaching  $1.3 \times 10^{-8}$  (Table 2), corre-




FIG. 1. Confirmation of the presence of plasmid pPZP111 in *A. tumefaciens* GM19023 transformants. Lanes: A, smart ladder (Eurogentec); ND, nondigested profile of extracted plasmids; D, *ScaI*-digested profile of extracted plasmids; 1, clone from filter experiments with *P. fluorescens* AK15(pPZP111) as donor; 2, clone from filter experiments with pPZP111 as donor; 3 and 4, clones from experiments with *P. fluorescens* AK15(pPZP111) as donor in nonsterile soil; 5 and 6, clones from experiments with pPZP111 as donor in sterile soil; 7, *A. tumefaciens* GM19023.

sponding to  $12 \pm 15$  transformants g of dry soil<sup>-1</sup>. We also carried out experiments in which the recipient bacteria were coinoculated with donor cells of *P. fluorescens* AK15 containing the plasmid pPZP111. In such cases, transformants were also detected (Fig. 1) at frequencies reaching  $2.7 \times 10^{-9}$  (Table 2), corresponding to  $6.7 \pm 13.3$  transformants g of dry soil<sup>-1</sup>.

Similar experiments were conducted with *P. fluorescens* LP59JG, which stabilized at  $2.4 \times 10^8 \pm 9.8 \times 10^7$  cells g of dry soil<sup>-1</sup> after 4 days. When the soils were inoculated with plasmid pKT230, transformants exhibiting resistance to kanamycin and streptomycin and carrying the plasmid were detected (Fig. 2) at frequencies reaching  $8.3 \times 10^{-8}$ , corresponding to  $20 \pm 16.3$  transformants g of dry soil<sup>-1</sup> (Table 2). We also carried out experiments in which the recipient bacteria were coinoculated with donor cells (*E. coli* containing the plasmid pKT230). In such cases, transformants were also detected (Fig. 2) at frequencies reaching  $5.8 \times 10^{-8}$  (Table 2), corresponding to  $33.3 \pm 36.5$  transformants g of dry soil<sup>-1</sup>. *E. coli* and *P. fluorescens* strains were identified by comparison of restriction patterns from PCR-restriction fragment length polymorphism analysis conducted on 16S DNA with restriction enzyme *AluI* (data not shown).

In nonsterile soil samples, the recipient *P. fluorescens* LP59JG sustained levels up to  $2.1 \times 10^7 \pm 7.3 \times 10^6$  cells g of dry soil<sup>-1</sup> after 4 days in soil, while the *E. coli* donor strain dropped to  $3.2 \times 10^5 \pm 1.2 \times 10^5$  cells g of dry soil<sup>-1</sup>. Among colonies growing on the selective medium ( $867 \pm 76$  cells g of dry soil<sup>-1</sup>), some were identified as *P. fluorescens*. This was assessed by the similarity of the 16S ribosomal restriction patterns of PCR products obtained from these clones compared to those of the recipient strain (data not shown). Moreover, these clones harbor an 11.9 kb plasmid, which exhibited the restriction patterns corresponding to the donor plasmid pKT230 (Fig. 2). The presence of indigenous soil bacteria resistant to the four selective antibiotics prevented us from determining the precise transformation rate. However, the transformation rate in nonsterile soil seems to be higher than that in sterile soils. We cannot justify this observation, but we suggest that it may be due to the presence of an

organic compound in soil that is in part destroyed by soil sterilization.

**Detection of transformants in vitro.** The two bacteria which exhibited evidence of transformation in soil, *A. tumefaciens* GM19023 and *P. fluorescens* LP59JG, as well as *E. coli* 1504 as the negative control, were tested in vitro for their ability to take up, internalize, and express extracellular DNA.

When transformed with the plasmid pPZP111 harboring the resistance genes to kanamycin and chloramphenicol (Table 1), *A. tumefaciens* provided recombinant clones at frequencies reaching  $6 \times 10^{-9}$  ( $14 \pm 10$  transformants ml<sup>-1</sup>), while the detection limit remained at  $1.6 \times 10^{-10}$  (Table 3). Moreover, detection of plasmids of the expected size (11.9 kb) (Fig. 1) helped confirm that a transformation event occurred, allowing the recipient strain to take up and replicate the donor plasmid DNA.

On the other hand, *E. coli* 1504 and *P. fluorescens* LP59JG failed to provide recombinant clones after they were incubated with the donor DNA solution or even with donor cells. As expected, the coculture of the recipient *E. coli* 1504 with *E. coli* DH10B(pKT230) or with the pure plasmid solution failed to provide detectable transformants, because *E. coli* is a nonnaturally competent bacterium for which transformation requires chemical or physical treatments.

More surprisingly, when tested in vitro, *P. fluorescens* LP59JG also failed to provide detectable transformants. However, two explanations are possible. This could be related to a

TABLE 2. Transformation rate and detection limit of experiments conducted in sterile soil microcosms

Donor strain or plasmid	Recipient strain	Transformation rate <sup>a</sup>	Detection limit
<i>E. coli</i> DH10B (pKT230)	<i>P. fluorescens</i> LP59JG	$5.8 \times 10^{-8}$	$9.8 \times 10^{-9}$
pKT230	<i>P. fluorescens</i> LP59JG	$8.3 \times 10^{-8}$	$2.0 \times 10^{-8}$
<i>P. fluorescens</i> AK15 (pPZP111)	<i>A. tumefaciens</i> GM19023	$2.7 \times 10^{-9}$	$2.2 \times 10^{-9}$
pPZP111	<i>A. tumefaciens</i> GM19023	$1.3 \times 10^{-8}$	$6.0 \times 10^{-9}$

<sup>a</sup> Efficiencies of transformation were expressed as the number of transformants relative to that of recipient cells.




FIG. 2. Confirmation of the presence of plasmid pKT230 in *P. fluorescens* LP59JG transformants. Lanes: A, smart ladder (Eurogentec); ND, nondigested profile of extracted plasmids; D, *Pst*I-digested profile of extracted plasmids; 1, *E. coli* DH10B(pKT230); 2, clone from experiments with *E. coli* DH10B(pKT230) as donor in sterile soil; 3, clone from experiments with *E. coli* DH10B(pKT230) as donor in nonsterile soil; 4 and 5, clones from experiments with pKT230 as donor in sterile soil; 6, *P. fluorescens* LP59JG.

sampling effect due to the number of replicates. But according to equation 1, we found that this hypothesis was refuted. We observed 76 negative plates and 30 positive plates in the various transformation events we tested with pKT230. The negative probability was thus  $76/106$ . As the *in vitro* experiment provided 15 negative plates and 0 positive plates with pKT230,  $P$  is equal to  $(76/106)^{15}$ , or 0.0068. In the same way, transformation experiments conducted with the donor *E. coli* DH10B (pKT230) provided a  $P$  of  $(41/60)^{35}$ , or 0.0000016. As  $P < 0.05$ , the test is significant, thus proving that the absence of transformant was not due to a sampling effect.

The second hypothesis concerns the way we monitored potential transformations. Conditions tested included liquid and solid LB media and plasmid solutions or living *E. coli* or *P. fluorescens* AK15 cells as sources of donor DNA. According to the detection limit, the potential transformation efficiency was thus below  $2.6 \times 10^{-9}$  (Table 3). However, development of competence in naturally transformable bacteria is known to depend upon the chemical composition of the medium, with requirements varying greatly. For some bacteria, specific nutrients are required to induce this physiological state, while in other cases the presence of particular substances inhibits the development of competence. For instance, media with limited carbon and energy sources or deficient in nitrogen and phosphorus have been found to favor the development of competence in *Pseudomonas stutzeri* (15). On the other hand, bacteria such as *Acinetobacter* sp. strain BD413 (14, 21) and *Ralstonia solanacearum* (5) were transformable in media with totally different requirements. In order to cover a wide range of conditions, we tested various media described as promoting competence development in most of the naturally transformable bacteria. This included the use of the P and the MM media favoring competence development in *Acinetobacter* sp. strain BD413 and *R. solanacearum*, respectively, but also media characterized by a specific deficiency to test whether inhibition of competence development would be related to the presence of a specific substance. Each medium was tested according to two protocols based on incubation of bacteria on agar-solidified media and in liquid. Finally, we also tested two soil-based media, including a rough medium containing the whole soil sample in water and a medium containing only the soil mineral fraction. None of the conditions tested enabled us to detect

transformants, which would have indicated that *P. fluorescens* can develop competence *in vitro*. In spite of our efforts and the use of rich, soil-based or deficient media, we were unable to determine whether an essential component for transformation present in the soil would be missing *in vitro* or whether the media we used contained an inhibiting product. Another hypothesis could involve a rapid degradation of the transforming DNA by excretion of an active nuclease before the cells became competent, as evidenced in *Bacillus subtilis* (2). However, in spite of a modification of the protocol to include two DNA inoculations separated by a 12-h lag time (see Materials and Methods), transformants remained undetectable. These data confirm that bacterial physiology remains largely unknown with numerous specific limitations. In the case of *P. fluorescens*, these limitations prevent the development of a competence state *in vitro*, while for more than 99% of the indigenous microflora it is the growth of colonies on petri dishes which is inhibited.

**Transformation mechanism.** A major question regarding these results is related to the mechanism by which bacteria can take up DNA. A first hypothesis deals with the presence of the molecular machinery permitting bacteria to develop a competence state. Tests should be conducted on bacteria belonging to the 99% nonculturable fraction of the soil microflora (28). Another hypothesis to explain the transformants we obtained

TABLE 3. Transformation rate and detection limit of experiments conducted *in vitro*

Donor strain or plasmid	Recipient strain	Transformation rate <sup>a</sup>	Detection limit
<i>E. coli</i> DH10B (pKT230)	<i>E. coli</i> 1504	$<4.2 \times 10^{-8}$	$4.2 \times 10^{-8}$
pKT230	<i>E. coli</i> 1504	$<5.0 \times 10^{-8}$	$5.0 \times 10^{-8}$
<i>E. coli</i> DH10B (pKT230)	<i>P. fluorescens</i> LP59JG	$<1.3 \times 10^{-7}$	$1.3 \times 10^{-7}$
pKT230	<i>P. fluorescens</i> LP59JG	$<2.6 \times 10^{-9}$	$2.6 \times 10^{-9}$
<i>P. fluorescens</i> AK15 (pPZP111)	<i>A. tumefaciens</i> GM19023	ND <sup>b</sup>	$4.4 \times 10^{-9}$
pPZP111	<i>A. tumefaciens</i> GM19023	$6.0 \times 10^{-9}$	$1.6 \times 10^{-10}$

<sup>a</sup> Efficiencies of transformation were expressed as the number of transformants relative to that of recipient cells.

<sup>b</sup> ND, *A. tumefaciens* GM19023 was not detectable when cultivated concomitantly with *P. fluorescens* AK15 *in vitro*.

with *A. tumefaciens* and *P. fluorescens* deals with a mechanism that would not be related to the well-defined natural transformation. *E. coli* transformation has been observed in various environments such as freshwater (4) and even foodstuffs (3). The naturally existing chemical ( $\text{Ca}^{2+}$  concentration) or physical (heat shock) parameters could provide *E. coli* with optimal conditions for a natural process such as transformation. Moreover, it has been hypothesized that induction of competence in *E. coli* would involve a physiological rather than a physicochemical mechanism (4). From our results, it can be supposed that soils, unlike other media, do not provide the required conditions for physiological or physicochemical transformation of *E. coli*. This could be due to the fact that *E. coli* is not a usual inhabitant of soils. When inoculated into nonsterile soil samples, the number of cells decreases rapidly, and so the competitive potential of *E. coli* becomes much lower than that of indigenous soil bacteria (23). However, the actual status leading to a physical, chemical, physiological, or genetic induction of competence in *A. tumefaciens* and *P. fluorescens*, which are natural inhabitants of soils, remains unknown. Additional experiments are necessary to determine whether the currently living strains of these bacteria are fitted with the genes involved in the development of competence, as observed in typical naturally competent bacteria, or whether they can take up DNA by other mechanisms. We can thus assume that the list of bacteria naturally capable of transformation is certainly longer than previously expected. Moreover, the fact that our findings revealed this property in well-known bacteria such as *A. tumefaciens* and *P. fluorescens* points out the interest of a more exhaustive study among available isolates, keeping in mind that development of competence was found to be species and even strain specific in bacteria (16).

Whatever the induction mechanisms operating in current bacterial strains, our results, based on experiments simulating natural conditions more closely than previous studies, demonstrate that transformation-mediated gene transfers can occur in soils. Whether such transformation-mediated gene exchange occurs at sufficiently high frequencies to contribute significantly to bacterial genome evolution remains to be investigated. However, factors such as the number of bacteria in the biosphere ( $5 \times 10^{30}$  bacterial cells) (30), but also the time scale (3 billion years) will have to be considered carefully when tracking bacterial genome evolution and the related mechanisms.

#### ACKNOWLEDGMENTS

We are grateful to Stéphane Peyrard for technical assistance.

This work was done as part of the Biotechnologies program supported by the Ministère Français de l'Enseignement Supérieur et de la Recherche (MENRT). S.D. and E.K. were funded by a grant from the MENRT.

#### REFERENCES

1. Bagdasarian, M., R. Lurz, B. Rückert, F. C. H. Franklin, M. M. Bagdasarian, J. Frey, and K. N. Timmis. 1981. Specific-purpose plasmid cloning vectors. II. Broad host range, high copy number, RSF1010-derived vectors, and a host-vector system for gene cloning in *Pseudomonas*. *Gene* **16**:237–247.
2. Basse, G., M. G. Lorenz, and W. Wackernagel. 1994. A biological assay for the sensitive and quantifiable detection of extracellular microbial DNases. *J. Microbiol. Methods* **20**:137–147.
3. Bauer, F., C. Hertel, and W. P. Hammes. 1999. Transformation of *Escherichia coli* in foodstuffs. *Syst. Appl. Microbiol.* **22**:161–168.
4. Baur, B., K. Hanselmann, W. Schlimme, and B. Jenni. 1996. Genetic transformation in freshwater: *Escherichia coli* is able to develop natural competence. *Appl. Environ. Microbiol.* **62**:3673–3678.
5. Bertolla, F., F. van Gijsegem, X. Nesme, and P. Simonet. 1997. Conditions for natural transformation of *Ralstonia solanacearum*. *Appl. Environ. Microbiol.* **63**:4965–4968.
6. Cochran, W. G. 1950. Estimation of bacterian densities by means of the most probable number. *Biometrics* **6**:105–116.
7. Demanéche, S., L. Jocteur-Monrozier, H. Quiquampoix, and P. Simonet. 2001. Evaluation of biological and physical protection against nuclease degradation of clay-bound plasmid DNA. *Appl. Environ. Microbiol.* **67**:293–299.
8. Drury, L. 1996. Transformation of bacteria by electroporation. In A. Harwood (ed.), *Methods in molecular biology*. Humana Press Inc., Totowa, N.J.
9. Edwards, U., T. Rogall, H. Blöcker, M. Emde, and E. C. Böttger. 1989. Isolation and direct complete nucleotide determination of entire genes. Characterization of a gene coding 16S ribosomal RNA. *Nucleic Acids Res.* **16**:7843–7853.
10. Gebhard, F., and K. Smalla. 1999. Monitoring field releases of genetically modified sugar beets for persistence of transgenic plant DNA and horizontal gene transfer. *FEMS Microbiol. Ecol.* **28**:261–272.
11. Ghiglione, J. F., F. Gourbière, P. Potier, L. Philippot, and R. Lensi. 2000. Role of respiratory nitrate reductase in ability of *Pseudomonas fluorescens* YT101 to colonize the rhizosphere of maize. *Appl. Environ. Microbiol.* **66**:4012–4016.
12. Ghiglione, J.-F., L. Philippot, P. Normand, R. Lensi, and P. Potier. 1999. Disruption of *narG*, the gene encoding the catalytic subunit of respiratory nitrate reductase, also affects nitrite respiration in *Pseudomonas fluorescens* YT101. *J. Bacteriol.* **181**:5099–5102.
13. Hajdukiewicz, P., Z. Svab, and P. Maliga. 1994. The small, versatile *pPZP* family of *Agrobacterium* binary vectors for plant transformation. *Plant Mol. Biol.* **25**:989–994.
14. Lorenz, M. G., K. Reipschläger, and W. Wackernagel. 1992. Plasmid transformation of naturally competent *Acinetobacter calcoaceticus* in non-sterile soil extract and groundwater. *Arch. Microbiol.* **157**:355–360.
15. Lorenz, M. G., and W. Wackernagel. 1991. High frequency of natural genetic transformation of *Pseudomonas stutzeri* in soil extract supplemented with carbon/energy and phosphorus source. *Appl. Environ. Microbiol.* **57**:1246–1251.
16. Lorenz, M. G., and W. Wackernagel. 1994. Bacterial gene transfer by natural genetic transformation in the environment. *Microbiol. Rev.* **58**:563–602.
17. Nielsen, K. M., A. M. Bones, and J. D. Van Elsas. 1997. Induced natural transformation of *Acinetobacter calcoaceticus* in soil microcosms. *Appl. Environ. Microbiol.* **63**:3972–3977.
18. Nielsen, K. M., M. D. M. van Weerelt, T. N. Berg, A. M. Bones, A. N. Hagler, and J. D. van Elsas. 1997. Natural transformation and availability of transforming DNA to *Acinetobacter calcoaceticus* in soils microcosms. *Appl. Environ. Microbiol.* **63**:1945–1952.
19. Paget, E., M. Lebrun, G. Freyssonnet, and P. Simonet. 1998. The fate of recombinant plant DNA in soil. *Eur. J. Soil Biol.* **34**:81–88.
20. Palmén, R., P. Buijsman, and K. J. Hellingwerf. 1994. Physiological regulation of competence induction for natural transformation in *Acinetobacter calcoaceticus*. *Arch. Microbiol.* **162**:344–351.
21. Palmén, R., B. Vosman, P. Buijsman, C. K. D. Breek, and K. J. Hellingwerf. 1993. Physiological characterization of natural transformation in *Acinetobacter calcoaceticus*. *J. Gen. Microbiol.* **139**:295–305.
22. Paul, J. H. 1999. Microbial gene transfer: an ecological perspective. *J. Mol. Microbiol. Biotechnol.* **1**:45–50.
23. Recorbet, G., C. Picard, P. Normand, and P. Simonet. 1993. Kinetics of the persistence of chromosomal DNA from genetically engineered *Escherichia coli* introduced into soil. *Appl. Environ. Microbiol.* **59**:4289–4294.
24. Romanowski, G., M. G. Lorenz, and W. Wackernagel. 1993. Use of polymerase chain reaction and electroporation of *Escherichia coli* to monitor the persistence of extracellular plasmid DNA introduced into natural soils. *Appl. Environ. Microbiol.* **59**:3438–3446.
25. Sikorski, J., S. Graupner, M. G. Lorenz, and W. Wackernagel. 1998. Natural genetic transformation of *Pseudomonas stutzeri* in a non-sterile soil. *Microbiology* **144**:569–576.
26. Smith, M. W., D.-F. Feng, and R. F. Doolittle. 1992. Evolution by acquisition: the case for horizontal gene transfers. *Trends Biochem. Sci.* **17**:489–493.
27. Stotzky, G., E. Gallori, and M. Khanna. 1996. Transformation in soil. In A. D. L. Akkermans, J. D. van Elsas, and F. J. de Bruijn (ed.), *Molecular microbial ecology manual*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
28. Strätz, M., M. Mau, and K. N. Timmis. 1996. System to study horizontal gene exchange among microorganisms without cultivation of recipients. *Mol. Microbiol.* **22**:207–215.
29. Trevors, J. T. 1996. DNA in soil: adsorption, genetic transformation, molecular evolution and genetic microchip. *Antonie Leeuwenhoek* **70**:1–10.
30. Whitman, W. B., D. C. Coleman, and W. J. Wiebe. 1998. Prokaryotes: the unseen majority. *Proc. Natl. Acad. Sci. USA* **95**:6578–6583.
31. Widmer, F., R. J. Seidler, and L. S. Watrud. 1996. Sensitive detection of transgenic plant marker gene persistence in soil microcosms. *Mol. Ecol.* **5**:603–613.
32. Ye, R. W., A. Arunakumari, B. A. Averill, and J. M. Tiedje. 1992. Mutants of *Pseudomonas fluorescens* deficient in dissimilatory nitrite reduction are also altered in nitric oxide reduction. *J. Bacteriol.* **174**:2560–2564.