

Molecular Biological Detection and Characterization of *Clostridium* Populations in Municipal Landfill Sites

M. I. Van Dyke* and A. J. McCarthy

School of Biological Sciences, University of Liverpool, Liverpool L69 7ZB, United Kingdom

Received 27 July 2001/Accepted 3 January 2002

Primer sets specific for 16S rRNA genes were designed for four phylogenetic groups of clostridia known to contain mesophilic cellulolytic species. Specific amplification of these groups from landfill leachate DNA extracts demonstrated the widespread occurrence of clostridia from the *Clostridium thermocellum* and *C. leptum* groups. In contrast, the *C. botulinum* group was never detected, and the *C. coccoides*-*C. lentocellum* group was only occasionally detected. Amplification products were analyzed by temporal thermal gel electrophoresis to generate profiles of the clostridial groups and to identify dominant bands. Sequence analysis of 17 landfill clones confirmed that the primers were specific for the clostridial subgroups and that the cloned sequences had a close relationship with known cellulose-degrading clostridia. The primers have therefore been authenticated for use in the rapid identification of clostridia in anaerobic environments.

Anaerobic degradation in landfills involves several coordinated groups of microorganisms and follows a process that is typical of waste degradation in anaerobic environments, such as soils, sediment, and sludge. As the primary stage of waste degradation, polysaccharide breakdown is an important limiting factor in anaerobic treatment of waste, which in municipal landfills primarily involves the decomposition of complexed polymers, including cellulose, hemicellulose, and lignin. Although cellulose is an important substrate in landfills, anaerobic degradation is poorly understood, and our knowledge is based on studies using culture-based methods (1). The enumeration of cellulolytic bacteria in landfills has often resulted in low cell counts (10), suggesting that culture-based methods may be underestimating bacterial numbers. The aim of the study reported here was to use information on 16S rRNA gene sequences to develop tools for the specific detection of cellulose-degrading bacteria in landfill sites.

It is likely that anaerobic cellulose degradation in landfills is due primarily to bacteria related to the genera *Clostridium* and *Eubacterium*. Although very few cellulolytic strains have been isolated from landfill sites, Westlake et al. (15) have identified isolates related to these two groups. The genus *Clostridium* and its relatives constitute an ancient group whose members exhibit a wide range of phenotypic characteristics. Phylogenetic analysis of 16S rRNA genes shows that the group is very diverse, with deeply branching clusters that include nonclostridial species. Comparison of 16S rRNA genes has allowed the division of the genus *Clostridium* into subgroups, and cellulose-degrading representatives from genera such as *Clostridium*, *Eubacterium*, and *Ruminococcus* can be found in a number of different clusters (3). However, mesophilic cellulose-degrading strains tend to be found in groups I, III, IV, and XIVab, with group III comprising only cellulose-degrading strains to date.

Consequently, we have exploited this clostridial 16S rRNA

database to investigate the presence of clostridial subgroups I, III, IV, and XIVab in landfill sites. Although these subgroups do not comprise solely cellulose-degrading species, their detection could be used to indicate the distribution of saccharolytic and proteolytic degrading bacteria in landfill sites. Specific PCR amplification, temporal thermal gel electrophoresis (TTGE), and sequence analysis are used to detect and profile these key groups of clostridia. TTGE can separate DNA fragments of the same length but with different sequence compositions. When combined with specific or nonspecific gene amplification, this method can rapidly profile the genetic diversity of microbial populations.

Pooled leachate samples from sites designated R, C, H, W, B, P, and S were obtained from landfill sites in the northwest of England that contained primarily municipal solid waste and were provided by UK Waste Ltd., Terry Adams Ltd., and Cleanaway Ltd. Samples So and Br were obtained from test cell reactors containing municipal solid waste and were provided by J. Wayne, Centre for Applied Microbiology Research, Porton Down, United Kingdom, and the Energy Technology Support Unit. One-liter samples of leachate were concentrated by centrifugation at $27,000 \times g$ for 40 min, and the solids were resuspended in 20 ml of 0.1 M K_2HPO_4 buffer. Aliquots derived from 75 ml of leachate were harvested by centrifugation at $16,000 \times g$ for 5 min, and the pellets were stored at $-70^\circ C$. A sample from sheep rumen, provided by D. Mercer, Rowett Research Institute, Aberdeen, United Kingdom, was used as a control known to contain a high concentration of cellulolytic bacteria. One-milliliter samples of rumen fluid were concentrated by centrifugation at $16,000 \times g$ for 5 min, and the pellets were stored at $-70^\circ C$. Reference strains and their sources are listed in Table 1.

DNA was extracted from pure cultures with a Hybaid RiboLyser. A 2-ml tube containing 0.5 g of glass beads (0.17- to 0.18-mm diameter), 0.5 ml of 0.12 M K_2HPO_4 (pH 8.0), 0.5 ml of saturated phenol (pH 8.0), and 0.5 ml of cell suspension was processed at 6 m/s for 30 s, placed on ice for 5 min, and centrifuged at $16,000 \times g$ for 5 min. Supernatant was extracted with phenol-chloroform-isoamyl alcohol (25:24:1) and chloro-

* Corresponding author. Present address: GAP EnviroMicrobial Services, 1020 Hargrieve Rd., Unit 14, London, Ontario, Canada N6E 1P5. Phone: (519) 681-0571. Fax: (519) 681-7150. E-mail: mvandyke@gapenviromic.com.




FIG. 1. TTGE profiles of 16S rRNA genes of *Clostridium* clusters III, IV, and XIVab. Amplification products using group-specific primers were reamplified using primer TTGE-1 (cluster III) or TTGE-2 (clusters IV and XIVab) and were separated by TTGE analysis. The profiles comprise amplification products from landfill sites R, C, H, W, B, P, Br, S, and So. Cloned sequences that matched bands marked with arrows on the TTGE gel were sequenced and used for phylogenetic analysis.

and ImageQuant software (Molecular Dynamics). DNA preparations from pure cultures were used to optimize the conditions for band separation. TTGE profiles from each landfill site were repeated at least twice, and the banding patterns were shown to be reproducible. Representative TTGE profiles for each *Clostridium* group are shown in Fig. 1.

As exemplified in Fig. 2, clones that gave bands of the same mobility as the most intense bands in landfill leachate (Fig. 1) were selected for sequencing. The entire cloned insert, obtained using group-specific primers, was sequenced. Plasmids containing cloned inserts were purified using the Qiagen plasmid miniprep kit. Sequence analysis was performed using the ABI 373 (Perkin-Elmer) and Li-Cor 4200 (MWG-Biotechnology) sequencing systems. Further 16S rRNA gene sequence data were obtained from the RDP. Sequence alignments were manipulated using the Wisconsin package version 8.1 and the Genetic Data Environment. Phylogenetic analysis was performed with TREECON for Windows version 1.3b (14) using the Jukes-Cantor (6) and neighbor-joining (13) distance calculations, with bootstrap analysis performed on 100 replicates.

No amplification products were obtained from any landfill leachate or rumen fluid samples with primers specific for the *Clostridium botulinum* group (cluster I), indicating that this group of bacteria is either not present or present in low numbers. Similarly, Franks et al. (4) could detect only low numbers of cells that hybridized to the cluster I probe in human fecal samples.

Both forward and reverse primers for the *C. thermocellum* group (cluster III) were designed in this study. Cluster III primers amplified DNA from landfill leachate from each site tested and from rumen fluid, resulting in gene products of the expected size. The *C. thermocellum* group is of particular interest in the context of this work, as to date it contains se-

quences only from cellulose-degrading strains isolated from environmental sources (8). TTGE analysis of *Clostridium* cluster III 16S rRNA genes showed that there were one or two dominant species present in each landfill site (Fig. 1). Phylo-




FIG. 2. TTGE profiles of 16S rRNA genes of *Clostridium* cluster III amplified from Br and So landfill leachate DNAs. The lanes labeled "clones" contain amplification products of DNAs cloned from sites Br and So.




FIG. 3. Phylogenetic tree showing relationships of 16S rRNA genes from *Clostridium* cluster III. Sequences recovered from landfill sites are in boldface. The tree was constructed using the Jukes-Cantor distance matrix and the neighbor-joining method. The scale bar represents a 2% difference in nucleotide sequence positions.

genetic analysis showed that the dominant sequences amplified from landfill sites grouped within sequences from pure-culture representatives (Fig. 3). Only one clone from site S (g3S5) was closely related to the thermophile *C. thermocellum*. Cloned sequences from sites H (g3H3) and Br (g3Br5) were similar to *C. aldricchii* and its relatives, and the remaining five clones from sites P, C, B, W, and R (g3P17, g3C1, g3B5, g3W3, and g3R1) were most closely related to the group containing *C. papyrosolvans*.

The *C. leptum* group (cluster IV) is closely related to cluster III (3) and contains representatives of a mixture of genera, including *Clostridium*, *Eubacterium*, and *Ruminococcus*. A number of species are mesophilic and cellulolytic, including *Ruminococcus albus*, *Ruminococcus flavefaciens*, and *C. cellulosi*. Noncellulolytic strains are also present, although many of these will degrade other polysaccharides (12). Using primers specific for the *C. leptum* group, amplification products were obtained from the rumen fluid sample and from all landfill sites except R and S. TTGE profiles showed that the landfill sites contained one or two strong bands with a number of less intensely stained bands (Fig. 1). Cloned sequences from landfill sites grouped among sequences from pure-culture representatives of this phylum (Fig. 4), confirming the specificity of the primer set. Clones from sites C, H, P, and Br (g4C2, g4H4, g4P3, and g4Br3) were all closely related to *Eubacterium plautii* and *Sporobacter termitidis*, clone g4H6 was closely related to *Ruminococcus bromii*, clone g4W7 was closely related to *Fusobacterium prausnitzii*, and g4B3 was not closely related to any

of the *Clostridium* cluster IV pure-culture sequences. Clones from site So were not analyzed.

Clostridium cluster XIVab is a large and diverse group, with isolates having both high and low G+C contents. Cluster XIVab contains both cellulolytic and noncellulolytic members from both human and animal gut, rumen, and environmental sources. In a study of human feces, bacteria that hybridized to the cluster XIVab probe Erec482 constituted 29% of the total (4). We could amplify cluster XIVab genes from sheep rumen fluid but could detect this group in only one landfill site sample (site Br). The clone from landfill site Br (g14Br5) was found to group within XIVa and was most closely related to *C. aminovalericum* (not shown).

The results of this study show that among the clostridial and eubacterial groups tested, the predominant and most ubiquitous groups in landfill are *C. thermocellum* (cluster III) and *C. leptum* (cluster IV). These two groups are closely related, suggesting that they may have derived from a precursor organism that was adapted to the conditions found in many anaerobic environments. For example, the low nitrogen availability in environments such as soils and landfills may select for bacteria with nitrogenase activity, which is present in the cluster III cellulose-degrading strains *C. papyrosolvans* and *C. cellobioparum* (8). The *C. thermocellum* group contains 16S rRNA sequences from only cellulolytic bacteria, which suggests that this phenotype is typical of the group. It is tempting to speculate that sequences obtained using cluster III-specific primers will also have a cellulolytic phenotype. However, to determine if the 16S rRNA sequences obtained using group-specific prim-




FIG. 4. Phylogenetic tree showing relationships of 16S rRNA genes from *Clostridium* cluster IV. Sequences recovered from landfill sites are in boldface. The tree was constructed as described in the legend to Fig. 3.

ers are saccharolytic, they will need to be compared with sequences from metabolically characterized pure-culture isolates.

Molecular detection of clostridial populations has been described for the human intestinal tract (4), but never in a landfill environment. Improved understanding of the microbial populations of landfill sites, and especially the groups involved in initial hydrolysis of waste material, can lead to an overall scheme to monitor bacterial populations involved in anaerobic degradation in situ. Changes in group-specific amplification and TTGE banding patterns can be used to monitor changing conditions at the landfill sites, especially if these techniques target RNA to give some measure of activity. The techniques will also have application to other areas of anaerobic ecology, such as soils, sediments, anaerobic digestors, and the gastrointestinal tract. The application of molecular biological techniques will therefore contribute to our understanding of the ecology of an important bacterial group that has been difficult to isolate and monitor.

Nucleotide sequence accession numbers. The GenBank accession numbers for the nucleotide sequences are AF401533 to AF401549.

This research was funded by the EPSRC/NERC (United Kingdom) Waste and Pollution Management Programme.

We are grateful to Paul Loughnane for technical assistance.

REFERENCES

1. **Barlaz, M. A., D. M. Schaefer, and R. K. Ham.** 1989. Bacterial population development and chemical characteristics of refuse decomposition in a simulated sanitary landfill. *Appl. Environ. Microbiol.* **55**:55–65.
2. **Benson, D. A., I. Karsch-Mizrachi, D. J. Lipman, J. Ostell, B. A. Rapp, and D. L. Wheeler.** 2000. GenBank. *Nucleic Acids Res.* **28**:15–18.
3. **Collins, M. D., P. A. Lawson, A. Willems, J. J. Cordoba, J. Fernandez-Garayzabal, P. Garcia, J. Cai, H. Hippe, and J. A. E. Farrow.** 1994. The phylogeny of the genus *Clostridium*: proposal of five new genera and eleven new species combinations. *Int. J. Syst. Bacteriol.* **44**:812–826.
4. **Franks, A. H., H. J. M. Harmsen, G. C. Raangs, G. J. Jansen, F. Schut, and G. W. Welling.** 1998. Variations of bacterial populations in human feces measured by fluorescent in situ hybridization with group-specific 16S rRNA-targeted oligonucleotide probes. *Appl. Environ. Microbiol.* **64**:3336–3345.
5. **Fry, N. K., J. K. Fredrickson, S. Fishbain, M. Wagner, and D. A. Stahl.** 1997. Population structure of microbial communities associated with two deep, anaerobic, alkaline aquifers. *Appl. Environ. Microbiol.* **63**:1498–1504.
6. **Jukes, T. H., and C. R. Cantor.** 1969. Evolution of protein molecules, p. 21–132. *In* H. N. Munro (ed.), *Mammalian protein metabolism*. Academic Press, New York, N.Y.
7. **Lane, D. J., B. Pace, G. J. Olsen, D. A. Stahl, M. L. Sogin, and N. R. Pace.** 1985. Rapid determination of 16S ribosomal RNA sequences for phylogenetic analyses. *Proc. Natl. Acad. Sci. USA* **82**:6955–6959.
8. **Leschine, S. B.** 1995. Cellulose degradation in anaerobic environments. *Annu. Rev. Microbiol.* **49**:399–426.
9. **Maidak, B. L., J. R. Cole, C. T. Parker, G. M. Garrity, N. Larsen, B. Li, T. G. Lilburn, M. J. McCaughey, G. J. Olsen, R. Overbeek, S. Pramanik, T. M. Schmidt, J. M. Tiedje, and C. R. Woese.** 1999. A new version of the RDP (Ribosomal Database Project). *Nucleic Acids Res.* **27**:171–173.
10. **Palmisano, A. C., D. A. Maruscik, and B. S. Schwab.** 1993. Enumeration of fermentative and hydrolytic micro-organisms from three sanitary landfills. *J. Gen. Microbiol.* **139**:387–391.
11. **Pearson, W. R.** 1990. Rapid and sensitive sequence comparison with FAST and FASTA. *Methods Enzymol.* **183**:63–98.
12. **Rainey, F. A., and P. H. Janssen.** 1995. Phylogenetic analysis by 16S ribosomal DNA sequence comparison reveals two unrelated groups of species within the genus *Ruminococcus*. *FEMS Microbiol. Lett.* **129**:69–74.
13. **Saitou, N., and M. Nei.** 1987. The neighbor-joining method: a new method for constructing phylogenetic trees. *Mol. Biol. Evol.* **4**:406–425.
14. **Van de Peer, Y., and R. De Wachter.** 1994. TREECON for Windows: a software package for the construction and drawing of evolutionary trees for the Microsoft Windows environment. *Comput. Appl. Biosci.* **10**:569–570.
15. **Westlake, K., D. B. Archer, and D. R. Boone.** 1995. Diversity of cellulolytic bacteria in landfill. *J. Appl. Bacteriol.* **79**:73–78.
16. **Woese, C. R.** 1987. Bacterial evolution. *Microbiol. Rev.* **51**:221–271.