

Molecular Characterization of the Leucine Cluster in *Buchnera* sp. Strain PSY, a Primary Endosymbiont of the Aphid *Pemphigus spyrothecae*

Beatriz Sabater-Muñoz,¹ Laura Gómez-Valero,¹ Roeland C. H. J. van Ham,² Francisco J. Silva,¹ and Amparo Latorre^{1*}

Institut Cavanilles de Biodiversitat i Biologia Evolutiva, Universitat de València, València 46071,¹ and Centro de Astrobiología, INTA-CSIC, Madrid,² Spain

Received 21 November 2001/Accepted 5 February 2002

Buchnera strains from most aphid subfamilies studied to date have been found to carry the leucine gene cluster (*leuA*, *-B*, *-C*, and *-D*) on a plasmid, an organization unique among bacteria. Here, however, we demonstrate a classical chromosomal location of the cluster in *Buchnera* sp. strain PSY from the aphid *Pemphigus spyrothecae* (subfamily Pemphiginae). The genes that flank *leuABCD* in *Buchnera* sp. strain PSY appear to be adjacent in the genome of *Buchnera* sp. strain APS, a strain carrying a leucine plasmid. We propose that the presence of a leucine plasmid predates the diversification of symbiotic *Buchnera* and that the chromosomal location observed in *Buchnera* sp. strain PSY arose by a transfer of the leucine genes from a plasmid to the chromosome.

Aphids are plant-sap-feeding insects that maintain a mutualistic association with bacteria from the genus *Buchnera* (Proteobacteria) harbored in specialized host cells called bacteriocytes (1, 11). The origin of the aphid-*Buchnera* association has been estimated at about 200 to 250 million years ago, and host and symbiont lineages have subsequently diverged strictly in parallel (10). The major role of *Buchnera* in the symbiosis is the provision of amino acids, nutrients that are in short supply in phloem sap (4). The complete genome of *Buchnera* from the aphid *Acyrtosiphon pisum* Harris (1776) (*Buchnera* sp. strain APS), has recently been sequenced (13). It is composed of a chromosome of 641 kb and two small plasmids. One of these plasmids contains genes for tryptophan biosynthesis (7, 12, 18), while the other contains genes for leucine biosynthesis (2, 3, 14, 16, 17, 19). Amplification on plasmids of essential amino acid biosynthesis genes is considered indicative of *Buchnera*'s capacity to overproduce these nutrients to the benefit of its host and hence an adaptation to its symbiotic lifestyle (2).

The first leucine plasmid in *Buchnera* (pRPE) was described using *Rhopalosiphum padi* (Linné 1758) (3), a member of the subfamily Aphidinae. It contains genes encoding the key enzymes in the pathway leading to leucine, in the same order as in *Escherichia coli* (*leuABCD*), two copies of the replication initiation gene *repA*, and ORF1, a putative integral membrane protein. Identical plasmids have been shown to be universally present in *Buchnera* from the subfamily Aphidinae (2, 16), while leucine plasmids with slightly different gene orders and/or gene contents are carried by *Buchnera* from the subfamilies Pterocommatinae, Thelaxinae, and Lachninae (14, 17, 19). Finally, *Buchnera* from the subfamily Pemphiginae has been found to carry cryptic plasmids of the same *repA1* family

that do not encode the structural leucine genes. These are very small plasmids containing only an origin of replication, one or two copies of the *repA* gene, and one additional gene (*ibp*, coding for a heat shock protein or ORF1), depending on the tribe within the subfamily (17).

In the present study, we have sequenced the chromosomally encoded leucine genes and flanking regions from *Buchnera* of the aphid *Pemphigus spyrothecae* (Passerini 1856) (*Buchnera* sp. strain PSY), a member of the tribe Pemphigini. This strain carries a small plasmid with the genes *repA1* and *ibp* (19). Why all lineages of *Buchnera* have not undergone amplification of essential amino acid biosynthesis genes via plasmids is an open and unresolved question but is likely to be related to the quality of phloem sap from different host plants and/or the growth rate of the aphids.

Amplification of the leucine gene cluster and flanking regions. *P. spyrothecae* was collected from galls of black poplar trees (*Populus nigra*). DNA was prepared by the cetyltrimethylammonium bromide-NaCl method, after isolation of bacteria from the aphids (5, 19).

Discrimination between plasmid or chromosomal location of the leucine genes, as well as determination of gene order, followed the procedure of van Ham et al. (17), which is based on a combination of structural PCRs (as shown in Fig. 1) and restriction mapping, followed by Southern blot hybridization. This procedure indicates a chromosomal location of the leucine cluster in *Buchnera* sp. strain PSY. The resulting restriction map of the leucine cluster chromosomal region showed a 7.4-kb band generated with *Xba*I that contains the complete cluster and flanking regions (data not shown), selecting this enzyme for the inverse PCR (iPCR) procedure. About 1 µg of *Buchnera* sp. strain PSY DNA was cut with *Xba*I and purified on a Qiagen anion-exchange column, and fragments were ligated with T4 DNA ligase (Stratagene). This DNA was subjected to iPCR, using primers *leuA__du2* (with a *Pst*I restriction site) and *leuA__dl3* (with a *Sal*I restriction site) and

* Corresponding author. Mailing address: Institut Cavanilles de Biodiversitat i Biologia Evolutiva, Universitat de València, Apartat oficial 22085, València 46071, Spain. Phone: (34) 963983649. Fax: (34) 963983670. E-mail: amparo.latorre@uv.es.




FIG. 1. Positions of primers in pRPE plasmid and *Buchnera* sp. strain PSY chromosome (PSY). The table shows combination of primers and results of structural PCRs according to van Ham et al. (17). +, successful amplification; -, no amplification. Primer names: 1, leuA_du2; 2, leuA_dl3; 3, leuD_up2; 4, repAd2.

the Expand Long Template PCR System (Roche) with a GeneAmp PCR System 2400 thermal cycler (Perkin-Elmer). PCR conditions were as follows: 92°C for 2 min; 10 cycles of 92°C for 10 s, 62°C for 30 s, and 68°C for 10 min; this was followed by 20 more cycles with an autoextension of 20 s/cycle at 68°C and a final extension of 68°C for 7 min.

Cloning and sequencing. The 7.0-kb fragment obtained by iPCR was purified with the GeneClean II kit (BIO-101), digested with *Pst*I/*Sal*I, and cloned into pUC18, also digested with *Pst*I/*Sal*I. One selected recombinant clone, named ps10, contained two internal *Eco*RV sites yielding three fragments of 2.6, 3.7, and 0.7 kb plus the vector. The two first fragments were then subcloned into pUC18 *Eco*RV digested and subjected to nested deletions (Nested Deletion kit; AP Biotech) for complete sequencing. The third one was religated and directly sequenced. Sequencing of all clones was carried out with an ABI 377 automated sequencer using the dRhodamine Dye Terminator Cycle Sequencing kit (Perkin-Elmer) and universal primers T3, T7, UNI17-mer, and UNIrev, as well as specific primers designed to close gaps.

The iPCR with outwardly oriented primers within *leuA* lack 389 nucleotides (nt) of the original *leuA* gene. For completion of this sequence, two specific primers were designed and used in a regular PCR to generate a 500-nt fragment containing the missing *leuA* fragment (PsleuA-F, 5'-CGT GAT GGA CAA GCT TTA ACA AG-3'; PsleuA-R, 5'-TGG AAT TGT AAA ACC GAC AGT ATC-3'). The PCR fragment was purified, cloned into pGEM-T Easy vector (Promega), and sequenced as above.

DNA sequence data were assembled with SEQUENCHER, version 4.0 (GeneCodes). BLAST-X, version 2.2.1, was used for gene assignment of ORF proteins (<http://www.ncbi.nlm.nih.gov/BLAST>).

Structure of the *Buchnera* sp. strain PSY leucine gene cluster. The 7.4-kb chromosome region characterized contains seven genes in the following order: *rep-leuA-leuB-leuC-leuD-trxA-rho* (Fig. 2c), whereas in the chromosome of *Buchnera* sp. strain APS, the genes *rep* and *trxA* are adjacent (13) and the leucine gene cluster is located on a plasmid (16).

In *Buchnera* sp. strain PSY, the intergenic region between

leuA and *leuB* is 441 bp long, suggesting a loss of the *leuABCD* operon structure found in most other bacteria. No ORF proteins were found in this region. In contrast, the intergenic regions between *leuB* and *leuC* and between *leuC* and *leuD* are very small (1 and 12 nt, respectively). This suggests that these genes are transcribed as a unique transcription unit, given the lack of space for additional promoters and the absence of putative *Buchnera* promoters in internal coding regions. Putative -35 and -10 promoter sequences, similar to those found for other *Buchnera* genes (17) were only found 42 and 22 nt upstream of *leuA* (TGTTATA and TTAAAAT, respectively). The sequence AGGA, located 7 nt upstream of the initiation codon of *leuA*, corresponds to the Shine Dalgarno sequence. A search for possible terminators of transcription did not show any inverted repeats downstream of *rep* and *leuD*. Such inverted repeats (short inverted repeat type 1) have been found downstream of *leuD* in various leucine plasmids (14).

Two possible scenarios. The different locations of the leucine gene cluster, either on the chromosome as in *Buchnera* sp. strain PSY or on a plasmid as in *Buchnera* sp. strain APS (13), can be explained by two possible scenarios.

(i) The last common ancestor of symbiotic *Buchnera* spp. carried the leucine operon on the chromosome and contained a cryptic plasmid with, at least, the genes *repA1* and *ibp*. After establishment of the symbiosis, the leucine genes were transferred to plasmids independently in several *Buchnera* lineages, resulting in leucine plasmids with different gene orders and/or gene contents (17, 19).

(ii) Alternatively, the transfer of the gene cluster to a *repA1* plasmid took place only once in the common ancestor of extant *Buchnera* lineages. Back transfer to the chromosome took place in the lineage leading to *Buchnera* sp. strain PSY, involving a recombination event between two direct repeat sequences, one in the intergenic region between the chromosomal genes *rep* and *trxA* and the other between the genes *leuD* and *leuA* of the plasmid (see Fig. 2).

Established aphid phylogenies, which postulate a basal split between the subfamily Pemphiginae and all remaining *Buchnera* lineages (2, 6), support the first scenario. *E. coli*, a close free-living relative of *Buchnera*, carries the leucine gene cluster




FIG. 2. The three steps of back transfer scenario. (a) ancestral chromosome; (b) intermediate stage; (c) *Buchnera* sp. strain PSY chromosome. The black line indicates the fragment sequenced; short open arrows indicate tRNA (13).

on the chromosome, in the order *leuA*, *-B*, *-C*, and *-D*. Although this organization is similar to the one described here for *Buchnera* sp. strain PSY, it is difficult to explain the great size (441 bp) of the intergenic region between *leuA* and *leuB* in the first scenario. Recent studies have estimated the median size of intergenic regions in *Buchnera* sp. strain APS to be 74 nt (9, 15).

The second scenario seems most plausible in the light of recent molecular phylogenetic studies. These have extended taxonomic sampling relative to previous studies and have questioned traditional aphid phylogenies (8, 17, 19). Most significantly, the subfamily Lachninae was shown to be basal to all other lineages, contrary to such a previous placement for the subfamily Pemphiginae. *Buchnera* strains from the subfamily Lachninae contain a leucine plasmid. A possible scenario of back transfer of the leucine gene cluster to the chromosome in *Buchnera* sp. strain PSY is presented in Fig. 2. This back transfer may have been followed by disintegration of some of the genes inserted between *leuA* and *leuB* (9, 15), leaving as a remnant of the process the relatively large intergenic region observed between *leuA* and *leuB*. Initially, nonrecombinant plasmid copies may have retained the leucine genes, but eventually these too may have disappeared through gene disintegration. Strikingly similar to the *leuA-leuB* intergenic region, the noncoding region between *ibp* and *repA1*, encoded by the plasmid of *Buchnera* sp. strain PSY, is unusually large, measuring 858 bp (19). This region might correspond to a remnant sequence of the leucine gene cluster's disintegration. Retention of the small, *ibp*-carrying plasmid could be due to disintegration of the chromosomal copy of this essential gene.

Nucleotide sequence accession number. The nucleotide sequences reported in this paper have been deposited in the GenBank and EMBL databases under accession no. AJ426489.

Financial support was provided by grants GR00-27 from Conselleria de Cultura y Educació, Comunidad Valenciana, to A.L. and BFM2000-1383 from Ministerio de Ciencia y Tecnología (MCYT) to A.L., F.J.S., and B.S.-M.

We are very grateful to J. M. Michelena for the identification of aphid species and to A. Moya for valuable comments on the manuscript. The facilities of the Servei Central de Suport a la Investigació Experimental of the University of València have been used.

REFERENCES

- Baumann, L., P. Baumann, C. Y. Lai, D. Rouhbakhsh, N. A. Moran, and M. A. Clark. 1995. Genetics, physiology, and evolutionary relationships of the genus *Buchnera*: intracellular symbionts of aphids. *Annu. Rev. Microbiol.* **49**:55–94.
- Baumann, L., P. Baumann, N. A. Moran, J. Sandström, and M. L. Thao. 1999. Genetic characterization of plasmids containing genes encoding enzymes of leucine biosynthesis in endosymbionts (*Buchnera*) of aphids. *J. Mol. Evol.* **48**:77–85.
- Bracho, A. M., D. Martínez-Torres, A. Moya, and A. Latorre. 1995. Discovery and molecular characterization of a plasmid localized in *Buchnera* sp. bacterial endosymbiont of the aphid *Rhopalosiphum padi*. *J. Mol. Evol.* **41**:67–73.
- Douglas, A. E. 1998. Nutritional interactions in insect-microbial symbioses: aphids and their symbiotic bacteria *Buchnera*. *Annu. Rev. Entomol.* **43**:17–37.
- Harrison, C. P., A. E. Douglas, and A. F. G. Dixon. 1989. A rapid method to isolate symbiotic bacteria from aphids. *J. Invertebr. Pathol.* **53**:427–428.
- Heie, O. E. 1987. Paleontology and phylogeny, p. 367–391. In A. K. Minks and P. Harrewijn (ed.), *World Crop Pests*, vol. 2A. Aphids: their biology, natural enemies and control. Elsevier, Amsterdam, The Netherlands.
- Lai, C.-Y., L. Baumann, and P. Baumann. 1994. Amplification of *trpEG*: adaptation of *Buchnera aphidicola* to an endosymbiotic association with aphids. *Proc. Natl. Acad. Sci. USA* **91**:3819–3823.
- Martínez-Torres, D., C. Buades, A. Latorre, and A. Moya. 2001. Molecular systematics of aphids and their primary endosymbionts. *Mol. Phylogenet. Evol.* **20**:437–449.
- Mira, A., H. Ochman, and N. Moran. 2001. Deletion bias and the evolution of bacterial genomes. *Trends Genet.* **17**:589–596.
- Moran, N. A., M. A. Munson, P. Baumann, and H. Ishikawa. 1993. A molecular clock in endosymbiotic bacteria is calibrated using the insect host. *Proc. R. Soc. Lond. B Biol. Sci.* **253**:167–171.
- Munson, M. A., M. A. Clark, L. Baumann, N. A. Moran, D. J. Voegtlin, and B. C. Campbell. 1991. Evidence for the establishment of aphid-eubacterium

- endosymbiosis in an ancestor of four aphid families. *J. Bacteriol.* **173**:6321–6324.
12. Rouhbakhsh, D., C.-Y. Lai, C. von Dohlen, M. A. Clark, L. Baumann, P. Baumann, N. A. Moran, and D. J. Voegtlin. 1996. The tryptophan biosynthesis pathway of aphid endosymbionts (*Buchnera*): genetics and evolution of plasmid-associated anthranilate synthase (*trpEG*) within the aphididae. *J. Mol. Evol.* **42**:414–421.
 13. Shigenobu, S., H. Watanabe, M. Hatori, Y. Sakaki, and H. Ishikawa. 2000. Genome sequence of the endocellular bacterial symbiont of aphids *Buchnera* sp. *APS. Nature* **407**:81–86.
 14. Silva, F. J., R. C. H. J. van Ham, B. Sabater, and A. Latorre. 1998. Structure and evolution of the leucine plasmids carried by the endosymbiont (*Buchnera aphidicola*) from aphids of the family Aphididae. *FEMS Microbiol. Lett.* **168**:43–49.
 15. Silva, F. J., A. Latorre, and A. Moya. 2001. Genome size reduction through multiple events of gene disintegration in *Buchnera* APS. *Trends Genet.* **17**:615–618.
 16. Soler, T., A. Latorre, B. Sabater, and F. J. Silva. 2000. Molecular characterization of the leucine plasmid from *Buchnera aphidicola*, primary endosymbiont of the aphid *Acyrtosiphon pisum*. *Curr. Microbiol.* **40**:264–268.
 17. van Ham, R. C. H. J., A. Moya, and A. Latorre. 1997. Putative evolutionary origin of plasmids carrying the genes involved in leucine biosynthesis in *Buchnera aphidicola* (endosymbiont of aphids). *J. Bacteriol.* **179**:4768–4777.
 18. van Ham, R. C. H. J., D. Martínez-Torres, A. Moya, and A. Latorre. 1999. Plasmid-encoded anthranilate synthase (*TrpEG*) in *Buchnera aphidicola* from aphids of the family Pemphigidae. *Appl. Environ. Microbiol.* **65**:117–125.
 19. van Ham, R. C. H. J., F. González-Candela, F. J. Silva, B. Sabater, A. Moya, and A. Latorre. 2000. Postsymbiotic plasmid acquisition and evolution of the *repA1*-replicon in *Buchnera aphidicola*. *Proc. Natl. Acad. Sci. USA* **97**:10855–10860.