

# Noninvasive Measurement of Bacterial Intracellular pH on a Single-Cell Level with Green Fluorescent Protein and Fluorescence Ratio Imaging Microscopy

Katja N. Olsen,<sup>1,2</sup> Birgitte B. Budde,<sup>2</sup> Henrik Siegumfeldt,<sup>2</sup> K. Björn Rechinger,<sup>2†</sup>  
Mogens Jakobsen,<sup>2</sup> and Hanne Ingmer<sup>1\*</sup>

Department of Veterinary Microbiology<sup>1</sup> and Department of Dairy and Food Science,<sup>2</sup> The Royal Veterinary and Agricultural University, DK-1870 Frederiksberg C, Denmark

Received 17 January 2002/Accepted 14 May 2002

**We show that a pH-sensitive derivative of the green fluorescent protein, designated ratiometric GFP, can be used to measure intracellular pH ( $pH_i$ ) in both gram-positive and gram-negative bacterial cells. In cells expressing ratiometric GFP, the excitation ratio (fluorescence intensity at 410 and 430 nm) is correlated to the  $pH_i$ , allowing fast and noninvasive determination of  $pH_i$  that is ideally suited for direct analysis of individual bacterial cells present in complex environments.**

Bacteria are often subjected to various forms of environmental stress, and in recent years, there has been increasing focus on the different mechanisms they employ to protect themselves against these environmental changes. One of the most extensively studied stress responses is the ability of bacteria to survive low pH by adjusting their intracellular pH ( $pH_i$ ) in response to changes in extracellular pH ( $pH_{ex}$ ) (1, 4). Several methods have been developed for measuring bacterial  $pH_i$  (2, 14, 17, 19). However, as these techniques require radioactive labeling and time-consuming staining procedures, noninvasive methods for continuous measurement of  $pH_i$  in bacteria are in demand. Since the discovery of green fluorescent protein (GFP), a wide range of mutant variants has been created. In eukaryotic cells, several GFP variants with altered excitation and emission spectra have been examined for their use as  $pH_i$  probes (8, 11, 12, 13, 16). One of these GFP variants, ratiometric GFP, was obtained by introducing specific amino acid substitutions to the chromophore, causing the resulting protein to alter its excitation spectrum according to the pH of the surrounding environment (13).

In order to express the ratiometric GFP protein in both gram-positive and gram-negative bacterial cells, we inserted the corresponding gene downstream of the P32 promoter in the chloramphenicol-resistant expression vector pMG36c, which replicates in both cell types (20). The ratiometric *gfp* gene (GenBank accession no. AF058694) was amplified by PCR by using the oligonucleotides C-GFP (5'-TAT CCC AAG CTT TTA TTT GTA TAG TTC ATC CAT GCC ATG TG-3') and N-GFP (5'-TGC TCT AGA GTA ATA AGG AGG AAA AAA TAT GAG TAA AGG AGA AGA ACT TTT CAC TGG AGT TGT CCC-3') (DNA Technology, Århus, Denmark) that additionally introduced an initiating ATG codon as well as a ribosomal binding site (5). The resulting 750-bp DNA

fragment was inserted into the *Xba*I- and *Eco*RI-digested pMG36c, and the correct DNA sequence of the ratiometric *gfp* gene present in the resulting plasmid (pGFP<sub>ratiometric</sub>) was confirmed by DNA sequence analysis (data not shown).

When we introduced pGFP<sub>ratiometric</sub> in the gram-positive bacterium *Lactococcus lactis* subsp. *lactis* CNRZ 157 as previously described (9) and grew cells at 30°C in M17 broth containing 0.5% (wt/vol) glucose and 5 µg of chloramphenicol per ml, we found that excitation at 410 nm gave a strong pH-dependent fluorescent signal. Furthermore, a pH-independent isosbestic point was seen at 430 nm, which is in accordance with previous results obtained in mammalian cells (13 and data not shown). It was therefore concluded that the excitation ratio, fluorescence intensity at 410 and 430 nm ( $R_{410/430}$ ), is suitable as a measure of  $pH_i$ .

To correlate  $R_{410/430}$  with  $pH_i$ , *L. lactis* cells carrying pGFP<sub>ratiometric</sub> were permeabilized with nisin (3, 7), which disrupts the proton gradient across the cytoplasmic membrane and allows the internal pH to become identical to the external pH. Correspondingly, *Escherichia coli* TOP 10 cells (Invitrogen) were grown at 37°C in Luria-Bertani broth, transformed with pGFP<sub>ratiometric</sub> (15), and pH-equilibrated by use of carbonyl cyanide-*m*-chlorophenylhydrazone (CCCP) (6). Subsequently, overnight cultures expressing ratiometric GFP were harvested by centrifugation (10,000 × *g* for 5 min), washed twice in potassium phosphate buffer (pH 7.5) containing glucose (10 mM), and finally resuspended in potassium phosphate buffer (10 mM glucose) with appropriate pH for 30 min. One hundred microliters of the bacterial suspension (approximately 10<sup>8</sup> cells) was applied to a coverslip coated with 0.01% (wt/vol) poly-L-lysine (Sigma) and allowed to settle. Prior to being used, the coverslips were cleaned by overnight submersion in chrome sulfuric acid (Sigma) and then rinsed in distilled water and stored in 70% (vol/vol) ethanol. Unattached bacteria were removed by rinsing with buffer. The fluorescence microscopy setup has previously been described (19), except that the emission was recorded on a Coolsnap fx charge-coupled device camera (Roper Scientific, Trenton, Pa.). The dichroic mirrors were 380 and 510 nm for ratiometric GFP and cFDAse [fluor-

\* Corresponding author. Mailing address: Department of Veterinary Microbiology, The Royal Veterinary and Agricultural University, Stigbøjlen 4, DK-1870 Frederiksberg C, Denmark. Phone: 45 35282773. Fax: 45 35282757. E-mail: ingmer@biobase.dk.

† Present address: Foss Electric A/S, DK-3400 Hillerød, Denmark.




FIG. 1. Correlation between  $\text{pH}_i$  and  $R_{410/430}$  of ratiometric GFP in pH-equilibrated cells. *L. lactis* and *E. coli* single cells were equilibrated with 10 kIU of nisin  $\text{ml}^{-1}$  (closed circles) and 10  $\mu\text{M}$  CCCP (closed triangles), respectively. Each point represents the mean value for 20 individual cells, with error bars indicating the standard deviations.

rochrome 5(6)-carboxyfluorescein diacetate succinimidyl ester], respectively, and the emission band-pass filters were 500 to 530 nm and 515 to 565 nm for ratiometric GFP and cFDase, respectively. Images were stored on a personal computer by using Metafluor 4.5 (Universal Imaging, Downingtown, Pa.), and data from regions of interest (i.e., single cells) were logged into a spreadsheet (Excel). In each experiment, at least 20 single cells were analyzed. Each experiment was carried out in duplicate.

On the basis of the results obtained with nisin- or CCCP-treated cells, a calibration curve was constructed by plotting  $R_{410/430}$  versus the pH of equilibrated cells in the pH range of 5.5 to 8.5 (Fig. 1). When *L. lactis* cells were permeabilized with nisin, we found that the relationship between  $R_{410/430}$  and  $\text{pH}_i$  was similar to that observed when *E. coli* cells treated with CCCP were investigated.

Subsequently, we used the calibration curve (Fig. 1) to determine  $\text{pH}_i$  in single bacterial cells that had been resuspended in 50 mM potassium phosphate buffers with various pHs (Fig. 2). Examination of the  $\text{pH}_i$  of *L. lactis* cells revealed that the pH gradient ( $\Delta\text{pH} = \text{pH}_i - \text{pH}_{\text{ex}}$ ) is approximately 1.8 at a  $\text{pH}_{\text{ex}}$  of 5.5. Increasing the  $\text{pH}_{\text{ex}}$  caused a decrease in the  $\Delta\text{pH}$ , which finally became negative at a  $\text{pH}_{\text{ex}}$  of 8.5. In order to validate these results, the pH measurements were also carried out by using the conventional pH-sensitive cFDase. *L. lactis* cells stained with cFDase were analyzed as previously described (19), except that the pH equilibration was performed with nisin as described above.

The results presented in Fig. 2 show a good correlation ( $R = 0.987$ ) between the pH responses obtained with the two methods. The observed deviations possibly result from the fact that both methods rely on a calculation of  $\text{pH}_i$  based on standard curves, which introduces a variation of 0.1 to 0.2 pH units (18).




FIG. 2. Determination of *L. lactis*  $\text{pH}_i$  as a function of  $\text{pH}_{\text{ex}}$  by use of ratiometric GFP and cFDase. The  $\text{pH}_i$  of cells transformed with ratiometric GFP (closed circles) was obtained by measuring the  $R_{410/430}$  and converting the obtained value to the correlating  $\text{pH}_i$ . For validation,  $\text{pH}_i$  was also obtained by staining the cells with cFDase (open circles) and measuring the  $R_{490/435}$ . Each point represents the mean value for 20 individual cells, with error bars indicating the standard deviations.

In addition, heterogeneities in the populations will influence the standard deviation. Therefore, we conclude that ratiometric GFP can be used as a  $\text{pH}_i$  probe for bacterial cells. When  $\text{pH}_i$  was measured in *E. coli* cells exposed to various pH values, we found that a large  $\Delta\text{pH}$  of approximately 1.5 was maintained at a low  $\text{pH}_{\text{ex}}$ , gradually decreasing with increasing  $\text{pH}_{\text{ex}}$  until finally becoming 0 at a  $\text{pH}_{\text{ex}}$  of 8 (Fig. 3). For both bacteria, we obtained similar standard deviations (Fig. 2 and 3), suggesting that ratiometric GFP is equally applicable for measurements with gram-negative and gram-positive bacteria.

To ensure that the presence of pGFP-ratiometric did not affect the growth of bacterial cells, we followed the growth of *L. lactis* and *E. coli* cells in the presence or absence of the plasmid and found no effect with *L. lactis*, while the presence of the plasmid in *E. coli* cells slightly reduced the growth rate (data not shown). We also investigated the stability of pGFP-ratiometric as previously described (10) and found that 74% of *L. lactis* and 54% of *E. coli* cells retained the plasmid after 60 generations, demonstrating that studies can be conducted in the absence of antibiotic selection of the plasmid. Furthermore, we investigated whether the type of acid used for acidification affects the ratiometric GFP excitation spectrum when expressed in *L. lactis* cells and found the same excitation spectrum with acetic, citric, or formic acid as we found with hydrochloric acid (data not shown). This result indicates that the method is also applicable when the pH is adjusted with organic acids.

In conclusion, we have applied a pH-sensitive derivative of the GFP, ratiometric GFP (13), for measurement of  $\text{pH}_i$  in single bacterial cells in the pH range of 5.5 to 8.5. While this




FIG. 3. Determination of *E. coli* p $H_i$  as a function of p $H_{ex}$ . Cells expressing ratiometric GFP were resuspended in phosphate buffers at pHs ranging from 5.5 to 8.5. The  $R_{410/430}$  was determined and was transformed to the corresponding p $H_i$ . Each point represents the mean value for 20 individual cells, with error bars indicating the standard deviations.

technique requires that the cells of interest express the ratiometric GFP protein, it provides a fast, noninvasive, and continuous determination of p $H_i$  which is ideally suited for kinetic studies of both gram-negative and gram-positive bacteria. As the ratio between the fluorescence intensities at two different excitation wavelengths is used, the determination is independent of the individual concentration of ratiometric GFP in each cell. Furthermore, we show that the excitation spectrum of ratiometric GFP when cells were acidified with various organic acids was identical to that when cells were acidified with hydrochloric acid, suggesting that the ratiometric GFP is well suited for studies of the p $H_i$  in bacteria residing in natural environments such as food products.

We thank G. Miesenböck for providing the ratiometric *gfp* gene, J. Kok for providing pMG36c, and M. Chopin for providing *L. lactis* subsp. *lactis* CNRZ 157. A. Arndal and C. Friis are thanked for excellent technical assistance. We thank J. Delves-Broughton (Aplin and Barrett Ltd., Danisco Cultor, Beaminster, England) for providing the purified nisin.

## REFERENCES

- Booth, I. R. 1985. Regulation of cytoplasmic pH in bacteria. *Microbiol. Rev.* **49**:359–378.
- Breuwer, P., J.-L. Drocourt, F. M. Rombouts, and T. Abee. 1996. A novel method for continuous determination of the intracellular pH in bacteria with the internally conjugated fluorescent probe 5 (and 6-)carboxyfluorescein succinimidyl ester. *Appl. Environ. Microbiol.* **62**:178–183.
- Budde, B. B., and M. Jakobsen. 2000. Real-time measurements of the interaction between single cells of *Listeria monocytogenes* and nisin on a solid surface. *Appl. Environ. Microbiol.* **66**:3586–3591.
- Cook, G. M., and J. B. Russell. 1994. The effect of extracellular pH and lactic acid on pH homeostasis in *Lactococcus lactis* and *Streptococcus bovis*. *Curr. Microbiol.* **28**:165–168.
- Denoya, C. D., D. H. Bechhofer, and D. Dubnau. 1986. Translational auto-regulation of ermC 23S rRNA methyltransferase expression in *Bacillus subtilis*. *J. Bacteriol.* **168**:1133–1141.
- Diez-Gonzalez, F., and J. B. Russell. 1997. Effects of carbonylcyanide-*m*-chlorophenylhydrazone (CCCP) and acetate on *Escherichia coli* O157:H7 and K-12: uncoupling versus anion accumulation. *FEMS Microbiol. Lett.* **151**:71–76.
- Duffes, F., P. Jenoe, and P. Boyaval. 2000. Use of two-dimensional electrophoresis to study differential protein expression in divercin V41-resistant and wild-type strains of *Listeria monocytogenes*. *Appl. Environ. Microbiol.* **66**:4318–4324.
- Elsiger, M. A., R. M. Wachter, G. T. Hanson, K. Kallio, and S. J. Remington. 1999. Structural and spectral response of green fluorescent protein variants to changes in pH. *Biochemistry* **38**:5296–5301.
- Holo, H., and I. F. Nes. 1989. High-frequency transformation, by electroporation, of *Lactococcus lactis* subsp. *cremoris* grown with glycine in osmotically stabilized media. *Appl. Environ. Microbiol.* **55**:3119–3123.
- Ingmer, H., and S. N. Cohen. 1993. Excess intracellular concentration of pSC101 RepA protein interferes with both plasmid DNA replication and partitioning. *J. Bacteriol.* **175**:7834–7841.
- Karagiannis, J., and P. G. Young. 2001. Intracellular pH homeostasis during cell-cycle progression and growth state transition in *Schizosaccharomyces pombe*. *J. Cell Sci.* **114**:2929–2941.
- Kneen, M., J. Farinas, Y. Li, and A. S. Verkman. 1998. Green fluorescent protein as a noninvasive intracellular pH indicator. *Biophys. J.* **74**:1591–1599.
- Miesenböck, G., D. A. De Angelis, and J. E. Rothman. 1998. Visualizing secretion and synaptic transmission with pH-sensitive green fluorescent proteins. *Nature* **394**:192–195.
- Nannen, N. L., and R. W. Hutkins. 1991. Intracellular pH effects in lactic acid bacteria. *J. Dairy Sci.* **74**:741–746.
- O'Callaghan, D., and A. Charbit. 1990. High-efficiency transformation of *Salmonella typhimurium* and *Salmonella typhi* by electroporation. *Mol. Gen. Genet.* **223**:156–158.
- Robey, R. B., O. Ruiz, A. V. P. Santos, J. Ma, F. Kear, L. J. Wang, C. J. Li, A. A. Bernardo, and J. A. L. Arruda. 1998. pH-dependent fluorescence of a heterologously expressed *Aequorea* green fluorescent protein mutant: in situ spectral characteristics and applicability to intracellular pH estimation. *Biochemistry* **37**:9894–9901.
- Roe, A. J., D. McLaggan, I. Davidson, C. O'Byrne, and I. R. Booth. 1998. Perturbation of anion balance during inhibition of growth of *Escherichia coli* by weak acids. *J. Bacteriol.* **180**:767–772.
- Siegmund, H., K. B. Rechinger, and M. Jakobsen. 1999. Use of fluorescence ratio imaging for intracellular pH determination of individual bacterial cells in mixed cultures. *Microbiology* **145**:1703–1709.
- Siegmund, H., K. B. Rechinger, and M. Jakobsen. 2000. Dynamic changes of intracellular pH in individual lactic acid bacterium cells in response to a rapid drop in extracellular pH. *Appl. Environ. Microbiol.* **66**:2330–2335.
- van de Guchte, M., J. M. van der Vossen, J. Kok, and G. Venema. 1989. Construction of a lactococcal expression vector: expression of hen egg white lysozyme in *Lactococcus lactis* subsp. *lactis*. *Appl. Environ. Microbiol.* **55**:224–228.