

Establishment of an Arbitrary PCR for Rapid Identification of Tn917 Insertion Sites in *Staphylococcus epidermidis*: Characterization of Biofilm-Negative and Nonmucoid Mutants

Johannes K.-M. Knobloch,* Max Nedelmann, Kathrin Kiel, Katrin Bartscht, Matthias A. Horstkotte, Sabine Dobinsky, Holger Rohde, and Dietrich Mack

Institut für Medizinische Mikrobiologie und Immunologie, Universitätsklinikum Hamburg-Eppendorf, D-20246 Hamburg, Germany

Received 14 March 2003/Accepted 15 July 2003

Transposon mutagenesis with the *Enterococcus faecalis* transposon Tn917 is a genetic approach frequently used to identify genes related with specific phenotypes in gram-positive bacteria. We established an arbitrary PCR for the rapid and easy identification of Tn917 insertion sites in *Staphylococcus epidermidis* with six independent, well-characterized biofilm-negative Tn917 transposon mutants, which were clustered in the *icaADBC* gene locus or harbor Tn917 in the regulatory gene *rsbU*. For all six of these mutants, short chromosomal DNA fragments flanking both transposon ends could be amplified. All fragments were sufficient to correctly identify the Tn917 insertion sites in the published *S. epidermidis* genomes. By using this technique, the Tn917 insertion sites of three not-yet-characterized biofilm-negative or nonmucoid mutants were identified. In the biofilm-negative and nonmucoid mutant M12, Tn917 is inserted into a gene homologous to the regulatory gene *purR* of *Bacillus subtilis* and *Staphylococcus aureus*. The Tn917 insertions of the nonmucoid but biofilm-positive mutants M16 and M20 are located in genes homologous to components of the phosphoenolpyruvate-sugar phosphotransferase system (PTS) of *B. subtilis*, *S. aureus*, and *Staphylococcus carnosus*, indicating an influence of the PTS on the mucoid phenotype in *S. epidermidis*.

The *Enterococcus faecalis* transposon Tn917 is a well-characterized tool in genetic manipulation of gram-positive bacteria. In a variety of bacteria, such as *Bacillus* species, *Streptococcus* species, enterococci, *Listeria monocytogenes*, and staphylococci, Tn917 was used to identify genes related to characteristic phenotypes or metabolic pathways (15, 25, 36, 46, 49, 51, 55). In the era before the sequencing of complete bacterial genomes, transposon insertion sites were analyzed by cloning or inverted PCR requiring the effort of chromosomal mapping. In recent years several bacterial genomes could be deciphered, including major gram-positive pathogens such as *Staphylococcus aureus*, *Staphylococcus epidermidis*, *L. monocytogenes*, and *Streptococcus pneumoniae*, as well as the gram-positive model organism *B. subtilis*. Therefore, in the post-genomic era short nucleotide or protein sequences are sufficient for the identification of genes in the respective genomes.

In *S. epidermidis* biofilm formation is the major pathogenic factor in foreign body-associated infections and leads to increased resistance against antibiotics (14, 22, 26). Biofilm-forming *S. epidermidis* strains display a mucoid phenotype with large humid colonies under certain culture conditions, as is observed for other bacteria expressing exopolysaccharides. Tn917 mutagenesis identified several genes involved in biofilm formation of *S. epidermidis*. Five of these mutants are clustered

with different orientations of Tn917 in the *icaADBC* gene locus (29, 30), which is coding for four synthetic enzymes of the polysaccharide intercellular adhesin (PIA) (13, 16). PIA is essential for cell accumulation during biofilm formation in *S. epidermidis* (28). PIA and biofilm formation as a virulence factor in *S. epidermidis* could be demonstrated by different animal models (43–45). In the sixth biofilm-negative and nonmucoid mutant, the Tn917 insertion is located in the first gene *rsbU* of the σ^B operon coding for a positive regulator of the alternative sigma factor σ^B (19). The lack of *icaADBC* transcription (30) in this mutant indicates a regulatory influence of σ^B on biofilm formation in *S. epidermidis*. In addition, transduction of this mutation in a methicillin-resistant genetic background leads to reduced susceptibility against oxacillin (31), which is, like biofilm formation, associated with invasive *S. epidermidis* strains (12). The regulation of biofilm formation by σ^B was confirmed recently by allelic gene replacement (J. K.-M. Knobloch et al., unpublished data). Indeed, biofilm formation of *S. epidermidis* is induced by a variety of environmental stress factors, including antibiotics and disinfectants (9, 11, 20, 39, 41, 42). However, at the moment it is still unknown which of these stress factors lead to induction of biofilm formation by activation of σ^B . Recently, the *icaADBC* gene locus was also described in *S. aureus* (8, 9, 33), and the σ^B dependence of biofilm formation in this staphylococcal species could be demonstrated (40). However, the regulation of biofilm formation seems to be quite different in these two staphylococcal species (21), and the specific differences should be further characterized.

Apparently, screening of *S. epidermidis* Tn917 mutants iden-

* Corresponding author. Mailing address: Institut für Medizinische Mikrobiologie und Immunologie, Universitätsklinikum Hamburg-Eppendorf, Martinistr. 52, D-20246 Hamburg, Germany. Phone: 49-40-42803-3147. Fax: 49-40-42803-4881. E-mail: knobloch@uke.uni-hamburg.de.

TABLE 1. Bacterial strains and plasmids

Strain	Comment(s)	Source or reference
1457	Isolate from infected central venous catheter harboring cryptic plasmid p1457	32, 35
1457c/pTV1ts	1457 cured of a cryptic plasmid and containing the temperature-sensitive plasmid pTV1ts harboring transposon Tn917	35
1457-M10	Isogenic biofilm-negative, nonmucooid Tn917 mutant of <i>S. epidermidis</i> 1457c (class I)	29
1457-M11	Isogenic biofilm-negative, nonmucooid Tn917 mutant of <i>S. epidermidis</i> 1457c (class I)	29
M12	Isogenic biofilm-negative, nonmucooid Tn917 mutant of <i>S. epidermidis</i> 1457c (class II)	30
M15	Isogenic biofilm-negative, nonmucooid Tn917 mutant of <i>S. epidermidis</i> 1457c (class III)	19, 30
M16	Isogenic biofilm-positive, nonmucooid Tn917 mutant of <i>S. epidermidis</i> 1457c	This study
M20	Isogenic biofilm-positive, nonmucooid Tn917 mutant of <i>S. epidermidis</i> 1457c	This study
M21	Isogenic biofilm-negative, nonmucooid Tn917 mutant of <i>S. epidermidis</i> 1457c (class I)	30
M22	Isogenic biofilm-negative, nonmucooid Tn917 mutant of <i>S. epidermidis</i> 1457c (class I)	30
M24	Isogenic biofilm-negative, nonmucooid Tn917 mutant of <i>S. epidermidis</i> 1457c (class I)	30

tifies gene loci of interest for virulence-associated phenotypes. To rapidly identify Tn917 insertions, we established in the present study an arbitrary PCR technique for rapid and easy identification of Tn917 insertion sites by investigation of these six well-characterized independent biofilm-negative and nonmucooid Tn917 mutants of *S. epidermidis* 1457 (19, 30). In addition, by using the newly established arbitrary PCR method, the Tn917 insertion sites of the biofilm-negative and nonmucooid mutant M12 (30) and two newly isolated nonmucooid but biofilm-positive mutants M16 and M20 were identified.

MATERIALS AND METHODS

Bacterial strains, growth conditions, and isolation of chromosomal DNA. The bacterial strains used in the present study are listed in Table 1. For the detection of biofilm formation, *S. epidermidis* cells were grown in 96-well tissue culture plates (NunclonDelta; Nunc, Roskilde, Denmark) or in Lab-Tek Chambered Coverglass cell culture system (Nalge Nunc, Naperville, Ill.) in Trypticase soy broth (TSB_{BBL}; Becton Dickinson, Cockeysville, Md.) at 37°C. Biofilm formation was quantified after staining with gentian violet by a BEPII ELISA reader (Behring), as described recently (27). For the detection of a mucooid phenotype, cells were grown on purple agar (Difco, Detroit, Mich.) supplemented with 0.4% *N*-acetylglucosamine (Sigma, St. Louis, Mo.). To investigate phenotypes on congo red agar (CRA) cells were grown on TSB_{BBL} supplemented with 1% glucose and 1% agar (Becton Dickinson), 0.08% Congo red (CRA_{TSB}; Merck, Darmstadt, Germany), and brain heart infusion broth (Oxoid) supplemented with 3.6% sucrose, 1% agar, and 0.08% Congo red (CRA_{BHI}) as described recently (21). Isolation of chromosomal DNA of *S. epidermidis* was performed as described previously (27). Erythromycin was used at a concentration of 150 µg/ml for selection of Tn917 mutants.

Transposon mutagenesis and phage transduction. Transposon mutagenesis was performed with the temperature-sensitive plasmid pTV1ts containing transposon Tn917 as described previously (27). *S. epidermidis* 1457c/pTV1ts was grown under nonpermissive temperature for pTV1ts under erythromycin selection, and the resulting mutants were screened for a biofilm-negative and/or

nonmucooid phenotype. Phage transduction of Tn917 insertions into wild-type *S. epidermidis* 1457 were performed with phage 71, kindly provided by V. T. Rosdahl, Statens Serum Institut, Copenhagen, Denmark, as described previously (27).

Arbitrary primed PCR and nucleotide sequence analysis. Amplification of short DNA-fragments was performed by using the DyNzyme DNA polymerase kit (Finzyme, Espoo, Finland) as described by the manufacturer. Oligonucleotides specific for the 5' end (917-5.1, 917-5.2, and 917-5.3; Table 2) and the 3 end of Tn917 (917-3.1, 917-3.2, and 917-3.3; Table 2), as well as the arbitrary primers (STAPHarb1, STAPHarb2, and arb3; Table 2), were synthesized by MWG Biotech (Munich, Germany). All PCRs were performed in a Primus 96 Thermocycler (MWG Biotech) without a mineral oil overlay.

The PCR of the first round was performed in a final volume of 50 µl. An arbitrary primer (STAPHarb1 or STAPHarb2; 200 pmol/reaction) was paired with one of the proximal Tn917 primers (homologous to the external nested-PCR primers [917-5.3, 917-5.2, 917-3.3, or 917-3.2]; 10 pmol/reaction). Portions (1 µl) of the chromosomal DNA preparations were used as templates, and PCR was performed under the following conditions: 95°C for 5 min; six cycles of 94°C for 30 s, 30°C for 30 s, and 72°C for 1 min; 30 cycles of 94°C for 30 s, 45°C for 30 s, and 72°C for 1 min; and finally 72°C for 5 min. Samples were then kept at 4°C.

The second round of PCR was performed in a final volume of 100 µl with 5 µl of the PCR of round 1 as a template. Primer arb3 (20 pmol/reaction) was paired with one of the respective distal Tn917 primers (homologous to internal nested PCR primers [917-5.2, 917-5.1, 917-3.2, or 917-3.1]; 20 pmol/reaction), and PCR was performed under the following conditions: 30 cycles of 94°C for 30 s, 45°C for 30 s, and 72°C for 1 min, followed by 72°C for 5 min. The samples were then kept at 4°C. The results of this PCR was visualized by agarose gel electrophoresis, and PCRs containing at least one distinct visible fragment were used for further characterization.

Nucleotide sequence analysis was performed by using the second PCR directly with the respective Tn917 internal primers on an ABI Prism 310 sequencer with capillary electrophoresis using the ABI Prism dGTP BigDye terminator ready reaction kit (PE Applied Biosystems, Foster City, Calif.), with 5 to 10 µl of the PCR as a template. Nucleotide sequences were analyzed subsequently with vector NTI suite II software. For identification of the Tn917 insertion sites, a basic local alignment search tool (BLAST) search with the sequence represent-

TABLE 2. Oligonucleotides used as primers

Primer	Localization (bp) ^a	Primer sequence
917-5.1	124–103	5'-ATC GAT ACA AAT TCC TCG TAG G-3'
917-5.2	274–255	5'-AAC CGT TAC CTG TTT GTG CC-3'
917-5.3	353–333	5'-CCA ATC ACT CTC GGA CAA TAC-3'
917-3.1	5298–5318	5'-TTT AGT GGG AAT TTG TAC CCC-3'
917-3.2	5194–5215	5'-GGG AGC ATA TCA CTT TTC TTG G-3'
917-3.3	5102–5122	5'-GAA CGC CGT CTA CTT ACA AGC-3'
STAPHarb1		5'-GGC CAC GCG TCG ACT AGT CAN NNN NNN NNN GAT AT-3'
STAPHarb2		5'-GGC CAC GCG TCG ACT AGT CAN NNN NNN NNN GAT CA-3'
arb3		5'-GGC CAC GCG TCG ACT AGT CA-3'

^a Localization in the published nucleotide sequence on Tn917 (accession number M11180).

TABLE 3. Functionality of arbitrary PCR in different Tn917 mutants

Mutant	Functionality ^a of PCR with first arbitrary primer:							
	STAPHarb1 and first/second Tn917 primers:				STAPHarb2 and first/second Tn917 primers:			
	917-5.3/917-5.2	917-5.3/917-5.1	917-5.2/917-5.1	917-3.3/917-3.2	917-5.3/917-5.2	917-5.3/917-5.1	917-5.2/917-5.1	917-3.3/917-3.2
M10	+	+	+	+	+	+	+	+
M11	–	–	–	+	–	+	–	+
M12	+	+	+	–	–	–	–	–
M15	+	+	+	+	–	–	+	+
M16	+	–	+	+	+	+	+	–
M20	–	–	+	+	+	+	+	–
M21	–	–	–	+	–	+	+	+
M22	+	–	–	+	+	+	–	+
M24	–	+	+	+	+	+	+	+

^a +, Arbitrary PCR fragments containing Tn917 flanking chromosomal DNA.

ing the chromosomal DNA of the amplified fragments was performed in the unfinished genome database for *S. epidermidis* RP62A at The Institute of Genomic Research (TIGR) homepage (<http://www.tigr.org>) and by using a standard nucleotide-nucleotide BLAST (blastn) search (1) in the recently annotated *S. epidermidis* ATCC 12228 genome (accession number NC_004461).

RESULTS

Evaluation of the arbitrary PCR. To establish a site-specific arbitrary PCR for the characterization of Tn917 insertions in *S. epidermidis*, we generated six Tn917-specific primers homologous to the 5'- and 3'-end of transposon Tn917. Two additional arbitrary primers, including a conserved 5' end of 20 bases (37), 10 random bases, and a specific 3' end of five bases adapted to the GC content of about 30% in staphylococci, were generated (Table 2). To evaluate the functionality of the arbitrary PCR the Tn917 insertion sites of six well-characterized biofilm-negative mutants (Table 1) of the biofilm-positive wild-type strain *S. epidermidis* 1457 were characterized with the arbitrary PCR. In the first round of PCR, a smear of amplified fragments could be detected for all of the six investigated mutants and primer combinations by gel electrophoresis (data not shown). To exclude nonspecific pairing of the Tn917 primers due to the low annealing temperatures in the first round of PCR, we used in the second round the respective distally located Tn917-specific primers. Despite the positive reaction for all primer pairs in the first round in the second, specific PCR round fragments containing Tn917 flanking chromosomal DNA could not be amplified with all primer pairs (Table 3).

However, with the respective Tn917-specific distal primers 917-5.1, 917-5.2, and 917-3.2 fragments containing Tn917, flanking chromosomal DNA could be amplified by the arbitrary PCR for all of the six investigated mutants (Table 3), whereas with primer 917-3.1 no sufficiently visible fragments could be obtained by arbitrary PCR. However, in a standard PCR assay this primer was fully functional (data not shown).

The Tn917 insertion sites were identified from the resulting sequences of arbitrary PCR fragments by analysis with the BLAST tool of the unfinished *S. epidermidis* RP62A genome at TIGR's homepage (www.tigr.org). All Tn917 insertions of the yet-characterized mutants could be identified correctly. Thus, the typical 5-bp duplication of Tn917 insertions could be identified in mutants M21, M22, and M24 (Table 4), whereas only the 5' end of Tn917 flanking regions could be cloned for the characterization of these mutants (30). All arbitrary PCRs were performed twice, and similar results were obtained in both analyses (data not shown).

Identification of the Tn917 insertion in mutant M12. The mutant M12 is characterized by a nonmucoid, biofilm- and hemagglutination-negative phenotype compared to its wild-type *S. epidermidis* 1457 (30). By using the arbitrary PCR technique, a 5' Tn917 flanking chromosomal DNA fragment of the Tn917 insertion site could be amplified, whereas for the 3' site no fragment containing flanking chromosomal DNA was obtained. However, the 5' Tn917 flanking fragment was sufficient for the identification of the Tn917 insertion site by a

TABLE 4. Nucleotide sequences flanking Tn917 insertions of *S. epidermidis* transposon mutants

Mutant	Gene inactivated	Tn917 orientation ^a	Insertion site sequence (position) ^b
M10	<i>icaA</i>	+	TAT GTT ATT (929) Tn917 TTA TTT ATC
M11	<i>icaA</i>	+	TTT ATT AAA (86) Tn917 TTA AAG AAA
M21	<i>icaC</i>	+	TTT ATA TAT (1001) Tn917 TAT ATT CAG
M22	<i>icaC</i>	+	TTT ATT TTC (154) Tn917 TTT TCG GCA
M24	<i>icaA</i>	–	GTC AAT AGA (331) Tn917 ATA GAG GTA
M15	<i>rsbU</i>	+	TCA CAA AAA (19) Tn917 AAA AAG TTA
M12	ND ^c	–	TTT AAA GTG (792) Tn917 AAG TGG AAC
M16	ND	–	GGT TGG TAT (110) Tn917 GGT ATG ACA
M20	ND	–	CGG TTT AAA (587) Tn917 TTA AAT GCA

^a Orientation of the *erm* gene of Tn917 with respect to the direction of the inactivated open reading frame.

^b The Tn917 flanking chromosomal region is displayed in the direction of the inactivated open reading frame. The typical 5-bp duplications of the Tn917 insertion are indicated in boldface. The insertion sites relative to the start point of the respective open reading frame are indicated in parentheses.

^c ND, not determined.

Consensus sequence	DGWK TND N ₁₂₋₁₅	GGRWAW
<i>Staphylococcus epidermidis</i>	AGTT TAA N ₁₃	GGGTAT
<i>Staphylococcus aureus</i>	AGTT TAA N ₁₃	GGGTAT
<i>Bacillus subtilis</i> (<i>purR</i>)	AGTT TAA N ₁₂	GGGAAT
<i>Bacillus subtilis</i> (<i>ctc</i>)	GGTT TAA N ₁₄	GGGTAT

FIG. 1. Alignment of the conserved σ^B promoter motif within the *purR* genes of *S. epidermidis*, *S. aureus*, and *B. subtilis* with the consensus sequence (38) and the *ctc* promoter of *B. subtilis*, a well-characterized σ^B -dependent promoter (38).

BLAST search in *S. epidermidis* RP62A, as well as in the recently annotated genome of the *icaADBC*-negative *S. epidermidis* ATCC 12228. The correct identification of the Tn917 insertion was confirmed by PCR with newly generated primers specific for the identified Tn917 flanking chromosomal regions (data not shown) and the typical 5-bp duplication of Tn917 insertions could thereby be characterized (Table 4). The insertion site of Tn917 in this mutant is located at position 792 of an 825-bp open reading frame (*orf2*) with high homology to the

purR gene of *Bacillus subtilis*, *Lactococcus lactis*, and *S. aureus*. Thus, the highest homology of the *S. epidermidis purR* homologue was observed with *S. aureus* with respect of the nucleotide and the putative amino acid sequence (data not shown). Interestingly, the Tn917 insertion site in *purR* is preceded by a region (bp 662 to 688 of *purR*) homologous to the consensus sequence of σ^B promoters in *B. subtilis* (38). This DNA motif is highly conserved between the *purR* genes of *S. epidermidis*, *S. aureus*, and *B. subtilis* (Fig. 1). In the published sequences of *S. aureus*, homologous genes could be identified in all annotated genomes (Fig. 2).

Consensus search for PurBox sequences. The PurR regulator homologues in *B. subtilis* and *L. lactis* regulate transcription of depending genes by binding to specific DNA regions, referred to as PurBoxes (17, 52), which are located in a palindromic way pairwise upstream of regulated genes in *B. subtilis* (47). Using the consensus sequence for PurBox sequences in *B. subtilis* and *L. lactis*, AWWWCCGAACWWT (17), no potential PurR regulatory sequences could be identified in the *icaADBC* operon by a consensus search with a tolerance of up to two mismatches in the low-importance weight bases or one mismatch of the high-importance weight bases of the consensus sequence (47). In addition, the negative regulator *icaR* (6,




FIG. 2. Chromosomal map of the *S. epidermidis glcA*, *glcT*, *icaADBC*, *purR*, and *rsbU* genes and their respective homologues in *S. aureus*. The origin of replication is marked by the gene *dnaA*. Except for *purR*, the genes involved in biofilm formation of *S. epidermidis* are rearranged between the *icaADBC*-positive and -negative strains, whereas in *S. aureus* the location of these genes are conserved in four independent strains.




FIG. 3. Phenotypes of different *S. epidermidis* mutants. (A) Biofilm formation on polystyrene (left) and glass (right) surfaces. Biofilm formation of the mutants M16 and M20 is comparable to that of wild-type *S. epidermidis* 1457, whereas the control mutant M10 displays a biofilm-negative phenotype. (B) Phenotypes on Congo red agar (CRA_{BHI}). Mutants M16 and M20, as well as the wild-type *S. epidermidis* 1457, displayed a dry crystalline morphology of black colonies, whereas the red colonies of mutant M10 display a smooth surface. Similar results were obtained on CRA_{TSB} (data not shown). (C) Phenotypes on Purple_{GlcNAc}. The wild-type *S. epidermidis* 1457 displays a mucoid phenotype, whereas the biofilm-positive Tn917 mutants M16 and M20, as well as the biofilm-negative mutant M10, displayed a mucoid-negative phenotype on this agar.

7) of the *icaADBC* operon is not preceded by a PurR regulatory sequence.

Generation and characterization of nonmucoid Tn917 mutants. Biofilm-negative mutants of the strongly biofilm-positive and mucoid *S. epidermidis* 1457 displayed a nonmucoid phenotype on purple agar supplemented with 0.5% *N*-acetylglucosamine (Purple_{GlcNAc}). However, the connection between biofilm formation and a mucoid phenotype is unclear at present. Therefore, we carried out transposon mutagenesis, and nonmucoid mutants were further characterized with respect to their ability to establish biofilms. Two mutants M16 and M20 were identified displaying a nonmucoid phenotype on Purple_{GlcNAc} compared to the wild-type *S. epidermidis* 1457 (Fig. 3B), whereas phenotypes on Congo red agar (Fig. 3C) and biofilm formation (Fig. 3A) were comparable between the wild type and mutants. The genetic linkage of phenotype and Tn917 insertion of these isogenic mutants was demonstrated by phage transduction of the Tn917 insertions into *S. epidermidis* 1457. The resulting transductants displayed a phenotype identical to that of the primary mutants (data not shown). The Tn917 insertion sites were subsequently characterized by the arbitrary PCR method, and 5' and 3' Tn917 flanking chromosomal DNA fragments could be amplified for M16 and M20 and the typical 5-bp duplication of Tn917 insertions could be identified by a BLAST search in *S. epidermidis* RP62A, as well as in *S. epidermidis* ATCC 12228 (Table 4). The correct identification of the Tn917 insertion was confirmed by PCR by using newly generated primers specific for the identified Tn917 flanking chromosomal regions (data not shown). In both mu-

tants the Tn917 insertion was located in open reading frames homologous to genes related to the phosphoenolpyruvate-sugar phosphotransferase system (PTS) of *B. subtilis*, *S. carnosus*, and *S. aureus*.

The Tn917 insertion of mutant M16 was localized at position 110 of an 846-bp open reading frame homologous to the gene of the rho-independent antiterminator GlcT of the *B. subtilis* and *S. carnosus* PTS (18, 50) and a respective homologue of *S. aureus* N315 (accession no. AP003133). Homologous genes could also be identified in all annotated *S. aureus* genomes (Fig. 2). Thus, the highest homology of the *S. epidermidis* *glcT* homologue was observed compared to the respective gene in *S. aureus* (Fig. 2). However, the homology to the respective *glcT* gene and the GlcT protein of *S. carnosus* is similar, and all staphylococcal homologues displayed only a moderate homology to the respective *B. subtilis* genes and proteins (data not shown).

The Tn917 insertion of mutant M20 was localized at position 587 of a 2,025-bp open reading frame homologous to EII proteins of the PTS in *B. subtilis* and *S. carnosus* (4, 5, 54) and a respective homologue of *S. aureus* N315 (accession no. AP003129). Homologous genes could also be identified in all annotated *S. aureus* genomes (data not shown). The homology of the respective putative gene products of *S. epidermidis* and *S. aureus* to the EII protein GlcA of *S. carnosus*, which is a glucose-specific component of the PTS, is strikingly high (data not shown).

DISCUSSION

Several genomes of *S. aureus* are published, and two complete genome sequences of *S. epidermidis* are available. The genome of the nonpathogenic *S. epidermidis* ATCC 12228 was annotated recently, and the genome of the pathogenic *S. epidermidis* isolate RP62A is available from TIGR. In addition, the complete genomes of *S. carnosus* and *S. haemolyticus* will be available in the near future. Therefore, in the postgenomic era, small chromosomal fragments are sufficient to identify regions or genes of interest in the published staphylococcal genomes. The *E. faecalis* transposon Tn917 is a well-characterized tool in genetic manipulation of gram-positive bacteria for identifying genes related to characteristic phenotypes or metabolic pathways. In the present study, we established a site-specific arbitrary PCR for the rapid and easy identification of Tn917 insertion sites in *S. epidermidis*. As a control of functionality of this method, we used six independent and well-characterized biofilm-negative Tn917 mutants of *S. epidermidis* 1457. For all six of these mutants, short 5' and 3' flanking chromosomal DNA fragments could be amplified by the arbitrary PCR. A set of different primer combinations were necessary to amplify fragments sufficient for a correct identification in all mutants. By using the BLAST tool to search TIGR's database of the unfinished *S. epidermidis* genome, all insertion sites could be correctly identified. For all mutants, the typical 5-bp duplication at the Tn917 insertion site could be detected, completing the data for mutants for which only the 5' Tn917 flanking chromosomal region could be identified by cloning (30). These results were reproducible and confirm the functionality of this arbitrary PCR method for identifying

Tn917 insertion sites in the genome of the used *S. epidermidis* 1457.

Comparison of the three annotated *S. aureus* genomes revealed that a large number of genes are only present in one of the three strains (2, 23). Therefore, it is possible that additional *S. epidermidis*-specific genes are missing in the RP62A and ATCC 12228 genomes, and the genome of additional clinical *S. epidermidis* isolates should be characterized to complete our knowledge of *S. epidermidis* variability and to optimize our understanding of biofilm formation and the pathogenesis of foreign body-related infections.

Using this arbitrary PCR technique, we identified the Tn917 insertion site of the biofilm-negative mutant M12 of wild-type strain *S. epidermidis* 1457. M12 harbors the Tn917 insertion in a homologue to the *purR* genes of *S. aureus*, *B. subtilis*, and *L. lactis*. In *B. subtilis* *purR* encodes a negative regulator of purine synthesis (52, 53), whereas in *L. lactis* the regulator PurR has a positive regulatory effect on purine synthesis (17). In *B. subtilis* a variety of other metabolic genes and operons are regulated by PurR (47). The direct regulation of the *icaADBC* operon by PurR is unlikely due to a missing putative PurR-binding site. In addition, the gene coding for the negative regulator IcaR (6, 7) of the *icaADBC* operon is not preceded by a PurR-binding site. Therefore, it could be speculated that purine synthesis plays a crucial role in biofilm formation. However, the Tn917 insertion site is preceded by a putative σ^B dependent promoter, which could explain the biofilm-negative phenotype by inactivation of genes downstream of *purR*. Interestingly, in *B. subtilis* the σ^B promoter within the *purR* gene seems to be silent (38). In *S. epidermidis*, regulation of biofilm expression by σ^B was predicted (19), suggesting that these genes could play an additional regulatory role or could act as the mediator of the σ^B regulation. Further experiments to elucidate the effect of these putative regulatory genes on the regulation of biofilm expression in *S. epidermidis* are under way.

The strongly biofilm-positive wild-type *S. epidermidis* 1457 displays a mucoid phenotype on Purple_{GlcNAc} and all as-yet-characterized biofilm-negative Tn917 mutants displayed a non-mucoid phenotype under these conditions. A mucoid phenotype is often associated with exopolysaccharide formation (3, 24) and is associated with pathogenicity of a variety of bacteria (3, 34, 48). However, the linkage between biofilm expression and a mucoid phenotype in *S. epidermidis* is unclear. To elucidate this linkage, we carried out a Tn917 transposon mutagenesis with a subsequent screening for nonmucoid mutants. Two mutants M16 and M20 with a nonmucoid but still biofilm-positive phenotype could be isolated, and the genetic linkage to the observed phenotypes could be demonstrated by phage transduction. The Tn917 insertions of M16 were identified in a homologue to the *glcT* gene coding for a rho-independent antiterminator in *S. carnosus* and *B. subtilis*. Homologous genes were also observed in *S. aureus*. The Tn917 insertions of M20 was identified in a homologue to the *glcA* and *glcB* genes of *S. carnosus* and the *ptsG* gene of *B. subtilis*, which represents the glucose-specific EII proteins in the PTS of these species. Homologous genes were also observed in *S. aureus*; however, the sugar specificity of the *S. aureus* homologue is not yet characterized. The homology of the inactivated genes in M16 and M20 to these components of the PTS of *S. carnosus* and *B.*

subtilis suggests that sugar transport via the PTS is necessary for a mucoid phenotype. Recently, it could be demonstrated that, besides *icaADBC*, a glucose-dependent factor is required for PIA synthesis and biofilm formation (10). However, sugar transport via this putative EII protein of the PTS is not essential for biofilm formation of *S. epidermidis*. Transcriptional analysis of *glcT* and *glcA* and investigations of the sugar specificity of the EII protein are under way to elucidate the mechanisms of the influence of the *S. epidermidis* PTS on the mucoid phenotype in *S. epidermidis*.

All of the newly identified genes involved in biofilm formation or expression of a mucoid phenotype of *S. epidermidis* are present in all published staphylococcal genomes (Fig. 2), indicating an important role for these genes in the cell biology of staphylococci. However, in *S. aureus* the localization of the homologous genes within the chromosome is conserved, whereas in *S. epidermidis* the genes belonging to the PTS system are localized in completely different chromosomal areas (Fig. 2), indicating major chromosomal rearrangements between *icaADBC*-positive and -negative *S. epidermidis* isolates.

The characterization of the new mutants gives evidence that the arbitrary PCR is a suitable tool for the rapid identification of Tn917 insertions in *S. epidermidis*. The primers for the established arbitrary PCR were adapted to the GC content of staphylococci. Therefore, we expect that this method is a useful tool for identifying Tn917 insertion sites of interest in all staphylococcal species.

ACKNOWLEDGMENTS

We thank Rainer Laufs for continuous support.

This work was supported by Deutsche Forschungsgemeinschaft grants given to D.M., J.K.-M.K., and H.R.

REFERENCES

- Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment search tool. *J. Mol. Biol.* **215**:403–410.
- Baba, T., F. Takeuchi, M. Kuroda, H. Yuzawa, K. Aoki, A. Oguchi, Y. Nagai, N. Iwama, K. Asano, T. Naimi, H. Kuroda, L. Cui, K. Yamamoto, and K. Hiramatsu. 2002. Genome and virulence determinants of high virulence community-acquired MRSA. *Lancet* **359**:1819–1827.
- Boucher, J. C., H. Yu, M. H. Mudd, and V. Deretic. 1997. Mucoid *Pseudomonas aeruginosa* in cystic fibrosis: characterization of muc mutations in clinical isolates and analysis of clearance in a mouse model of respiratory infection. *Infect. Immun.* **65**:3838–3846.
- Christiansen, I., and W. Hengstenberg. 1996. Cloning and sequencing of two genes from *Staphylococcus carnosus* coding for glucose-specific PTS and their expression in *Escherichia coli* K-12. *Mol. Gen. Genet.* **250**:375–379.
- Christiansen, I., and W. Hengstenberg. 1999. Staphylococcal phosphoenolpyruvate-dependent phosphotransferase system—two highly similar glucose permeases in *Staphylococcus carnosus* with different glucoside specificity: protein engineering in vivo? *Microbiology* **145**:2881–2889.
- Conlon, K. M., H. Humphreys, and J. P. O'Gara. 2002. *icaR* encodes a transcriptional repressor involved in environmental regulation of *ica* operon expression and biofilm formation in *Staphylococcus epidermidis*. *J. Bacteriol.* **184**:4400–4408.
- Conlon, K. M., H. Humphreys, and J. P. O'Gara. 2002. Regulation of *icaR* gene expression in *Staphylococcus epidermidis*. *FEMS Microbiol. Lett.* **216**:171–177.
- Cramton, S. E., C. Gerke, N. F. Schnell, W. W. Nichols, and F. Götz. 1999. The intercellular adhesion (*ica*) locus is present in *Staphylococcus aureus* and is required for biofilm formation. *Infect. Immun.* **67**:5427–5433.
- Cramton, S. E., M. Ulrich, F. Götz, and G. Döring. 2001. Anaerobic conditions induce expression of polysaccharide intercellular adhesin in *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Infect. Immun.* **69**:4079–4085.
- Dobinsky, S., K. Kiel, H. Rohde, K. Bartscht, J. K. M. Knobloch, M. A. Horstkotte, and D. Mack. 2003. Glucose-related dissociation between *icaADBC* transcription and biofilm expression by *Staphylococcus epidermidis*: evidence for an additional factor required for polysaccharide intercellular adhesin synthesis. *J. Bacteriol.* **185**:2879–2886.
- Fitzpatrick, F., H. Humphreys, E. Smyth, C. A. Kennedy, and J. P. O'Gara.

2002. Environmental regulation of biofilm formation in intensive care unit isolates of *Staphylococcus epidermidis*. *J. Hosp. Infect.* **52**:212–218.
12. Frebourg, N. B., S. Lefebvre, S. Baert, and J. F. Lemeland. 2000. PCR-based assay for discrimination between invasive and contaminating *Staphylococcus epidermidis* strains. *J. Clin. Microbiol.* **38**:877–880.
 13. Gerke, C., A. Kraft, R. Sussmuth, O. Schweitzer, and F. Götz. 1998. Characterization of the *N*-acetylglucosaminyltransferase activity involved in the biosynthesis of the *Staphylococcus epidermidis* polysaccharide intercellular adhesin. *J. Biol. Chem.* **273**:18586–18593.
 14. Götz, F. 2002. *Staphylococcus* and biofilms. *Mol. Microbiol.* **43**:1367–1378.
 15. Handwerker, S. 1994. Alterations in peptidoglycan precursors and vancomycin susceptibility in Tn917 insertion mutants of *Enterococcus faecalis* 221. *Antimicrob. Agents Chemother.* **38**:473–475.
 16. Heilmann, C., O. Schweitzer, C. Gerke, N. Vanittanakom, D. Mack, and F. Götz. 1996. Molecular basis of intercellular adhesion in the biofilm-forming *Staphylococcus epidermidis*. *Mol. Microbiol.* **20**:1083–1091.
 17. Kilstrop, M., and J. Martinussen. 1998. A transcriptional activator, homologous to the *Bacillus subtilis* PurR repressor, is required for expression of purine biosynthetic genes in *Lactococcus lactis*. *J. Bacteriol.* **180**:3907–3916.
 18. Knezevic, I., S. Bachem, A. Sickmann, H. E. Meyer, J. Stülke, and W. Hengstenberg. 2000. Regulation of the glucose-specific phosphotransferase system (PTS) of *Staphylococcus carnosus* by the antiterminator protein GlcT. *Microbiology* **146**:2333–2342.
 19. Knobloch, J. K. M., K. Bartscht, A. Sabottke, H. Rohde, H. H. Feucht, and D. Mack. 2001. Biofilm formation by *Staphylococcus epidermidis* depends on functional RsbU, an activator of the *sigB* operon: differential activation mechanisms due to ethanol and salt stress. *J. Bacteriol.* **183**:2624–2633.
 20. Knobloch, J. K. M., M. A. Horstkotte, H. Rohde, P. M. Kaulfers, and D. Mack. 2002. Alcoholic ingredients in skin disinfectants increase biofilm expression of *Staphylococcus epidermidis*. *J. Antimicrob. Chemother.* **49**:683–687.
 21. Knobloch, J. K. M., M. A. Horstkotte, H. Rohde, and D. Mack. 2002. Evaluation of different detection methods of biofilm formation in *Staphylococcus aureus*. *Med. Microbiol. Immunol.* **191**:101–106.
 22. Knobloch, J. K. M., H. von Osten, M. A. Horstkotte, H. Rohde, and D. Mack. 2002. Minimal attachment killing (MAK): a versatile method for susceptibility testing of attached biofilm-positive and -negative *Staphylococcus epidermidis*. *Med. Microbiol. Immunol.* **191**:107–114.
 23. Kuroda, M., T. Ohta, I. Uchiyama, T. Baba, H. Yuzawa, I. Kobayashi, L. Cui, A. Oguchi, K. Aoki, Y. Nagai, J. Lian, T. Ito, M. Kanamori, H. Matsumaru, A. Maruyama, H. Murakami, A. Hosoyama, Y. Mizutani-Ui, N. K. Takahashi, T. Sawano, R. Inoue, C. Kaito, K. Sekimizu, H. Hirakawa, S. Kuhara, S. Goto, J. Yabuzaki, M. Kanehisa, A. Yamashita, K. Oshima, K. Furuya, C. Yoshino, T. Shiba, M. Hattori, N. Ogasawara, H. Hayashi, and K. Hiramoto. 2001. Whole genome sequencing of methicillin-resistant *Staphylococcus aureus*. *Lancet* **357**:1225–1240.
 24. Lee, C. Y., and J. C. Lee. 2000. *Staphylococcus* capsule, p. 361–366. *In* V. A. Fischetti, R. P. Novick, J. J. Ferretti, D. A. Portnoy, and J. I. Rood (ed.), Gram-positive pathogens. ASM Press, Washington, D.C.
 25. Li, Z., D. D. Sledjeski, B. Kreikemeyer, A. Podbielski, and M. D. Boyle. 1999. Identification of *pel*, a *Streptococcus pyogenes* locus that affects both surface and secreted proteins. *J. Bacteriol.* **181**:6019–6027.
 26. Mack, D. 1999. Molecular mechanisms of *Staphylococcus epidermidis* biofilm formation. *J. Hosp. Infect.* **43**(Suppl.):S113–S125.
 27. Mack, D., K. Bartscht, C. Fischer, H. Rohde, C. de Grahl, S. Dobinsky, M. A. Horstkotte, K. Kiel, and J. K. M. Knobloch. 2001. Genetic and biochemical analysis of *Staphylococcus epidermidis* biofilm accumulation. *Methods Enzymol.* **336**:215–239.
 28. Mack, D., W. Fischer, A. Krokotsch, K. Leopold, R. Hartmann, H. Egge, and R. Laufs. 1996. The intercellular adhesin involved in biofilm accumulation of *Staphylococcus epidermidis* is a linear β -1,6-linked glucosaminoglycan: purification and structural analysis. *J. Bacteriol.* **178**:175–183.
 29. Mack, D., J. Riedewald, H. Rohde, T. Magnus, H. H. Feucht, H. A. Elsner, R. Laufs, and M. E. Rupp. 1999. Essential functional role of the polysaccharide intercellular adhesin of *Staphylococcus epidermidis* in hemagglutination. *Infect. Immun.* **67**:1004–1008.
 30. Mack, D., H. Rohde, S. Dobinsky, J. Riedewald, M. Nedelmann, J. K. M. Knobloch, H. A. Elsner, and H. H. Feucht. 2000. Identification of three essential regulatory gene loci governing expression of *Staphylococcus epidermidis* polysaccharide intercellular adhesin and biofilm formation. *Infect. Immun.* **68**:3799–3807.
 31. Mack, D., A. Sabottke, S. Dobinsky, H. Rohde, M. A. Horstkotte, and J. K. M. Knobloch. 2002. Differential expression of methicillin resistance by different biofilm-negative *Staphylococcus epidermidis* transposon mutant classes. *Antimicrob. Agents Chemother.* **46**:178–183.
 32. Mack, D., N. Siemssen, and R. Laufs. 1992. Parallel induction by glucose of adherence and a polysaccharide antigen specific for plastic-adherent *Staphylococcus epidermidis*: evidence for functional relation to intercellular adhesion. *Infect. Immun.* **60**:2048–2057.
 33. McKenney, D., J. Hubner, E. Muller, Y. Wang, D. A. Goldmann, and G. B. Pier. 1998. The *ica* locus of *Staphylococcus epidermidis* encodes production of the capsular polysaccharide/adhesin. *Infect. Immun.* **66**:4711–4720.
 34. Nassif, X., J. M. Fournier, J. Arondel, and P. J. Sansonetti. 1989. Mucoid phenotype of *Klebsiella pneumoniae* is a plasmid-encoded virulence factor. *Infect. Immun.* **57**:546–552.
 35. Nedelmann, M., A. Sabottke, R. Laufs, and D. Mack. 1998. Generalized transduction for genetic linkage analysis and transfer of transposon insertions in different *Staphylococcus epidermidis* strains. *Zentbl. Bakteriol.* **287**:85–92.
 36. Neubauer, H., I. Pantel, and F. Götz. 1999. Molecular characterization of the nitrite-reducing system of *Staphylococcus carnosus*. *J. Bacteriol.* **181**:1481–1488.
 37. O'Toole, G. A., L. A. Pratt, P. I. Watnick, D. K. Newman, V. B. Weaver, and R. Kolter. 1999. Genetic approaches to study of biofilms. *Methods Enzymol.* **310**:91–109.
 38. Petersohn, A., J. Bernhardt, U. Gerth, D. Hoper, T. Koburger, U. Völker, and M. Hecker. 1999. Identification of σ^B -dependent genes in *Bacillus subtilis* using a promoter consensus-directed search and oligonucleotide hybridization. *J. Bacteriol.* **181**:5718–5724.
 39. Rachid, S., S. Cho, K. Ohlsen, J. Hacker, and W. Ziebuhr. 2000. Induction of *Staphylococcus epidermidis* biofilm formation by environmental factors: the possible involvement of the alternative transcription factor sigB. *Adv. Exp. Med. Biol.* **485**:159–166.
 40. Rachid, S., K. Ohlsen, U. Wallner, J. Hacker, M. Hecker, and W. Ziebuhr. 2000. Alternative transcription factor σ^B is involved in regulation of biofilm expression in a *Staphylococcus aureus* mucosal isolate. *J. Bacteriol.* **182**:6824–6826.
 41. Rachid, S., K. Ohlsen, W. Witte, J. Hacker, and W. Ziebuhr. 2000. Effect of subinhibitory antibiotic concentrations on polysaccharide intercellular adhesion expression in biofilm-forming *Staphylococcus epidermidis*. *Antimicrob. Agents Chemother.* **44**:3357–3363.
 42. Rohde, H., J. K. M. Knobloch, M. A. Horstkotte, and D. Mack. 2001. Correlation of biofilm expression types of *Staphylococcus epidermidis* with polysaccharide intercellular adhesin synthesis: evidence for involvement of *icaADBC* genotype-independent factors. *Med. Microbiol. Immunol.* **190**:105–112.
 43. Rupp, M. E., P. D. Fey, C. Heilmann, and F. Götz. 2001. Characterization of the importance of *Staphylococcus epidermidis* autolysin and polysaccharide intercellular adhesin in the pathogenesis of intravascular catheter-associated infection in a rat model. *J. Infect. Dis.* **183**:1038–1042.
 44. Rupp, M. E., J. S. Ulphani, P. D. Fey, K. Bartscht, and D. Mack. 1999. Characterization of the importance of polysaccharide intercellular adhesin/hemagglutinin of *Staphylococcus epidermidis* in the pathogenesis of biomaterial-based infection in a mouse foreign body infection model. *Infect. Immun.* **67**:2627–2632.
 45. Rupp, M. E., J. S. Ulphani, P. D. Fey, and D. Mack. 1999. Characterization of *Staphylococcus epidermidis* polysaccharide intercellular adhesin/hemagglutinin in the pathogenesis of intravascular catheter-associated infection in a rat model. *Infect. Immun.* **67**:2656–2659.
 46. Saklani-Jusforgues, H., E. Fontan, and P. L. Goossens. 2001. Characterisation of a *Listeria monocytogenes* mutant deficient in D-arabitol fermentation. *Res. Microbiol.* **152**:175–177.
 47. Saxild, H. H., K. Brunstedt, K. I. Nielsen, H. Jarmer, and P. Nygaard. 2001. Definition of the *Bacillus subtilis* PurR operator using genetic and bioinformatic tools and expansion of the PurR regulon with *glyA*, *guaC*, *pbuG*, *xpt-pbuX*, *yqhZ-fold*, and *pbuO*. *J. Bacteriol.* **183**:6175–6183.
 48. Schrage, H. M., S. Alberti, C. Cywes, G. J. Dougherty, and M. R. Wessels. 1998. Hyaluronic acid capsule modulates M protein-mediated adherence and acts as a ligand for attachment of group A streptococcus to CD44 on human keratinocytes. *J. Clin. Invest.* **101**:1708–1716.
 49. Sebulsky, M. T., D. Hohnstein, M. D. Hunter, and D. E. Heinrichs. 2000. Identification and characterization of a membrane permease involved in iron-hydroxamate transport in *Staphylococcus aureus*. *J. Bacteriol.* **182**:4394–4400.
 50. Stulke, J., I. Martin-Verstraete, M. Zagorec, M. Rose, A. Klier, and G. Rapoport. 1997. Induction of the *Bacillus subtilis* *ptsGHI* operon by glucose is controlled by a novel antiterminator, GlcT. *Mol. Microbiol.* **25**:65–78.
 51. Vats, N., and S. F. Lee. 2000. Active detachment of *Streptococcus mutans* cells adhered to epon-hydroxylapatite surfaces coated with salivary proteins in vitro. *Arch. Oral Biol.* **45**:305–314.
 52. Weng, M., P. L. Nagy, and H. Zalkin. 1995. Identification of the *Bacillus subtilis* *pur* operon repressor. *Proc. Natl. Acad. Sci. USA* **92**:7455–7459.
 53. Weng, M., and H. Zalkin. 2000. Mutations in the *Bacillus subtilis* purine repressor that perturb PRPP effector function in vitro and in vivo. *Curr. Microbiol.* **41**:56–59.
 54. Zagorec, M., and P. W. Postma. 1992. Cloning and nucleotide sequence of the *ptsG* gene of *Bacillus subtilis*. *Mol. Gen. Genet.* **234**:325–328.
 55. Zagorec, M., and M. Steinmetz. 1991. Construction of a derivative of Tn917 containing an outward-directed promoter and its use in *Bacillus subtilis*. *J. Gen. Microbiol.* **137**(Pt. 1):107–112.