

Release of Extracellular Transformable Plasmid DNA from *Escherichia coli* Cocultivated with Algae

Kazuaki Matsui,^{1*} Nobuyoshi Ishii,² and Zen'ichiro Kawabata¹

Center for Ecological Research, Kyoto University, Otsu 520-2113,¹ and Environmental and Toxicological Science Research Group, National Institute of Radiological Sciences, Inage-ku, Chiba-shi 263-8555,² Japan

Received 12 August 2002/Accepted 13 January 2003

We studied the effects of cocultivation with either *Euglena gracilis* (Euglenophyta), *Microcystis aeruginosa* (Cyanophyta), *Chlamydomonas neglecta* (Chlorophyta), or *Carteria inversa* (Chlorophyta) on the production of extracellular plasmid DNA by *Escherichia coli* LE392(pKZ105). Dot blot hybridization analysis showed a significant release of plasmid DNA by cocultivation with all the algae tested. Further analysis by electrotransformation confirmed the release of transformable plasmid DNA by cocultivation with either *E. gracilis*, *M. aeruginosa*, or *C. inversa*. These results suggest algal involvement in bacterial horizontal gene transfer by stimulating the release of transformable DNA into aquatic environments.

Dissolved DNA (dDNA), DNA that passes through a 0.2- μm -pore-size filter, is a form of extracellular DNA commonly observed in natural aquatic environments and has been found at concentrations from 0.2 to 88 $\mu\text{g liter}^{-1}$ in both freshwater and seawater (7, 16, 27, 28, 32). dDNA was initially thought to be an attractive source of nutrients for microorganisms (15, 31), but finding naturally competent bacterial strains suggested that the occurrence of bacterial dDNA might also act as a source for the uptake of genetic material (9, 14). We recently reported on the stability of the transformability of plasmid DNA in natural lake water and suggested that dDNA may have a genetic function as a gene pool for further natural transformation (22).

DNA from bacteria has been shown to form a major component of the dDNA pool (27, 33), while algal DNA may also be a significant constituent (25). The production of dDNA is controlled by both physicochemical (26) and biological factors (27). Grazing by bacteriovores (11, 17, 33) and cell lysis by viral infection (1, 29, 35) seem to be the main forces releasing bacterial cellular DNA into the surrounding water. However, the presence of bacterial chromosomal DNA in culture liquids (3, 10) and the release of transformable plasmid and chromosomal DNA from cultured bacteria (20) have indicated that active DNA excretion may occur without disruption of the cells.

Photosynthetic algae are major biological constituents of aquatic environments and are also important biological factors in stimulating bacterial growth via their secretion of dissolved organic carbon (5). Although the influence of algae on bacterial growth has been studied, there has been no study of their impact on the production of dDNA by bacteria. Algal blooms are frequently observed in eutrophic waters where antibiotic-resistant coliform bacteria, including *Escherichia coli*, are delivered via wastewater (13) and are probably an important reservoir of antibiotic resistance genes. Thus, one effect of

algae on the production of dDNA by bacteria may be involved in the dissemination of bacterial antibiotic resistance genes via further natural transformation. In this study, we examine the involvement of algae in the production of extracellular plasmid DNA by *E. coli* and the ability of the DNA released to transform in the bacteria.

A 974-bp *PvuII-EcoRI* fragment of pEGFP (Clontech, Palo Alto, Calif.) containing the *lac* promoter and *egfp*, a green fluorescent protein gene, was inserted into the *SmaI-EcoRI* site of pHY300PLK (12), a shuttle vector between *Bacillus subtilis* and *E. coli*. The resulting plasmid, pKZ105, which is expressed as green fluorescence in *E. coli* but not in *B. subtilis*, was transferred into *E. coli* LE392 (30) and used to trace the excretion of cellular plasmid DNA. A slightly modified version of C medium (8) was used as a culture medium with the following composition (in milligrams per liter of distilled water): $\text{Ca}(\text{NO}_3)_2 \cdot \text{H}_2\text{O}$, 150; KNO_3 , 100; $\beta\text{-Na}_2$ glycerophosphate, 50; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 40; vitamin B_{12} , 0.0001; biotin, 0.0001; thiamine HCl, 0.01; and yeast extract, 200. C medium also contained 3 ml of a solution of trace metals (pH adjusted to 7.5). The trace metal solution consisted of the following (in milligrams per liter of distilled water): $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$, 196; $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$, 36; $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 22; $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$, 4; $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 2.5; and $\text{Na}_2\text{EDTA} \cdot 2\text{H}_2\text{O}$, 1,000. Yeast extract was sterilized separately from the other components, and they were mixed together before use. The medium was decanted into a series of autoclaved test tubes (10 ml each) before inoculation of the organisms. *E. coli* LE392(pKZ105) was precultured overnight at 37°C in the modified C medium and directly inoculated into each test tube at 10^5 CFU ml^{-1} . *E. coli* LE392(pKZ105) was also cocultured with either euglenophyte *Euglena gracilis* Z, cyanophyte *Microcystis aeruginosa* (NIES 298), or chlorophyte *Chlamydomonas neglecta* (NIES 439) or chlorophyte *Carteria inversa* (NIES 422) to evaluate the effects of algae on DNA excretion by this bacterium. *E. gracilis* was kindly provided by N. Nakano, Department of Agricultural Chemistry, University of Osaka Prefecture, Sakai, Japan, and the other algal strains were obtained from the National Institute for Environmental Studies (NIES), Environmental Agency, Tsukuba, Japan. All algal

* Corresponding author. Mailing address: Center for Ecological Research, Kyoto University, Kamitanakami Hirano-cho 509-3, Otsu 520-2113, Japan. Phone: 81-077-549-8234. Fax: 81-077-549-8201. E-mail: k-matsui@ecology.kyoto-u.ac.jp.





FIG. 1. (a) Dot blots of plasmid from experimental cultures. Each dot contains DNA (equivalent to the amount of DNA in 1 ml of culture). No signal was detected from the samples of monospecies cultures. Pure isolated plasmid DNA was used as a standard (Std.), and the amount of DNA applied on each dot is shown to the right of the gel. (b) Regressions of the intensity of hybridization signals (y axis) against the amount of pKZ105 DNA in each dot (x axis). The correlation coefficient for the linear regression was 0.996 over the range of 10 to 1,000 pg of standard pKZ105 DNA.

strains were axenic and had been precultured for 10 days in the modified C medium. Precultured algae were concentrated by centrifugation ($3,000 \times g$, 10 min) and then directly inoculated into each test tube at 4.5×10^4 organisms ml^{-1} (*E. gracilis*), 1.8×10^6 organisms ml^{-1} (*M. aeruginosa*), 4.2×10^5 organisms

ml^{-1} (*C. neglecta*), and 7.7×10^3 organisms ml^{-1} (*C. inversa*). All the test tubes were incubated for 17 days at 25°C with a light regimen of 12 h of light and 12 h of dark under cool-white fluorescent lamps.

The population of each organism was assessed by counting




FIG. 2. Changes in the concentration of total extracellular DNA (shaded bars) and pKZ105 aliquot (black bars) and changes in the number of each organism (lines) in cultures of *E. coli* LE392(pKZ105) (○) grown with *E. gracilis* (▲) (a), *C. neglecta* (◆) (b), *C. inversa* (▼) (c), or *M. aeruginosa* (■) (d) and in *E. coli* LE392(pKZ105) monoculture (e). The means \pm standard deviations (error bars) from three experiments are shown. Statistically significant release of pKZ105 (at 5% level) is denoted by an asterisk.

E. coli CFU on Luria-Bertani (LB) plates (30) and by directly counting algae under a microscope. At each observation time, extracellular DNA in the culture fluids was separated from the microorganisms by brief centrifugation ($3,000 \times g$, 5 min), followed by filtration through a 0.22- μm -pore-size Millex-GV25 filter unit (Millipore, Bedford, Mass.). The filtrate was concentrated using 30,000-molecular-weight-cutoff centrifugal filter devices (CENTRIPLUS; Millipore) and then purified by utilizing QIAquick PCR purification kit (QIAGEN K. K., Tokyo, Japan). The concentration of total extracellular DNA was determined by a modified version of the fluorescence method of DeFlaun et al. (7), using λ -HindIII digest (Takara Shuzo, Tokyo, Japan) as standard DNA. To increase the sensitivity of the measurement, we used SYBR green I (Molecular Probes, Eugene, Oreg.) instead of Hoechst 33258 dye. pKZ105 was quantified by dot blot hybridization, using part of the *egfp* fragment as a probe, by the method of Matsui et al. (22). Significant differences in the release of plasmid DNA were examined statistically by one-way analysis of variance with post hoc Fisher's protected least significant difference test, using Stat View 5.0 for Macintosh (SAS Institute Inc. Cary, N.C.).

An aliquot of purified DNA, equivalent to the DNA in 0.25 ml of culture, was also used to electrotransform *E. coli* JM109 by the method of Sambrook and Russell (30). A series of transformations with different amounts of pure, isolated pKZ105 from *E. coli* LE392(pKZ105) was also conducted to obtain a standard curve of transformation. Isolation of pKZ105 was conducted by alkaline lysis (30) using the CONCERT rapid plasmid purification system (Life Technologies, Inc.,

Rockville, Md.). Fish sperm DNA (molecular biology grade; Roche Diagnostics, Tokyo, Japan) was mixed with pKZ105 as a mimic of total extracellular DNA to test the interference effects on transformation. Transformants were selected for resistance to 20 μg of tetracycline per liter in LB agar medium. Expression of the green fluorescence gene was confirmed by color change of colonies and expression of green fluorescence.

A dot blot and the standard curve of dot blot hybridization are presented in Fig. 1. The DNA detection limits of the methods employed were 10 $\mu\text{g ml}^{-1}$ (dot blot hybridization analysis; plasmid pKZ105) and 200 $\mu\text{g ml}^{-1}$ (fluorometric analysis; total extracellular DNA). The time courses of population density and the release of extracellular DNA were examined in monocultures and cocultures of *E. coli* LE392(pKZ105) (Fig. 2). Significant extracellular DNA production was observed from cocultures of *E. coli* LE392(pKZ105) with algae, while the amount of DNA detected and the timing of DNA production varied with the algal species.

When *E. coli* was cocultured with *E. gracilis*, release of total extracellular DNA was observed after 7 days of cultivation (Fig. 2a), and all the total DNA detected must have originated from *E. coli* LE392(pKZ105), since no DNA was detected in the monoculture of *E. gracilis* (Fig. 3a). The data from dot blot hybridization demonstrated that the pKZ105 plasmid was also present in the extracellular DNA fraction and confirmed the release of DNA from *E. coli* LE392(pKZ105) when it was cocultivated with *E. gracilis*. Extracellular DNA production was not observed in the monoculture of *E. coli* LE392(pKZ105), and the population changes of *E. coli* LE392(pKZ105) were no




FIG. 3. Changes in the concentration of total extracellular DNA (shaded bars) and in the number of organisms (lines) in monocultures of *E. gracilis* (▲) (a), *C. neglecta* (◆) (b), *C. inversa* (▼) (c), and *M. aeruginosa* (■) (d). Note that there was no *E. coli*(pKZ105) in these cultures, so no pKZ105 was recorded. The means \pm standard deviations (error bars) from three experiments are shown.

different when *E. gracilis* was included in the culture (Fig. 2a and e). Thus, the production of extracellular DNA would not have been caused by cell lysis of *E. coli*, and indirect effects, such as the involvement of metabolite-mediated effects of *E. gracilis*, were indicated.

Coculturing *E. coli* LE392(pKZ105) with *C. neglecta* or *C. inversa* also resulted in a gradual increase of total extracellular DNA during cultivation (Fig. 2b and c), but the detection of similar total amounts of extracellular DNA in the monocultures of *C. neglecta* and *C. inversa* suggested that in this case the majority of the extracellular DNA in the cocultures was released from the algae (Fig. 3b and c). However, detection of pKZ105 DNA by dot blot hybridization indicated that the plasmid DNA was also released from *E. coli* LE392(pKZ105) when it was cultivated with *C. neglecta* or *C. inversa*. The sudden decline in the cell density of *E. coli* after 7 days of culture suggested the release of *E. coli* DNA by cell lysis (Fig. 2b and c). It is possible that *C. neglecta* or *C. inversa* produced antibacterial substances, since some algal species are known to produce substances which inhibit the growth and respiration of bacteria (4). However, the detection of *E. coli* plasmid DNA after 3 days of culture, when the *E. coli* population density was at its highest, suggested that there was an alternative effect of these cocultured algae on the release of extracellular DNA. In either case, extracellular DNA was released from *E. coli* LE392(pKZ105) by cocultivation with *C. neglecta* or *C. inversa*.

When *E. coli* LE392(pKZ105) was cocultured with *M. aeruginosa*, extracellular DNA was detected only after the first 3 days of culture (Fig. 2d). The increase in the population

density of *E. coli* during the first 3 days and the fact that no extracellular DNA was detected after 7 days of cultivation (when the *E. coli* cell density had declined) suggested that cocultivation with *M. aeruginosa* had a stimulative effect on the release of *E. coli* DNA. In the *M. aeruginosa* monoculture, extracellular DNA was detected at 7, 10, and 17 days of culture but not at 3 days (Fig. 3d). Although the cellular condition of the *M. aeruginosa* cocultured with *E. coli* LE392(pKZ105) seemed different from the *M. aeruginosa* in monoculture, the detection of pKZ105 DNA in the coculture did indicate that there had been some release of extracellular DNA from *E. coli* LE392(pKZ105) when it was cocultured with *M. aeruginosa*.

No extracellular DNA was detected in monocultures of *E. coli* LE392(pKZ105), and the culture conditions employed here were clearly not suitable for *E. coli* LE392(pKZ105) to release extracellular DNA, at least for the 17 days of cultivation (Fig. 2e). Thus, all of the four algal species tested were involved in the production of extracellular DNA from *E. coli* under our experimental conditions.

Recent advances in studies of molecular microbial ecology and genome analysis indicate that horizontal gene transfer is involved in bacterial evolution and adaptation to changing environments (6, 24). As a means of gene transfer, transformation may be the most important mechanism by which bacteria acquire DNA from distantly related organisms, including eukaryotes (23). We think that there may be natural transformation-mediated horizontal gene transfer in an aquatic ecosystem from a study of a naturally transformable bacterium, *B. subtilis* (K. Matsui, N. Ishii, and Z. Kawabata, submitted for

TABLE 1. Electrotransformation of *E. coli* JM109 with concentrated extracellular DNA from 0.25 ml of cocultured *E. coli* LE392(pKZ105)

Culture condition of <i>E. coli</i> LE392(pKZ105)	Cultivation period (no. of days)	No. of transformants (CFU) (mean \pm SD)	
		Expected ^a	Detected
Cocultured with <i>E. gracilis</i>	3	0	0
	7	$(5.7 \pm 2.7) \times 10^1$	$(1.0 \pm 0.5) \times 10^0$
	10	$(2.7 \pm 0.5) \times 10^2$	$(4.0 \pm 2.0) \times 10^0$
	17	$(1.2 \pm 0.2) \times 10^3$	0
Cocultured with <i>C. neglecta</i>	3	$(2.1 \pm 0.2) \times 10^2$	0
	7	$(1.9 \pm 0.9) \times 10^2$	0
	10	$(1.0 \pm 0.5) \times 10^2$	0
	17	$(1.2 \pm 0.3) \times 10^2$	0
Cocultured with <i>C. inversa</i>	3	$(2.6 \pm 0.2) \times 10^1$	$(2.8 \pm 1.3) \times 10^1$
	7	$(2.7 \pm 0.8) \times 10^1$	0
	10	$(1.6 \pm 0.1) \times 10^1$	0
	17	$(1.8 \pm 0.1) \times 10^1$	0
Cocultured with <i>M. aeruginosa</i>	3	$(2.7 \pm 1.7) \times 10^1$	$(1.0 \pm 0.5) \times 10^0$
	7	0	0
	10	0	0
	17	0	0

^a Expected numbers of transformants were calculated from the equation of linear regression of the standard curve, using the amount of pKZ105 detected from each coculture.

publication). The availability of a transformable gene is a necessity for successful horizontal gene transfer by transformation and for completion of the transformation event (21).

To determine whether the detected plasmid (pKZ105) maintained its transformability, we conducted further transformation experiments (Table 1). A standard curve was obtained by a series of transformations with pKZ105 which were isolated from *E. coli* LE392(pKZ105) (Fig. 4). By applying the amount of pKZ105 detected to the standard curve, the number of transformants on each electrotransformation was also estimated. No transformants were obtained with the DNA from cocultivation with *C. neglecta*, while significant amounts of extracellular pKZ105 DNA were detected. Since the total extracellular DNA concentration was high in coculture with *C. neglecta*, that level of total extracellular DNA might suppress the transformation event. This hypothesis was tested by conducting the transformation experiments using pure pKZ105 DNA mixed with fish sperm DNA (Fig. 4). The addition of fish sperm DNA slightly enhanced, and did not reduce, the transformation efficiency. Thus, the total extracellular DNA would not have interfered with the transformation, and failure to transform with the DNA from cocultivation with *C. neglecta* would result in the release of no transformable DNA by cocultivation with this strain. However, in contrast to the result from cocultivation with *C. neglecta*, the successful electrotransformation did indicate the release of transformable DNA from *E. coli* LE392(pKZ105) when in cocultivation with *E. gracilis*, *C. inversa*, or *M. aeruginosa*.

Previous studies have also suggested active excretion of DNA from several bacteria, but the mechanism by which those bacteria excrete cellular DNA is not yet clear (3, 10, 20, 26). Lorenz et al. (20) reported the active release of transformable plasmids and chromosomal DNA from competent *B. subtilis* during growth in minimal media. Their observation suggested

that DNA excretion was influenced by the physiological state of the bacterial cells. In aquatic environments, both biotic and abiotic factors are involved in determining the physiological state of bacteria. A recent study has demonstrated that extracellular products from a green alga, *Chlamydomonas reinhardtii*, suppress heterocyst formation in the cyanobacterium *Anabaena flos-aquae* (18) and suggested that there are chemical interactions which can alter the physiological state of other organisms and even organisms in other kingdoms. Paul and David (26) showed that the production of extracellular DNA from *E. coli* HB101(R388) was altered by physicochemical factors such as salinity changes. These studies suggested that the existence of chemical interactions which alter the physiological state of *E. coli* LE392(pKZ105) and that the subsequent release of transformable DNA could have occurred by cocultivation with algae.

Compounds, such as acylated homoserine lactones and oligopeptides, are known to be involved in cell-to-cell communication and the alternation of both intra- and interspecific behaviors in bacteria (2, 34). As one example of interspecies interaction, compounds produced from the macroalga *Delisea pulchra* alter the behavior of bacteria by interfering with the bacterial signaling system mediated by acylated homoserine lactones (19, 34). Thus, there may be a signal compound which stimulates DNA excretion from *E. coli*. Characterization of this compound(s) would clarify this hypothesized-chemical-mediated interaction between algae and bacteria.

The reason why algae stimulate the release of bacterial DNA is another question to which there is no clear answer. One possibility is that stimulation of the release of bacterial DNA, which is rich in phosphorus, may be an alternative way for algae to ameliorate the competition with bacteria for nutrients,




FIG. 4. Relationship between the number of transformants and the amount of applied pKZ105 DNA mixed with fish sperm DNA. pKZ105 DNA was mixed with the following amounts of fish sperm DNA: 0 (○), 0.1 (△), 1 (▽), 10 (◇), and 100 (■) ng. The solid line indicates the regression; the correlation coefficient was 0.997. The means \pm standard deviations (error bars) from three experiments are shown.

since uptake of phosphate is commonly dominated by bacteria (>50 to nearly 100%) in many freshwater environments (36).

The exact mechanisms by which, and the reasons why, the algae tested could induce the excretion of *E. coli* plasmid DNA still remain to be clarified, but this study has clearly demonstrated algal involvement in the production of transformable plasmid DNA by *E. coli* bacteria, which may suggest a new ecological interaction facilitating horizontal gene transfer in aquatic environments.

This work was supported in part by a Japanese Ministry of Education, Culture, Sports, Science and Technology Grant-in-Aid for Creative Basic Research (09NP1501) and a Grant-in-Aid for Basic Research (A) (13309009). K.M. was also supported by a Research Fellowship of the Japan Society for the Promotion of Science for Young Scientists.

We thank Masaya Ueki for cultivation of algae, Atsushi Maruyama and Yasushi Miyamoto for statistical analysis, and Mary Morris for improving the English.

REFERENCES

- Alonso, M. C., V. Rodriguez, J. Rodriguez, and J. J. Borrego. 2000. Role of ciliates, flagellates and bacteriophages on the mortality of marine bacteria and on dissolved-DNA concentration in laboratory experimental systems. *J. Exp. Mar. Biol. Ecol.* **244**:239–252.
- Bassler, B. L. 2002. Small talk: cell-to-cell communication in bacteria. *Cell* **109**:421–424.
- Catlin, B. W. 1956. Extracellular deoxyribonucleic acid of bacteria and a deoxyribonuclease inhibitor. *Science* **124**:441–442.
- Chróst, R. J. 1975. Inhibitors produced by algae as an ecological factor affecting bacteria in water. II. Antibacterial activity of algae during blooms. *Acta Microbiol. Pol. B* **7**:167–176.
- Chróst, R. J. 1983. Plankton photosynthesis, extracellular release and bacterial utilization of released dissolved organic carbon (RDOC) in lakes of different trophic. *Acta Microbiol. Pol.* **32**:275–287.
- Davison, J. 1999. Genetic exchange between bacteria in the environment. *Plasmid* **42**:73–91.
- DeFlaun, M. F., J. H. Paul, and D. Davis. 1986. Simplified method for dissolved DNA determination in aquatic environments. *Appl. Environ. Microbiol.* **52**:654–659.
- Erata, M. 1997. Global Environmental Forum (GEF) list of strains, 5th ed., p. 128. National Institute for Environmental Studies, Environmental Agency, Tsukuba, Japan.
- Frischer, M. E., G. J. Stewart, and J. H. Paul. 1994. Plasmid transfer to indigenous marine bacterial populations by natural transformation. *FEMS Microbiol. Ecol.* **15**:127–136.
- Hara, T., and S. Ueda. 1981. A study on the mechanisms of DNA excretion from *P. aeruginosa* KYU-1: effect of mitomycin C on extracellular DNA production. *Agric. Biol. Chem.* **45**:2457–2461.
- Ishii, N., Z. Kawabata, S. Nakano, M. G. Min, and R. Takata. 1998. Microbial interactions responsible for dissolved DNA production in a hypereutrophic pond. *Hydrobiologia* **380**:67–76.
- Ishiwa, H., and H. Shibahara. 1985. New shuttle vectors for *Escherichia coli* and *Bacillus subtilis*. II. Plasmid pHY300PLK, a multipurpose cloning vector with a polylinker, derived from pHY460. *Jpn. J. Genet.* **60**:235–243.
- Iwane, T., T. Urase, and K. Yamamoto. 2001. Possible impact of treated wastewater discharge on incidence of antibiotic resistant bacteria in river water. *Water. Sci. Technol.* **43**:91–99.
- Jeffrey, W. H., J. H. Paul, and G. J. Stewart. 1990. Natural transformation of a marine *Vibrio* species by plasmid DNA. *Microb. Ecol.* **19**:259–269.
- Jørgensen, N. O. G., and C. S. Jacobsen. 1996. Bacterial uptake and utilization of dissolved DNA. *Aquat. Microb. Ecol.* **11**:263–270.
- Karl, D. M., and M. D. Bailiff. 1989. The measurement and distribution of dissolved nucleic acids in aquatic environments. *Limnol. Oceanogr.* **34**:543–558.
- Kawabata, Z., N. Ishii, M. Nasu, and M. G. Min. 1998. Dissolved DNA produced through a prey-predator relationship in a species-defined aquatic microcosm. *Hydrobiologia* **385**:71–76.
- Kearns, K. D., and M. D. Hunter. 2002. Algal extracellular products suppress *Anabaena flos-aeque* heterocyst spacing. *Microb. Ecol.* **43**:174–180.
- Kjelleberg, S., P. Steinberg, M. Givskov, L. Gram, M. Manefield, and R. de Nys. 1997. Do marine natural products interfere with prokaryotic AHL regulatory systems? *Aquat. Microb. Ecol.* **13**:85–93.
- Lorenz, M. G., D. Gerjets, and W. Wackernagel. 1991. Release of transforming plasmid and chromosomal DNA from two cultured soil bacteria. *Arch. Microbiol.* **156**:319–326.
- Lorenz, M. G., and W. Wackernagel. 1994. Bacterial gene transfer by natural genetic transformation in the environment. *Microbiol. Rev.* **58**:563–602.
- Matsui, K., M. Honjo, and Z. Kawabata. 2001. Estimation of the fate of dissolved DNA in thermally stratified lake water from the stability of exogenous plasmid DNA. *Aquat. Microb. Ecol.* **26**:95–102.
- Mazodier, P., and J. Davies. 1991. Gene transfer between distantly related bacteria. *Annu. Rev. Genet.* **25**:147–171.
- Ochman, H., J. G. Lawrence, and E. A. Groisman. 2000. Lateral gene transfer and the nature of bacterial innovation. *Nature* **405**:299–304.
- Paul, J. H., L. Cazares, and J. Thurmond. 1990. Amplification of *rbcL* gene from dissolved and particulate DNA from aquatic environments. *Appl. Environ. Microbiol.* **56**:1963–1966.
- Paul, J. H., and A. W. David. 1989. Production of extracellular nucleic acids by genetically altered bacteria in aquatic-environment microcosms. *Appl. Environ. Microbiol.* **55**:1865–1869.
- Paul, J. H., W. H. Jeffrey, and M. F. DeFlaun. 1987. Dynamics of extracellular DNA in the marine environment. *Appl. Environ. Microbiol.* **53**:170–179.
- Paul, J. H., S. C. Jiang, and J. B. Rose. 1991. Concentration of viruses and dissolved DNA from aquatic environments by vortex flow filtration. *Appl. Environ. Microbiol.* **57**:2197–2204.
- Paul, J. H., J. B. Rose, S. C. Jiang, C. A. Kellogg, and L. Dickson. 1993. Distribution of viral abundance in the reef environment of Key Largo, Florida. *Appl. Environ. Microbiol.* **59**:718–724.
- Sambrook, J., and D. W. Russell. 2000. Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Siuda, W., R. J. Chróst, and H. Güde. 1998. Distribution and origin of dissolved DNA in lakes of different trophic states. *Aquat. Microb. Ecol.* **15**:89–96.
- Siuda, W., and H. Güde. 1996. Determination of dissolved deoxyribonucleic acid concentration in lake water. *Aquat. Microb. Ecol.* **11**:193–202.
- Turk, V., A. S. Rehnstam, E. Lundberg, and Å. Hagström. 1992. Release of bacterial DNA by marine nanoflagellates, an intermediate step in phosphorus regeneration. *Appl. Environ. Microbiol.* **58**:3744–3750.
- Whitehead, N. A., A. M. L. Barnard, H. Slater, N. J. L. Simpson, and G. P. C. Salmund. 2001. Quorum-sensing in gram-negative bacteria. *FEMS Microbiol. Rev.* **25**:365–404.
- Weinbauer, M. G., D. Fuks, and P. Peuzzi. 1993. Distribution of viruses and dissolved DNA along a coastal trophic gradient in the northern Adriatic Sea. *Appl. Environ. Microbiol.* **59**:4074–4082.
- Wetzel, R. G. 2001. Limnology. Academic Press, San Diego, Calif.