

Aerobic Anoxygenic Photosynthesis in *Roseobacter* Clade Bacteria from Diverse Marine Habitats

Martin Allgaier,[†] Heike Uphoff, Andreas Felske, and Irene Wagner-Döbler*

Department of Microbiology, German Research Institute for Biotechnology, D-38124 Braunschweig, Germany

Received 17 December 2002/Accepted 26 May 2003

The marine *Roseobacter* clade comprises several genera of marine bacteria related to the uncultured SAR83 cluster, the second most abundant marine picoplankton lineage. Cultivated representatives of this clade are physiologically heterogeneous, and only some have the capability for aerobic anoxygenic photosynthesis, a process of potentially great ecological importance in the world's oceans. In an attempt to correlate phylogeny with ecology, we investigated the diversity of *Roseobacter* clade strains from various marine habitats (water samples, biofilms, laminariae, diatoms, and dinoflagellate cultures) by using the 16S rRNA gene as a phylogenetic marker gene. The potential for aerobic anoxygenic photosynthesis was determined on the genetic level by PCR amplification and sequencing of the *pufLM* genes of the bacterial photosynthesis reaction center and on the physiological level by detection of bacteriochlorophyll (Bchl) *a*. A collection of ca. 1,000 marine isolates was screened for members of the marine *Roseobacter* clade by 16S rRNA gene-directed multiplex PCR and sequencing. The 42 *Roseobacter* clade isolates found tended to form habitat-specific subclusters. The *pufLM* genes were detected in two groups of strains from dinoflagellate cultures but in none of the other *Roseobacter* clade isolates. Strains within the first group (the DFL-12 cluster) also synthesized Bchl *a*. Strains within the second group (the DFL-35 cluster) formed a new species of *Roseovarius* and did not produce Bchl *a* under the conditions investigated here, thus demonstrating the importance of genetic methods for screening of cultivation-dependent metabolic traits. The *pufL* genes of the dinoflagellate isolates were phylogenetically closely related to *pufL* genes from *Betaproteobacteria*, confirming similar previous observations which have been interpreted as indications of gene transfer events.


Phototrophy through bacteriochlorophyll (Bchl) *a*-mediated aerobic anoxygenic photosynthesis (40) has been estimated (based on in situ measurements of Bchl *a*) to be responsible for as much as 5 to 10% of the energy generation in the upper layers of the tropical oceans (12, 18, 19). The bacteria responsible for the process are thought to be related to the uncultivated marine SAR83 cluster within the alpha subclass of the *Proteobacteria*, which represents the second most abundant lineage of marine picoplankton bacteria after SAR11 (2, 11, 29). However, aerobic phototrophs have also been found in other genera of *Alphaproteobacteria* (*Erythrobacter*, *Roseivivax*, and *Sphingomonas*) and *Betaproteobacteria* (*Roseateles*) as well as among as yet uncultivated marine bacteria (2). In recent years, a large number of bacteria which are distantly related to the SAR83 cluster, and which form the *Roseobacter* clade, have been cultivated. Presently, 12 genera are recognized within this clade: *Ketogulonicigenium*, *Marinosulfonomonas*, *Jannaschia*, *Octadecabacter*, *Ruegeria*, *Staleyia*, *Roseobacter*, *Sulfitobacter*, *Antarctobacter*, *Sagittula*, *Roseovarius*, and *Silicibacter*. The genus *Roseivivax* (36) forms a separate lineage outside the *Roseobacter* clade (see Fig. 1) (E. Stackebrandt, personal communication). These bacteria were isolated from diverse marine or saline habitats, e.g., from marine sand, sediments, and mac-

roalgae (31), toxin-producing dinoflagellates (22, 27), the Black Sea (33), the North Sea (39), a meromictic lake (42), black smokers (40), and hot springs (26). Accordingly, their described physiological properties are highly diverse, ranging from sulfite reduction (33) to metabolism of dimethylsulfoniopropionate (43, 44) and production of toxins (10, 15). Only four species with the capability for aerobic anoxygenic photosynthesis are known to date, namely, *Roseobacter litoralis* and *Roseobacter denitrificans*, isolated from marine algae and sediments (31), and *Staleyia guttiformis* (21) and *Roseovarius tolerans* (20), both from the hypersaline antarctic Ekho Lake.

Speciation in bacteria is thought to be driven by the occupation of a new ecological niche (4). Thus, within a described species, ecotypes can be found with highly similar or even identical 16S rRNA gene sequences but clearly different ecological adaptations. Several examples have been described among pathogenic bacteria (7). In *Neisseria meningitidis*, 10 ecotypes, which occurred in different geographical areas and had different outbreak characteristics, could be identified by multilocus sequence typing (6). In surveys of as yet uncultured bacteria, clusters of highly related 16S rRNA gene sequences have frequently been described. In some instances, experimental evidence showed that these corresponded to different populations inhabiting distinct microenvironments. For example, clusters of uncultured *Synechococcus* strains from Yellowstone hot springs inhabited distinct microniches defined by temperature, photic zone, and stage of succession (28). Similar observations have been made among cultured marine organisms; small differences in 16S rRNA gene sequences of *Prochlorococcus* strains have been shown to correspond to high-light- and low-light-adapted populations (24), which have major

* Corresponding author. Mailing address: German Research Institute for Biotechnology (GBF), Department of Microbiology, Mascheroder Weg 1, D-38124 Braunschweig, Germany. Phone: 49 531 6181 408. Fax: 49 531 6181 974. E-mail: iwd@gbf.de.

[†] Present address: Leibniz-Institut für Gewässerökologie und Binnenfischerei (IGB), Abteilung Limnologie geschichteter Seen, 16775 Stechlin-Neuglobsow, Germany.



genomewide genetic differences (<http://web.mit.edu/chisholm/www/index.html>).

The spectrum of ecological niches described in the literature for *Roseobacter* clade bacteria is huge, ranging from oligotrophic picoplankton (an uncultivated SAR83 cluster) to microalgal symbioses, but no systematic survey has been undertaken so far. Here we attempted to correlate ecology with phylogeny, by using the 16S rRNA gene as a phylogenetic marker gene and the genetic potential for aerobic anoxygenic photosynthesis as a cultivation-independent functional trait, for a large collection of marine isolates from various North Sea habitats. Our aims were to identify phylogenetic clusters of *Roseobacter* clade bacteria which were typical for certain habitats and to determine the capability for aerobic anoxygenic photosynthesis throughout these clusters.

MATERIALS AND METHODS

Sample collection and processing. Samples were obtained from the North Sea around the island of Helgoland during sampling trips in 1998, 1999 and 2002. The research vessel Aade or Diker was used for sampling except for the rocky littoral, which could be reached by foot. Water samples were taken with a Ruttner sampler at the specified depth. A list of strains, sampling dates, sampling locations, and media used for first cultivation of isolates is given in Table 1. With the exception of strains designated PIC (for picoplankton), isolates were obtained by direct plating of aliquots (50 μ l) of the highest sample dilution on the media indicated. Serial dilutions were made in artificial seawater or phosphate-buffered saline (PBS) (93.2 ml of 1 M Na₂HPO₄, 6.8 ml of 1 M NaH₂PO₄, and 5.8 g of NaCl per 1,000 ml).

For enrichment of PIC strains, 50 liters of the water sample was filtered through 5- μ m-pore-size nylon mesh filters on the sampling day and stored in polyvinyl chloride containers in the dark at 4°C until enrichment was started (38). The isolates from the *Roseobacter* clade were enriched as follows. Aliquots (20 ml) of the filtered North Sea water sample were amended with tryptone and yeast extract (each at 1.0 g/liter) and incubated either at room temperature in natural daylight without shaking (PIC-68, PIC-69, and PIC-70), at room temperature in natural daylight with shaking (PIC-74), at room temperature in the dark without shaking (PIC-76), or at 4°C in the dark without shaking (PIC-82). After growth was visible, serial dilutions were plated on medium 1 (see Table 2).

For isolation of strain BIO-7, a water sample of 7 liters was taken from a depth of 3 m and allowed to settle overnight. Sedimented particles were collected by using a glass pipette, serially diluted, and plated. For strains BIO-11, BIO-13, BIO-24, BIO-204, and BIO-205, biofilms were grown in situ by exposing sterilized glass plates (200 by 200 by 3 mm) mounted in a biofilm collector (3) to the North Sea on 26 July 1999 at two different locations at a depth of 3 m. After 1 or 2 weeks, biofilms were aseptically scraped off, serially diluted, and plated.

For isolation of strain DT-12, a plankton sample was taken by using a 170- μ m-mesh-size net pulled behind a boat for 5 min. Diatoms (*Thalassiosira* sp.) were picked under a dissecting microscope and washed three times (see below). Approximately 30 cells were suspended in 100 μ l of artificial seawater, and aliquots were plated without dilution.

For isolates designated LM, seaweeds (a *Laminaria* sp.) from a small pool (depth, 60 cm) in the rocky littoral zone of the island of Helgoland ("Felswatt") were sampled during low tide. Cauloids were cut off. Phylloids were washed several times in artificial seawater. The surface biofilm was scraped off with a sterile blade, suspended in 5 ml of artificial seawater, mixed vigorously, serially diluted, and plated.

Strains designated DFL were obtained from the following dinoflagellate cultures of the Biological Research Institute of Helgoland: *Prorocentrum lima* ME130, *Alexandrium lusitanicum* ME207, and *Alexandrium ostenfeldii* KO287. Cultures were grown in F-2 medium (14) at 18°C with a cycle of 12 h of light and 12 h of darkness. The algal cells were picked by using a 1.5- μ l Gilson pipette and transferred to a petri dish with artificial seawater (ca. 5 ml). They were washed by vigorous mixing, and the individual cells were subsequently transferred to a new petri dish. This procedure was repeated twice. Approximately 10 washed cells were finally pooled in 100 μ l of artificial seawater and plated without dilution.

Isolation and cultivation of strains. Strains were isolated on solid marine media with relatively low concentrations of carbon sources (Table 2). Samples for isolation of HEL strains were plated on medium 1, 2, 3, or 9; those for PIC and BIO strains were plated on medium 1, 3, 9, or 10 or the medium given by Uphoff et al. (38); and those for DT, DFL, and LM strains were plated on media 4 to 8. For the composition of artificial seawater, Hutner's salt solution, and vitamin solution, see the information on DSMZ media 607, 590, and 603, respectively, on the Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ) website (<http://www.dsmz.de/media/media.htm>). All media contained cycloheximide (25 mg liter⁻¹) to prevent the growth of eukaryotes and were solidified with agar (15 g liter⁻¹). Media were sterilized by autoclaving at 121°C for 20 min. Unfiltered seawater was autoclaved before use. Heat-sensitive components, such as antibiotics and vitamins, were added to autoclaved media from filter-sterilized (pore size, 0.2 μ m) stock solutions. Picked colonies were restreaked on the same medium two to three times until they were pure. Subsequently, they were cultivated on Marine Broth 2216 (Difco). *Roseovarius tolerans* (DSMZ 11457), *S. guttiformis* (DSMZ 11458), *Roseobacter litoralis* (DSMZ 6996), and *Roseobacter denitrificans* (DSMZ 7001) were used as reference strains. The *Roseobacter* strains were grown on DSMZ medium 695 (<http://www.dsmz.de/media/media.htm>), while *Roseovarius tolerans* and *S. guttiformis* were grown on medium 8 (Table 2) with 25 and 10% artificial seawater, respectively. In addition to the type strain of *Roseovarius tolerans* (EL-172¹), seven other isolates of *Roseovarius tolerans* were investigated (EL-52, EL-78, EL-83, EL-90, EL-164, EL-171, and EL-222) (personal gifts from Matthias Labrenz).

DNA extraction. Genomic DNA was extracted from a loopful of bacterial cells from a fresh plate by using the NucleoSpin tissue kit (Macherey-Nagel, Düren, Germany) with an additional step for RNase digestion after cell lysis. To this end, 4 μ l of RNase (Roche, Basel, Switzerland) was added and incubated at 37°C for 30 min, followed by incubation at 95°C for 10 min to denature the enzyme. Genomic DNA was eluted by using 100 μ l of elution buffer and stored at -20°C. Strains which did not yield enough genomic DNA by this method were extracted by using the FastDNA SPIN Kit for Soil from Bio 101 (now called Qbiogene, Heidelberg, Germany). For cell lysis, a bead beater (Retsch, Haan, Germany) was used instead of the FastPrep instrument. Genomic DNA was eluted with 50 μ l of DNase-pyrogen-free water and stored at -20°C.

Amplification of 16S rRNA and *pufLM* genes. The primer pair 16F 27 (5'-A GAGTTTGATCCTGGCTCAG-3') and 16R 1492 (5'-TAC GGY TAC CTT GTT ACG ACT T-3') was used for PCR amplification of the nearly complete 16S rRNA gene (23). The reaction mixture contained 1.5 mM MgCl₂, 0.25 μ M each deoxynucleoside triphosphate, 0.5 μ M each primer, 1 μ l of template DNA, and 5 U of *Taq* DNA polymerase (Qiagen, Hilden, Germany) in a total volume of 50 μ l. PCR was performed in a Landgraf (Langenhagen, Germany) thermocycler or a Mastercycler gradient (Eppendorf, Hamburg, Germany) by using an initial denaturation step at 94°C (2 min), followed by 30 cycles of denaturation at 94°C (1 min), annealing at 54°C (1 min), and extension at 72°C (2 min). A final extension at 72°C (10 min) and subsequent cooling at 4°C completed the reaction. The amplified DNA was purified using the QIAquick PCR Purification kit (Qiagen), eluted with 50 μ l of H₂O, and stored at -20°C.

The primer pair *pufLF* (5'-CTK TTC GAC TTC TGG GTS GG-3') and *pufMR* (5'-CCA TSG TCC AGC GCC AGA A-3') (K is G or T and S is G or

FIG. 1. Phylogenetic tree of the *Roseobacter* clade within the alpha subclass of the *Proteobacteria*. Reference organisms are italicized, and marine isolates described in this work are boldfaced. Representative environmental clones from uncultivated organisms of the Sargasso Sea (env.SAR), the Pacific Ocean (env.OCS), the Mediterranean Sea (env.Adriatic), the continental shelf of the Atlantic Ocean (env.OM), and the Columbia River estuary (env.CRE) were included (29). In addition, isolates from toxic dinoflagellates (667-19 and 253-13) and a marine sponge (HRV3) were also included. Strains in which the *pufLM* genes as well as *Bchl a* were detected are marked with asterisks. Strains which contained the *pufLM* genes but not *bchl a* are marked with a Δ . Except for environmental clones and isolate HRV3, the tree is based on near-complete 16S rRNA gene sequences and was calculated using the maximum-likelihood algorithm (8). Bootstrap values above 50% are given. The bar corresponds to 10 base substitutions per 100 nucleotide positions. *Rhodobacter veldkampii* was used as an outgroup.

TABLE 1. List of isolated *Roseobacter* clade strains and sampling information^a

Strain ^b	Type of sample	Sampling date (mo/day/yr)	Sampling site	Medium ^c
Bacterioplankton				
HEL-8	Water sample	9/23/98	Deep ridge ^d	1
HEL-10				1
HEL-26				1
HEL-43				1
HEL-45				1
HEL-46				1
HEL-76				2
HEL-77				2
HEL-78				2
PIC-68	Filtered water sample	9/23/98	Deep ridge	1
PIC-69				1
PIC-70				1
PIC-72				1
PIC-74				1
PIC-76				1
PIC-82				1
Biofilms				
BIO-7	Suspended particles	7/26/99	Dune ^e	3
BIO-11	In situ biofilm (1 wk) ^f		South harbor ^g	3
BIO-13	In situ biofilm (1 wk)		Dune	1
BIO-24	In situ biofilm (2 wk)		Dune	1
BIO-204	In situ biofilm (1 wk)		Dune	10
BIO-205	In situ biofilm (2 wk)		Dune	9
Diatoms, DT-12				
	<i>Thalassiosira</i> sp.	4/4/02	Cable ton ^h	4
Macroalgal surface				
LM-4	<i>Laminaria</i> sp.	4/7/02	Rocky littoral ⁱ	5
LM-7				8
LM-9				8
LM-10				8
LM-11				8
LM-16				8
Dinoflagellates				
DFL-10	<i>A. ostenfeldii</i>	April 2002	Culture KO287	8
DFL-12	<i>P. lima</i>		Culture ME130	6
DFL-14	<i>A. ostenfeldii</i>		Culture KO287	6
DFL-16	<i>A. ostenfeldii</i>		Culture KO287	6
DFL-23	<i>A. lusitanicum</i>		Culture ME207	5
DFL-24	<i>A. ostenfeldii</i>		Culture KO287	8
DFL-27	<i>A. ostenfeldii</i>		Culture KO287	7
DFL-30	<i>A. ostenfeldii</i>		Culture KO287	6
DFL-31	<i>A. ostenfeldii</i>		Culture KO287	6
DFL-35	<i>A. ostenfeldii</i>		Culture KO287	8
DFL-36	<i>A. ostenfeldii</i>		Culture KO287	6
DFL-38	<i>A. ostenfeldii</i>		Culture KO287	6
DFL-41	<i>A. ostenfeldii</i>		Culture KO287	6

^a Samples were taken in the North Sea around the island of Helgoland during trips in 1998, 1999, and 2002. For details on isolation of strains, see Materials and Methods.

^b For each set of related strains (with designations beginning with the same letters), the type of sample, sampling date, or sampling site is given only in the first row if it is the same in all cases.

^c For descriptions of media, see Table 2.

^d For the deep ridge (Tiefe Rinne), samples were taken at 54°08'N, 7°52'E; depth, 10 m; temperature, 15.5°C; oxygen, 8.1 mg/liter; Secchi disk visibility, 5.5 m.

^e For the dune (Düne), samples were taken at 54°11.42'N, 7°53.51'E; depth, 3 m; temperature, 19.6°C; pH 8.22; conductivity, 44 mS/cm; oxygen, 6.2 mg/liter.

^f In situ biofilms were grown for 1 or 2 weeks.

^g For the south harbor (Südhafen), samples were taken at 54°10.13'N, 7°53.55'E; depth, 3 m.

^h For the cable ton (Kabeltonne), samples were taken at 54°11.18'N, 7°54.00'E; depth, 0.5 to 1.5 m; temperature, 6.5°C; Secchi disk visibility, 2.8 m.

ⁱ For the rocky littoral (Felswatt), samples were taken at a depth of 0.5 m and a water temperature of 8.5°C.

C) was used for amplification of the *pufLM* genes from the bacterial photosynthesis reaction center (2). The reaction mixture was the same as that described above, except that 2 U of *Taq* DNA polymerase was used. The reaction volume was 20 μ l (for screening of strains) or 50 μ l (for excision of bands). The PCR program was the same as that described above, except that initial denaturation at 94°C was performed for 3 min, and the annealing temperature was 60°C. The amplified *pufLM* fragment (ca. 1.5 kb) was excised from the agarose gel under

UV light and extracted by using the QIAquick Gel Extraction kit (Qiagen) according to the manufacturer's instructions. Extracted DNA was eluted with 30 μ l of H₂O and stored at -20°C.

Sequencing. Sequencing was performed with an automated capillary sequencer, the ABI PRISM 3100 Genetic Analyzer (Applied Biosystems, Weiterstadt, Germany), according to the manufacturer's instructions. The sequencing reaction contained 5 μ l of DNA, 1 μ l of primer (10 μ M), 5 μ l of the reagent from

TABLE 2. List of media^a used for the first cultivation of marine *Roseobacter* clade strains

Medium no.	Medium description	Main carbon sources	Other components	Salts
1	Cytophaga (marine) medium, DSMZ medium 172 ^b	Tryptone (1.0 g), yeast extract (1.0 g)		Sea salts
2	Marine R2A medium ^c	Peptone (0.5 g), yeast extract (0.5 g)	Casamino Acids (0.5 g), dextrose (0.5 g), soluble starch (0.5 g), sodium pyruvate (0.3 g)	75% artificial seawater
3	M13 Verrucomicrobium medium, DSMZ medium 607	Peptone (0.25 g), yeast extract (0.25 g)	Glucose (0.25 g), vitamin solution (10 ml), Hutner's salts (20 ml)	25% artificial seawater
4	Marine broth 2216 (Difco), 1/10 strength	Peptone (0.5 g), yeast extract (0.1 g)		Tap water
5	Luria-Bertani broth (Sigma) in North Sea water	Tryptone (10.0 g), yeast extract (5.0 g)		Unfiltered North Sea water
6	Luria-Bertani broth (Sigma) in North Sea water, 1/10 strength	Tryptone (1.0 g), yeast extract (0.5 g)		Unfiltered North Sea water
7	Peptone, yeast extract, North Sea water	Peptone (0.04 g), yeast extract (0.008 g)		Filtered North Sea water (pore size, 3 µm)
8	Peptone, yeast extract, glucose, vitamins, 30% North Sea water	Peptone (0.25 g), yeast extract (0.25 g)	Glucose (0.25 g), vitamin solution (10 ml), Hutner's salts (20 ml)	Unfiltered North Sea water (300 ml)
9	Hg medium	Peptone (0.25 g), yeast extract (0.25 g)	HgCl ₂ (1.0 mg)	Filtered North Sea water (pore size, 0.2 µm)
10	North Sea water			Unfiltered North Sea water

^a Amounts are given for a total volume of 1 liter. All media additionally contained 15 g of agar (Difco) and 25 mg of cycloheximide (Sigma).

^b Information on DSMZ media is available at <http://www.dsmz.de/media/media.htm>.

^c From the work of Suzuki et al. (35).

the sequencing kit (ABI PRISM Dye Terminator Cycle Sequencing Ready Reaction kit; Applied Biosystems), 1 µl of dimethyl sulfoxide, and 8 µl of H₂O. The sequencing PCR was performed with an Eppendorf (Hamburg, Germany) Mastercycler gradient by using 25 cycles consisting of denaturation (at 96°C for 15 s), annealing (at 60°C for 15 s), and extension (at 60°C for 240 s). For sequencing of the 16S rRNA gene, primers 16F 27 and 16R 1492 were used; for sequencing of the *pufLM* genes, primers *pufLF* and *pufMR* were used. After amplification, the DNA was cleaned by ethanol precipitation as follows. The reaction mixture was transferred to a 0.5-ml Eppendorf reaction tube containing 10 µl of sodium acetate (3 M) and 250 µl of ethanol (98%, vol/vol). After vigorous mixing, the DNA was precipitated by centrifugation (at 14,000 × *g* for 30 min). The supernatant was discarded, and the pellet was washed in 300 µl of ethanol (70%, vol/vol). The DNA was precipitated by centrifugation (at 14,000 × *g* for 10 min), the supernatant was discarded, and the pellet was dried for 5 min in a SpeedVac.

Phylogenetic inferences. Sequences were analyzed by using the ARB software package (<http://www.arb-home.de>). Partial 16S rRNA gene sequences were imported into the ARB database of ca. 12,600 aligned reference sequences. Partial sequences were automatically aligned and assembled into full sequences, and alignments were corrected manually. Full sequences were integrated into the ARB phylogenetic tree by using the maximum parsimony algorithm tool (9). This tool does not correct for evolutionary distances and does not change the tree topology. It served as a basis for identification of the reference organisms. These were subsequently selected for tree calculation by means of the maximum-likelihood algorithm (8) by using the fastDNAmI program (25). Partial *pufLM* sequences were aligned using ClustalW (37). The alignment was imported into ARB. A phylogenetic tree was calculated in ARB using the maximum-likelihood method. Almost full length 16S rRNA gene sequences and partial *pufLM* sequences were checked for homologies to published sequences by BLAST searches (www.ncbi.nlm.nih.gov). Similar sequences were downloaded, imported into ARB, and used for calculation of phylogenetic trees.

Determination of Bchl *a*. Bchl *a* was detected spectrophotometrically in vivo or in vitro. For in vivo measurements, 2 ml of liquid culture (Marine Broth 2216; Difco) incubated in the dark or in a natural daylight rhythm was washed (at 7,000 rpm for 5 min, with resuspension of the pellet in 1 ml of PBS). To reduce light scattering (32), 1 ml of bovine serum albumin (30%, wt/vol; Sigma) was added. For in vitro measurements, Bchl *a* was extracted by the procedure of Cohen-Bazire et al. (5). Two microliters of a liquid culture was centrifuged (at 7,000 rpm for 5 min), and the pellet was resuspended in a drop of remaining medium. A 1.5-ml volume of an ice-cold (−20°C) acetone-methanol solution (7:2, vol/vol) was added, mixed well, and incubated at room temperature in the dark for 12 h.

Cell fragments were then removed by centrifugation (at 2,500 rpm for 5 min). Spectrophotometric measurements were performed with an Ultraspec K 4053 instrument from LKB Biochrom (Cambridge, United Kingdom). Relative absorption was determined after calibration at 900 nm against air in the range from 400 to 900 nm (in vivo) or 600 to 900 nm (in vitro).

Nucleotide sequence accession numbers. Sequences of *Roseobacter* clade bacterium 16S rRNA genes sequenced in this study can be found in the EMBL data bank under accession numbers AJ534205 to AJ534244 and AJ532580. *pufL* sequences can be found under accession numbers AJ532572 to AJ532579.

RESULTS AND DISCUSSION

Isolation of strains. A total of 1,019 isolates were obtained from different marine habitats, mainly from the North Sea around the island of Helgoland. Of these, 71 were isolated from an untreated seawater sample (strains designated HEL), 412 were enriched from the same water sample after filtering (PIC strains), 463 were obtained from biofilms grown in situ in the North Sea on glass plates (BIO strains), 12 were isolated from the surfaces of *Thalassiosira* sp. diatoms (DT strains), 44 were isolated from cultures of *A. ostentfeldii*, *A. lusitanicum*, or *P. lima* (Dinophyta) (DFL strains), and 17 were isolated from *Laminaria* sp. surfaces (LM strains). In the latter three cases, pigmented colonies were picked selectively, while HEL, BIO, and PIC strains were picked independently of their pigmentation. Bacteria were isolated by direct plating of the fresh, untreated sample or the highest sample dilution on solid marine media, with the exception of PIC strains, which were enriched under a wide spectrum of incubation and amendment conditions (38). The media used for isolation of strains had carbon contents that were relatively low, though high in comparison to those used for the first cultivation of SAR11 (30), and differed in their salt contents (Table 2). Unamended, autoclaved North Sea water was also used as a culture medium.

Screening for *Roseobacter* clade strains. All strains were screened for *Roseobacter* clade phylogenetic affiliation by multiplex PCR targeting the 16S rRNA gene. This PCR results in amplicons of a certain size for selected phylogenetic groups. The primer mix used produced a 650-bp amplicon in *Alphaproteobacteria* and a 250-bp fragment in *Roseobacter* clade bacteria (38). Positive strains were partially sequenced, and false positives were excluded from further analysis. Forty-two strains that were assigned to the *Roseobacter* clade based on partial 16S rRNA gene sequencing were fully sequenced.

Phylogenetic affiliation of strains. Of the 42 strains analyzed, 9 had been isolated from a North Sea water sample, 7 from picoplankton enrichment cultures, 6 from biofilms, 6 from *Laminaria* surfaces, 1 from *Thalassiosira* surfaces, and 13 from dinoflagellate cultures (Table 1). Figure 1 shows that the isolates tended to form clusters of related organisms from the same habitat. These clusters were distributed over the whole *Roseobacter* clade and represented new genera (e.g., *Jannaschia helgolandensis* [39]) or species. Only in one case did a cluster contain a strain highly related, but not identical, to a strain which had been isolated from a different habitat: BIO-205, from biofilms in the cluster around HEL-45 from the water column. Among isolates from dinoflagellates, strains derived from *A. ostenfeldii* cultures (DFL-36, DFL-38, DFL-30, DFL-27, DFL-16, and DFL-31) were more closely related to each other than to the isolate from a *P. lima* culture (DFL-12). Similar observations have been interpreted previously as the result of coevolution (1). The data might point to a process of evolution driven by ecological adaptation which has resulted in clusters of phylogenetically related strains specifically adapted to the ecological niche in question, e.g., biofilm, dinoflagellate, or picoplankton.

Almost all cloned environmental sequences within the *Roseobacter* clade fell into a separate branch containing SAR83 from the Sargasso Sea as well as sequences from the Mediterranean Sea, Atlantic Ocean, and Pacific Ocean. They formed three subclusters, which were affiliated with recently described genera, i.e., *Octadecabacter* (13), isolated from arctic and antarctic ice; *Marinosulfonomonas methylotropa* (16); and *J. helgolandensis* from the North Sea (39). Even more similar to the third and largest cluster of SAR83-related sequences than *J. helgolandensis* were isolates obtained from dinoflagellate cultures (the DFL-12 cluster). The level of similarity to *J. helgolandensis* was 92 to 93%, indicating at least genus level divergence. However, the phylogenetic affiliation of the environmental sequences, as well as the grouping of genera within the *Roseobacter* clade, was not robust, as indicated by the low bootstrap values, and thus should not be overestimated.

Genetic potential for aerobic anoxygenic photosynthesis in *Roseobacter* clade marine isolates. All marine isolates shown in Fig. 1 as well as several type strains were screened for the presence of the *pufLM* genes of the photosynthesis reaction center by PCR amplification and sequencing. Their presence was confirmed in *Roseobacter denitrificans*, *Roseobacter litoralis*, and *S. guttiformis*. Interestingly, the *pufLM* genes were found in all eight strains of *Roseovarius tolerans* tested. When first describing *Roseovarius tolerans*, Labrenz et al. (20) could detect Bchl *a* in only four strains (EL-78, EL-83, EL-171, and EL-172^T) and therefore named the genus *Roseovarius*. In addition, *pufLM* genes were found in two dinoflagellate isolates

(DFL-24 and DFL-35) within the genus *Roseovarius* (probably representing a new species) and in the DFL-12 cluster of strains, which were also isolated from dinoflagellate cultures. Interestingly, none of the other *Roseobacter* clade marine isolates had *pufLM* genes. However, these genes were also detected in other *Alphaproteobacteria* (Fig. 2A). Some strains that had the *pufLM* genes and were also isolated from dinoflagellates formed a cluster related to *Ahrensia kielensis* (DFL-13, DFL-44, DFL-42, and DFL-43), and another isolate was related to *Roseibium hamelinense* (DFL-11). Moreover, a water sample isolate related to *Sphingomonas alaskensis* (HEL-42) also had *pufLM* genes.

Production of Bchl *a*. Bchl *a* was detected in freshly grown cells and methanol extracts of all DFL-12 cluster strains. It was not found in the two other *Roseobacter* clade strains that had the *pufLM* genes (DFL-24 and DFL-35) or in the isolates outside the *Roseobacter* clade that had the *pufLM* genes (DFL-13, DFL-44, DFL-42, DFL-43, DFL-11, and HEL-42). Moreover, of the eight strains of *Roseovarius tolerans*, only four were reported to produce Bchl *a* (20). Thus, of the 16 strains analyzed in this study which had the *pufLM* genes, only 7 produced Bchl *a* under the conditions investigated, i.e., growth in liquid culture in the dark or in the natural light-dark cycle in full-strength marine broth. Expression of Bchl *a* synthesis in aerobic, anoxygenic phototrophs is regulated by environmental parameters (41). It is completely suppressed by light. In addition, oxygen concentration and carbon limitation have been shown to be involved in *Roseateles depolymerans* photosynthesis (34). It might be expected that production of Bchl *a* can be induced in most of the isolates that have *pufLM* genes if the "right" conditions are found. For example, by growing the dinoflagellate isolates in 1/10 marine broth, pigmented cultures could be obtained for some strains, indicating that nutrient concentration and/or salt concentration may play a role in induction of Bchl *a* synthesis.

Phylogeny of *pufL*. Sequence alignment and phylogenetic analysis of the *pufL* fragments from the marine isolates and of closely related published *pufL* sequences (Fig. 2) showed several differences between the 16S rRNA gene phylogenetic tree and the *pufL*-based tree, except for the highest level of similarity. Thus, clusters of related or identical *pufL* sequences (the DFL-12 cluster, *Roseovarius* cluster, DFL-35 cluster, and DFL-13 cluster) corresponded to clusters of related or identical 16S rRNA gene sequences. However, the DFL-35 cluster was most closely related to *Roseovarius tolerans* on the basis of 16S rRNA gene sequence analysis but to *Roseobacter denitrificans* on the basis of *pufL* alignment. Similarly, *S. guttiformis* was monophyletic with *Roseobacter litoralis* and *Roseobacter denitrificans* on the basis of 16S rRNA gene analysis, while *Rhodobacter* formed a separate lineage. By contrast, *S. guttiformis* and *Rhodobacter sphaeroides* were monophyletic in the *pufL* gene-based tree, while *Roseobacter litoralis*, *Roseobacter denitrificans*, and *Roseovarius tolerans* formed a separate lineage. For phylogenetic groups at even higher taxonomic levels, the tree topology showed differences between the two marker genes. Subgroups 1 to 4 of the alpha proteobacteria were intermixed in the *pufL* gene-based tree, and the *pufL* sequence from the beta proteobacterium *Roseateles depolymerans* affiliated with those of *Alphaproteobacteria*. This phenomenon has been observed before (17) and has been interpreted as the




FIG. 2. Sequence similarities among *pufL* partial sequences (A) and 16S rRNA gene sequences (B) from reference strains of phototrophic alpha- and beta-subclass proteobacteria, marine bacteria described in this study (boldfaced), and phototrophic bacteria described by Beja et al. (2). The tree was calculated by using the ARB software package (<http://www.arb-home.de>) and is based on the maximum-likelihood algorithm (8). For the 16S rRNA gene-based tree, near-complete sequences were used and a protobacterial filter was applied. Bootstrap values above 70% are given in the *pufL* gene-based tree, and those above 50% are given in the 16S rRNA gene-based tree. The bar corresponds to 10 base substitutions per 100 nucleotide positions. The gamma proteobacterium *Allochromatium vinosum* was used as an outgroup in both trees.

result of gene transfer of the photosynthesis gene cluster from an ancient alpha proteobacterium into the *Betaproteobacteria*, especially since it turned out that the photosynthesis genes are highly clustered and arranged in a similar fashion in different phototrophic bacteria (17). Recently it was suggested that there is a large diversity of uncultured aerobic phototrophic proteobacteria, some of them within the beta-proteobacterial subgroup (2). It remains to be seen how these new sequences will affect overall tree topology.

Conclusions. The data show that dinoflagellates represent an important ecological niche for aerobic anoxygenic phototrophic *Alphaproteobacteria*. Since dinoflagellates are photoautotrophic organisms themselves and can swim actively, they probably provide a suitable light regimen for the associated bacteria.

ACKNOWLEDGMENTS

We thank Gunnar Gerdts for logistic support. Thanks to the divers of the BAH Helgoland, Udo Schilling and Carsten Wanke, and the crew of the research vessels Diker and Ade for sampling. Special thanks to Matthias Labrenz for providing strains and to Hanno Biebl for many helpful discussions.

REFERENCES

- Ashen, J. B., and L. J. Goff. 2000. Molecular and ecological evidence for species specificity and coevolution in a group of marine algal-bacterial symbioses. *Appl. Environ. Microbiol.* **66**:3024–3030.
- Beja, O., M. T. Suzuki, J. F. Heidelberg, W. C. Nelson, C. M. Preston, T. Hamada, J. A. Eisen, C. M. Fraser, and E. F. DeLong. 2002. Unsuspected diversity among marine aerobic anoxygenic phototrophs. *Nature* **415**:630–633.
- Brümmer, I. H. M., W. Fehr, and I. Wagner-Dobler. 2000. Biofilm community structure in polluted rivers: abundance of dominant phylogenetic groups over a complete annual cycle. *Appl. Environ. Microbiol.* **66**:3078–3082.
- Cohan, F. M. 2002. What are bacterial species? *Annu. Rev. Microbiol.* **56**:457–487.
- Cohen-Bazire, G., W. R. Sistrom, and R. Y. Stanier. 1957. Kinetic studies of pigment synthesis by non-sulfur purple bacteria. *J. Cell. Comp. Physiol.* **49**:25–68.
- Feil, E. J., M. C. Maiden, M. Achtman, and B. G. Spratt. 1999. The relative contributions of recombination and mutation to the divergence of clones of *Neisseria meningitidis*. *Mol. Biol. Evol.* **16**:1496–1502.
- Feil, E. J., and B. G. Spratt. 2001. Recombination and the population structures of bacterial pathogens. *Annu. Rev. Microbiol.* **55**:561–590.
- Felsenstein, J. 1981. Evolutionary trees from DNA sequences: a maximum likelihood approach. *J. Mol. Evol.* **17**:368–376.
- Fitch, W. 1971. Toward defining the course of evolution: minimum change for a specific tree topology. *Syst. Zool.* **20**:406–416.
- Gallacher, S., K. J. Flynn, J. M. Franco, E. E. Brueggemann, and H. B. Hines. 1997. Evidence for production of paralytic shellfish toxins by bacteria associated with *Alexandrium* spp. (Dinophyta) in culture. *Appl. Environ. Microbiol.* **63**:239–245.
- Giovannoni, S. J., and M. S. Rappé. 2000. Evolution, diversity and molecular ecology of marine prokaryotes, p. 47–84. *In* D. L. Kirchman (ed.), *Microbial ecology of the ocean*. John Wiley & Sons, Inc., New York, N.Y.
- Goerick, R. 2002. Bacteriochlorophyll *a* in the ocean: is anoxygenic bacterial photosynthesis important? *Limnol. Oceanogr.* **47**:290–295.
- Gosink, J. J., R. P. Herwig, and J. T. Staley. 1997. *Octadecabacter arcticus* gen. nov., sp. nov., and *O. antarcticus*, sp. nov., nonpigmented, psychrophilic gas vacuolate bacteria from polar sea ice and water. *Syst. Appl. Microbiol.* **20**:356–365.
- Guillard, R. R. L. 1975. Culture of phytoplankton for feeding marine invertebrates, p. 26–60. *In* W. L. Smith and M. H. Chanley (ed.), *Culture of marine invertebrate animals*. Plenum Press, New York, N.Y.
- Hold, G. L., E. A. Smith, M. S. Rappé, E. W. Maas, E. R. B. Moore, C. Stroempl, J. R. Stephen, J. Prosser, T. H. Birkbeck, and S. Gallacher. 2001. Characterisation of bacterial communities associated with toxic and non-toxic dinoflagellates: *Alexandrium* spp. and *Scrippsiella trochoidea*. *FEMS Microbiol. Ecol.* **37**:161–173.
- Holmes, A. J., D. P. Kelly, S. C. Baker, A. S. Thompson, P. DeMarco, E. M. Kenna, and J. C. Murrell. 1997. *Methylsulfolobomonas methylavora* gen. nov., sp. nov., and *Marinosulfolobomonas methylotropa* gen. nov., sp. nov.: novel methylotrophs able to grow on methanesulfonic acid. *Arch. Microbiol.* **167**:46–53.
- Igarashi, N., J. Harada, S. Nagashima, K. Matsuura, K. Shimada, and K. V. Nagashima. 2001. Horizontal transfer of the photosynthesis gene cluster and operon rearrangement in purple bacteria. *J. Mol. Evol.* **52**:333–341.
- Kolber, Z. S., C. L. Van Dover, R. A. Niederman, and P. G. Falkowski. 2000. Bacterial photosynthesis in surface waters of the open ocean. *Nature* **407**:177–179.
- Kolber, Z. S., F. G. Plumley, A. S. Lang, J. T. Beatty, R. E. Blankenship, C. L. Van Dover, C. Vetriani, M. Koblizek, C. Rathgeber, and P. G. Falkowski. 2001. Contribution of aerobic photoheterotrophic bacteria to the carbon cycle in the ocean. *Science* **292**:2492–2495.
- Labrenz, M., M. D. Collins, P. A. Lawson, B. J. Tindall, P. Schumann, and P. Hirsch. 1999. *Roseovarius tolerans* gen. nov., sp. nov., a budding bacterium with variable bacteriochlorophyll *a* production from hypersaline Ekho Lake. *Int. J. Syst. Evol. Microbiol.* **49**:137–147.
- Labrenz, M., B. J. Tindall, P. A. Lawson, M. D. Collins, P. Schumann, and P. Hirsch. 2000. *Staleyia guttiformis* gen. nov., sp. nov. and *Sulfitobacter brevis* sp. nov., alpha-3-Proteobacteria from hypersaline, heliothermal and meromictic antarctic Ekho Lake. *Int. J. Syst. Evol. Microbiol.* **50**:303–313.
- Lafay, B., R. Ruimy, C. R. Detraubenberg, V. Breitmayer, M. J. Gauthier, and R. Christen. 1995. *Roseobacter algicola* sp. nov., a new marine bacterium isolated from the phycosphere of the toxin-producing dinoflagellate *Prorocentrum lima*. *Int. J. Syst. Bacteriol.* **45**:290–296.
- Lane, D. J. 1991. 16S–23S rRNA sequencing, p. 125–175. *In* E. Stackebrandt and M. Goodfellow (ed.), *Nucleic acid techniques in bacterial systematics*. Wiley, Chichester, United Kingdom.
- Moore, L. R., G. Rocap, and S. W. Chisholm. 1998. Physiology and molecular phylogeny of coexisting *Prochlorococcus* ecotypes. *Nature* **393**:464–467.
- Olsen, G. J., H. Matsuda, R. Hagstrom, and R. Overbeek. 1994. fastDNAm1: a tool for construction of phylogenetic trees of DNA sequences using maximum likelihood. *Comput. Appl. Biosci.* **10**:41–48.
- Petursdottir, S. K., and J. K. Kristjansson. 1997. *Silicibacter lacuscaerulensis* gen. nov., sp. nov., a mesophilic moderately halophilic bacterium characteristic of the Blue Lagoon geothermal lake in Iceland. *Extremophiles* **1**:94–99.
- Prokic, I., F. Brummer, T. Brigge, H. D. Gortz, G. Gerdts, C. Schutt, M. Elbrachter, and W. E. G. Muller. 1998. Bacteria of the genus *Roseobacter* associated with the toxic dinoflagellate *Prorocentrum lima*. *Protist* **149**:347–357.
- Ramsing, N. B., M. J. Ferris, and D. M. Ward. 2000. Highly ordered vertical structure of *Synechococcus* populations within the one-millimeter-thick photic zone of a hot spring cyanobacterial mat. *Appl. Environ. Microbiol.* **66**:1038–1049.
- Rappé, M. S., K. Vergin, and S. J. Giovannoni. 2000. Phylogenetic comparisons of a coastal bacterioplankton community with its counterparts in open ocean and freshwater systems. *FEMS Microbiol. Ecol.* **33**:219–232.
- Rappé, M. S., S. A. Connon, K. L. Vergin, and S. J. Giovannoni. 2002. Cultivation of the ubiquitous SAR11 marine bacterioplankton clade. *Nature* **418**:630–633.
- Shiba, T. 1991. *Roseobacter litoralis* gen. nov. sp. nov., and *Roseobacter denitrificans* sp. nov., aerobic pink-pigmented bacteria which contain bacteriochlorophyll *a*. *Syst. Appl. Microbiol.* **14**:140–145.
- Sojka, G., H. H. Freeze, and H. Gest. 1970. Quantitative estimation of bacteriochlorophyll in situ. *Arch. Biochem. Biophys.* **136**:578–580.
- Sorokin, D. Y. 1995. *Sulfitobacter pontiacus* gen. nov., sp. nov., a new heterotrophic bacterium from the Black Sea, specialised on sulfite oxidation. *Microbiology* **64**:295–305.
- Suyama, T., T. Shigematsu, T. Suzuki, Y. Tokiwa, T. Kanagawa, K. V. Nagashima, and S. Hanada. 2002. Photosynthetic apparatus in *Roseateles depolymerans* 61A is transcriptionally induced by carbon limitation. *Appl. Environ. Microbiol.* **68**:1665–1673.
- Suzuki, M. T., M. S. Rappé, Z. W. Haimberger, H. Winfield, N. Adair, J. Strobel, and S. J. Giovannoni. 1997. Bacterial diversity among small-subunit rRNA gene clones and cellular isolates from the same seawater sample. *Appl. Environ. Microbiol.* **63**:983–989.
- Suzuki, T., Y. Muroga, M. Takahama, and Y. Nishimura. 1999. *Rosevivax halodurans* gen. nov., sp. nov., and *Rosevivax halotolerans* sp. nov., aerobic bacteriochlorophyll-containing bacteria isolated from a saline lake. *Int. J. Syst. Bacteriol.* **49**:629–634.
- Thompson, J. D., D. G. Higgins, and T. J. Gibson. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
- Uphoff, H., A. Felske, and I. Wagner-Döbler. 2001. The microbial diversity in picoplankton enrichment cultures: a molecular screening of marine isolates. *FEMS Microbiol. Ecol.* **35**:249–258.
- Wagner-Döbler, I., H. Rheims, A. Felske, R. Pukall, and B. Tindall. 2003.

- Jannaschia helgolandensis*, gen. nov., sp. nov., a novel abundant member of the marine *Roseobacter* clade from the North Sea. *Int. J. Syst. Evol. Microbiol.* **53**:731–738. (First published 4 October 2002; 10.1099/ijs.0.02377–0.)
40. **Yurkov, V., and J. T. Beatty.** 1998. Isolation of aerobic anoxygenic photosynthetic bacteria from black smoker plume waters of the Juan de Fuca Ridge in the Pacific Ocean. *Appl. Environ. Microbiol.* **64**:337–341.
41. **Yurkov, V. V., and J. T. Beatty.** 1998. Aerobic anoxygenic phototrophic bacteria. *Microbiol. Mol. Biol. Rev.* **62**:695–724.
42. **Yurkova, N., C. Rathgeber, J. Swiderski, E. Stackebrandt, J. T. Beatty, K. J. Hall, and V. Yurkov.** 2002. Diversity, distribution and physiology of the aerobic phototrophic bacteria in the mixolimnion of a meromictic lake. *FEMS Microbiol. Ecol.* **40**:191–204.
43. **Zubkov, M. V., B. M. Fuchs, S. D. Archer, R. P. Kiene, R. Amann, and P. H. Burkil.** 2001. Linking the composition of bacterioplankton to rapid turnover of dissolved dimethylsulphoniopropionate in an algal bloom in the North Sea. *Environ. Microbiol.* **3**:304–311.
44. **Zubkov, M. V., B. M. Fuchs, S. D. Archer, R. P. Kiene, R. Amann, and P. H. Burkil.** 2002. Rapid turnover of dissolved DMS and DMSP by defined bacterioplankton communities in the stratified euphotic zone of the North Sea. *Deep Sea Res. Part I* **49**:3017–3038.