

## Salinity Decreases Nitrite Reductase Gene Diversity in Denitrifying Bacteria of Wastewater Treatment Systems

Sachiko Yoshie,<sup>1</sup> Naohiro Noda,<sup>2</sup> Satoshi Tsuneda,<sup>1\*</sup> Akira Hirata,<sup>1</sup> and Yuhei Inamori<sup>3</sup>

Department of Chemical Engineering, Waseda University, Shinjuku-ku, Tokyo 169-8555,<sup>1</sup> National Institute of Advanced Industrial Science and Technology, Tsukuba, Ibaraki 305-8566,<sup>2</sup> and National Institute for Environmental Studies, Tsukuba, Ibaraki 305-0053,<sup>3</sup> Japan

Received 14 October 2003/Accepted 23 January 2004

**Investigation of the diversity of *nirK* and *nirS* in denitrifying bacteria revealed that salinity decreased the diversity in a nitrate-containing saline wastewater treatment system. The predominant *nirS* clone was related to *nirS* derived from marine bacteria, and the predominant *nirK* clone was related to *nirK* of the genus *Alcaligenes*.**

Nitrogen removal from wastewater is accomplished by bacterial denitrification. Some types of industrial wastewater, such as metallurgic wastewater, contain large amounts of nitrate and saline (11). Although we have developed a nitrogen removal system for saline wastewater, nitrite often accumulates, particularly under relatively high-saline conditions (32). Therefore, it is important to understand the ecology of nitrite-reducing bacterial communities to determine stable operational conditions for denitrification processes.

A few researchers and we have studied microbial communities in anaerobic reactors for the denitrification of saline wastewater (13, 31, 32) and used an approach based on taxonomic analysis of the 16S rRNA gene to identify all of the bacterial species within a community. These studies suggested that members of the gamma subdivision of the class *Proteobacteria* are important in such wastewater treatment systems. However, 16S rRNA gene-based approaches are unable to detect denitrifying bacteria particularly in such systems because bacterial groups possessing denitrifying abilities are phylogenetically diverse (33). Furthermore, although aerobic halophilic species have been reported (7, 29), the microbial ecology of moderately halophilic denitrifying bacteria has hardly been reported.

Recently, primer sets specific for functional genes involved in denitrification, namely, *nirK*, *nirS* (3, 9), and *nosZ* (22), have been developed. Thus, PCR-based approaches revealed denitrifying microbial communities in activated sludge (9, 27), marine sediment (4, 5, 15), and soil (17, 19) on the basis of *nirK* and *nirS* and in marine sediment on the basis of *nosZ* (23, 24). These approaches have contributed to the improvement of the complete or partial sequences of nitrite reductase genes and nitrous oxide reductase genes in the database. We expect that the information on these genes detected from various environments will contribute to further studies on the ecophysiology of denitrifying communities.

In this study, we focused on the microbial ecology of nitrite-reducing bacteria in two series of metallurgic wastewater treat-

ment systems (MWTs) with different fluidity conditions; one of the MWTs was composed of an anaerobic packed bed, and the other was composed of an anaerobic fluidized bed (31). The *nirK* and *nirS* heterogeneity in the anaerobic reactors of MWTs was investigated by cloning, sequencing, and phylogenetic analysis to determine the actual denitrifying bacterial community. Furthermore, the anaerobic packed bed and the fluidized bed were compared for *nirK* and *nirS* diversity to investigate the influence of fluidity conditions on the denitrifying microbial community.

**Sludge samples and isolates from MWTs.** Sludge samples were collected from two series of laboratory scale anaerobic-aerobic circulating bioreactors used as MWTs, as described previously (31). These systems consisted of an anaerobic reactor (2 liters) and an aerobic fluidized reactor (1 liter). The anaerobic reactor of run 1 was packed with sponge cubic medium without mixing. The anaerobic reactor of run 2 was a completely mixed fluidized bed in which polyvinyl alcohol particles coated with activated carbon were used as a carrier. The metallurgic wastewater used in this study was from a factory that recovers precious metals from industrial waste. The wastewater was diluted with tap water, and acetic acid (approximately 3,000 mg liter<sup>-1</sup>) was added as a carbon source for denitrification (C/N ratio = 1.5) before being supplied to the anaerobic reactors. The composition of the inlet was NO<sub>x</sub>-N (nitrate and nitrite) (1,800 to 2,500 mg liter<sup>-1</sup>), NH<sub>4</sub>-N (320 to 1,690 mg liter<sup>-1</sup>), and saline (14,000 to 32,000 mg liter<sup>-1</sup>). For DNA extraction, sludge samples adhering and not adhering to bed materials were collected and suspended in TE buffer (1 mM EDTA, 10 mM Tris-HCl [pH 8.0]) and stored at -20°C. Some bacteria were isolated from runs 1 and 2 of MWTs by using Trypticase soy agar (Becton Dickinson, Mountain View, Calif.) medium under aerobic conditions, and their denitrification activity was tested. The isolates were identified by partial 16S rRNA gene sequence analysis (31).

**DNA extraction, PCR amplification, and sequence analysis.** DNA extraction from sludge samples was performed by the method of Smalla (26), with slight modifications. The isolates were added to TE buffer and heated at 95°C for 5 min to extract DNA. The oligonucleotide primer pairs nirK1F-nirK5R and nirS1F-nirS6R (3) were used in PCR amplifications per-

\* Corresponding author. Mailing address: Department of Chemical Engineering, Waseda University, 3-4-1 Ohkubo, Shinjuku-ku, Tokyo 169-8555, Japan. Phone: 81-3-5286-3210. Fax: 81-3-3209-3680. E-mail: stsuneda@waseda.jp.

formed with a model 9700 thermal cycler (Applied Biosystems, Foster City, Calif.). The presence of PCR products, whose sizes were approximately 514 and 890 bp for *nirK* and *nirS*, respectively, was confirmed by running 8  $\mu$ l of the product on 2% agarose gels and then staining the agarose gels with ethidium bromide. The PCR products were purified by eluting the bands from the agarose gels with a Wizard SV gel and a PCR cleanup system (Promega Corp., Madison, Wis.). The eluted *nirK* or *nirS* PCR products were cloned with a ZERO Blunt TOPO PCR cloning kit (Invitrogen Corp., Carlsbad, Calif.) for *nirK* and a QIAGEN PCR cloning kit (QIAGEN, Hilden, Germany) for *nirS* in accordance with the manufacturer's instructions. The DNA insert was amplified and used as template DNA in a cycle sequencing reaction with a Big Dye Terminator cycle sequencing kit (Applied Biosystems) and a DYEnamic ET Terminator cycle sequencing kit (Amersham Biosciences, Freiburg, Germany) in accordance with the manufacturer's instructions. The *nirK* and *nirS* fragments were sequenced with an ABI PRISM 377 and ABI PRISM 3100-Avant DNA sequencing system (Applied Biosystems). Sequences with more than 97% similarity were considered to belong to the same operational taxonomic unit (OTU). The Shannon-Weaver index ( $H$ ) (25) was calculated with the formula  $H = -\sum(p_i)(\log_2 p_i)$ , where  $p_i$  is the proportion of each phylogenetic group to the total number of clones detected. Evenness (16) based on the Shannon-Weaver index was calculated with the formula  $E = H/\log_2 S$ . A database search was conducted with BLAST (1) from the DDBJ (DNA Data Bank of Japan). The sequences determined in this study and those retrieved from the databases were aligned by using CLUSTAL W (28). Phylogenetic trees were constructed with CLUSTAL W and Tree-View (18) by the neighbor-joining method (21).

**Phylogenetic relationship of *nirK* and *nirS*.** The phylogenetic tree based on the *nirS* sequence showed three major clusters (I to III) and 13 OTUs (A to M) (Fig. 1). The nucleotide sequence similarities between the *nirS* clones and those in the database ranged from 74.3 to 83.1%, and the deduced amino acid sequence similarities ranged from 73.3 to 88.9%. Most of the *nirS* clones obtained in this study exhibited less than 80% nucleotide sequence similarity to the *nirS* sequence in the database, suggesting that these *nirS* genes derived from novel denitrifying bacteria are indigenous to MWTSSs.

Cluster II included five OTUs of *nirS* clones from MWTSSs, and these *nirS* clones were associated with the *nirS* gene of *Pseudomonas stutzeri*. The fractions of the *nirS* clone affiliated with cluster II relative to all of the *nirS* clones detected were 32.3 and 52.1% in runs 1 and 2, respectively. OTUs B to E were most similar to the *nirS* gene of marine denitrifying isolate D7-6 (accession no. AJ248396) described by Braker et al. (5). The *nirS* genes of isolates R1-Apr-MIB-5 and R1-Dec-MIB-5, which were affiliated with the genus *Pseudomonas* (31), were included in OTUs B and C, respectively. OTUs D and E were unique to run 1, and their fractions were 15.6 and 8.9%, respectively. All of the *nirS* clones of MWTSSs did not belong to cluster IIb, including the *nirS* clones found in a marine environment by Braker et al. (5). For cluster IIIb, the most abundant *nirS* clones from run 1 were affiliated with OTU J and accounted for 53.3 and 18.1% of all of the *nirS* clones detected in runs 1 and 2, respectively. OTU J was the most similar to the *nirS* gene of marine denitrifying isolate B9-12 (accession no.

AJ248393) (98.6% similarity to *Halomonas variabilis*) found by Braker et al. (5). The *nirS* clones similar to the *nirS* gene of *Marinobacter* sp. were found, although their fraction was small (OTU H). The sequences of these clones were almost the same as the *nirS* sequence of isolate R1-Apr-MIB-4. Cluster IIIc consisted mainly of *nirS* clones of unknown denitrifying bacteria. OTU K was similar to the *nirS* gene of *Azoarcus* sp. at similarities of 13.3 and 13.8% to all of the *nirS* clones detected in runs 1 and 2, respectively. Clusters IIa and IIIb, whose fractions were large relative to all of the detected *nirS* clones, were related to the marine bacteria isolated by Braker et al. (5). The NaCl concentration of the influent appropriately diluted for MWTSSs was 1.5 to 3.5%, which is similar to that of seawater. Many *nirS* clones similar to the *nirS* gene of marine bacteria were detected from MWTSSs regardless of the differences in composition except that of saline, suggesting that a microbial community possessing *nirS* is affected by salinity. Although the physiology of moderately halophilic bacteria under anaerobic conditions has hardly been reported, *Halomonas elongata* (30), *Halomonas desiderata* (2), and *Bacillus halodenitrificans* (6) are well-known denitrifying bacteria. It was reported that *H. elongata* can reduce both nitrate and nitrite and that *B. halodenitrificans* is so unusually tolerant to nitrite as to grow in 0.58 M NaNO<sub>2</sub>.

It is said that *nirS* phylogenetic trees generally show the same clustering as 16S rRNA gene phylogenetic trees (5). The *nirS* genes of the bacteria isolated from MWTSSs, which are affiliated with *Pseudomonas* sp. (R1-MIB-Apr-5, Dec-5) and *Marinobacter* sp. (R1-MIB-Apr-4), branched with the same clusters including the same species isolated from a marine environment. However, no nitrite reductase genes have been detected from the isolated bacterium identified as *Halomonas* sp. by partial 16S rRNA gene sequence analysis (31), although *nirS* clone analysis detected clones similar to the *nirS* gene of *Halomonas* sp. (OTUs I and J). One possible explanation for these results is that the nitrite reductase genes of isolated *Halomonas* sp. might have some mismatch to the primers used in this study whereas uncultured bacteria had no mismatch to the primers used. Our previous study revealed that the group of bacteria belonging to the gamma subdivision of the class *Proteobacteria* including *Halomonas* sp. is dominant in MWTSSs (31). Simultaneous detection of the 16S rRNA and the functional gene will make it possible to combine the phylogenetic information of each gene.

For *nirK*, the number of clusters detected was lower than that for *nirS*; there were three major clusters and seven OTUs (Fig. 2). According to the BLAST search, the nucleotide sequence similarities between the *nirK* clones and those in the database ranged from 74.6 to 91.1% and the deduced amino acid sequence similarities ranged from 70.8 to 97.5%. In contrast to *nirS* clones, all of the *nirK* clones did not cluster with any marine clones. The fractions of clones including OTUs D, E, and F, which were not similar to any *nirK* sequence in the database, were high relative to all of the *nirK* genes detected, suggesting that those clones are inherent in MWTSSs. For both reactors, most of the *nirK* clones were affiliated with cluster II, including OTUs C, D, and E, similar to the *nirK* gene of *Alcaligenes xylosoxidans* (accession no. AF051831). In particular, the percentages of *nirK* clones affiliated with OTU C relative to all of the clones were 63.5 and 27.1% in runs 1 and 2,




FIG. 1. Phylogenetic tree constructed by the neighbor-joining method on the basis of partial *nirS* sequences cloned from run 1 (R1), run 2 (R2), or MWTS isolates (MIB). The *nirN* sequence of *Pseudomonas aeruginosa* was used as an outgroup to root the tree. A single sequence from each system was chosen as a representative of a cluster with more than 97% similarity. Bootstrap numbers indicate the value of 1,000 replicate trees supporting order (8), and values of less than 500 were omitted. Scale bar = 10% nucleotide substitution. The values in parentheses are the percent distributions of the clones within the *nirS* population.




FIG. 2. Phylogenetic tree constructed by the neighbor-joining method on the basis of partial *nirK* sequences cloned from run 1 (R1) or run 2 (R2) of MWTS. The *aniA* sequence of *Neisseria gonorrhoeae* was used as an outgroup to root the tree. A single sequence from each system was chosen as a representative of a cluster with more than 97% similarity. Bootstrap numbers indicate the value of 1,000 replicate trees supporting order (8), and values of less than 500 were omitted. Scale bar = 10% nucleotide substitution. The values in parentheses are the percent distributions of the clones within the *nirK* population.

respectively. The *nirK* gene of isolate R1-Apr-MIB-6, which clustered in the genus *Alcaligenes* in the phylogenetic tree on the basis of the partial 16S rRNA gene sequence, was included in OTU D. The fraction of cluster III, which consisted of the *nirK* of *Rhodobacter* sp. and *Azospirillum* sp., included the most abundant OTU in run 2 (F), and the fractions of cluster III were 5.7 and 41.6%, respectively. The *nirK* clones affiliated with cluster Ic, which were similar to the *nirK* gene of *Mesorhizobium* sp., were derived from run 2 only.

It was mentioned that great care should be taken to detect *nirK* fragments from environments with high salinity because the *nirK* primer set used in this study shows no sequence homology to the *nirK* gene of an archaean denitrifier, which has been recently sequenced (12). PCR amplification was carried out with primers specific for the archaean 16S rRNA gene to examine whether archaea exist in MWTSs, resulting in the lack of PCR products from these systems (data not shown). Therefore, there is no possibility of underestimating *nirK* diversity in MWTSs, which is consistent with the fact that halophilic archaea generally inhabit a hypersaline environment

with more than 2.5 M NaCl. Nevertheless, we should pay attention to the anticipated problem of *nirK* primer specificity, which is limited to the amplification of *nirK* similar to those that were referred to in the primer design, as mentioned in a previous study (5).

**Diversity of *nirK* and *nirS* from MWTSs.** Table 1 shows the phylotype richness (*S*), Shannon-Weaver diversity index (*H*), and evenness (*E*) of *nirK* and *nirS* in MWTSs. The Shannon-Weaver diversity indices of *nirS* and *nirK* were 1.96 and 1.22 for run 1 and 2.77 and 2.04 for run 2, respectively. Statistical analyses showed that the diversity level in run 2 was higher than that in run 1 for both *nirK* and *nirS*. Furthermore, the diversity of *nirS* was higher than that of *nirK* in both reactors. However, the diversities of *nirK* and *nirS* in MWTSs were lower than those in natural environments such as soil, in which the diversity indices of *nirK* and *nirS* are 3.55 and 5.27, respectively (19). This was probably due to the differences in salinity and carbon source between MWTSs and soil. For the latter reason, the influent of MWTSs included only acetic acid as a carbon source utilizable by bacteria, and any other organic

TABLE 1. Diversity analysis of *nirK* and *nirS* gene fragments from anaerobic packed bed and anaerobic fluidized bed of MWTs

Gene and reactor	<i>S</i> <sup>a</sup>	<i>H</i> <sup>b</sup>	<i>E</i> <sup>c</sup>
<i>nirS</i>			
Packed bed (run 1)	6	1.96	0.76
Fluidized bed (run 2)	10	2.77	0.84
<i>nirK</i>			
Packed bed (run 1)	5	1.22	0.52
Fluidized bed (run 2)	7	2.04	0.82

<sup>a</sup> Phylotype richness, *S* was the total number of OTUs in a sludge.

<sup>b</sup> The Shannon-Weaver diversity index (25) was calculated as follows:  $H = -\sum(p_i)(\log_2 p_i)$ , where  $p_i$  is the proportion of each phylogenetic groups to the number of all detected clones.

<sup>c</sup> Evenness (16) was calculated as follows from the Shannon-Weaver index:  $E = H/\log_2 S$ .

compounds originally included in metallurgic wastewater could not be biologically utilized. Therefore, it is presumed that salinity and small carbon source variations decrease the diversity of microbial communities. In fact, the microbial community in a methanol-fed denitrification process (13) was completely different from that in MWTs (31). Labbe et al. reported that members of the genus *Methylophaga* are predominant in a methanol-fed denitrification system for treating seawater (13). However, it was suggested that members of the genus *Halomonas* are dominant denitrifying bacteria in an acetate-fed saline wastewater treatment system in this study. Therefore, the diversity of nitrite reductase genes may also vary depending on the types of carbon sources used. Moreover, the *nirK* and *nirS* diversities in run 2 were higher than those in run 1 (Table 1). This difference in nitrite reductase gene diversity between runs 1 and 2 may arise from the differences between the biofilm community and the suspended microbial community caused by fluidity conditions. It was observed that most of the microorganisms in run 1 formed a biofilm in and on the medium, while those in run 2 were suspended regardless of the medium in the reactor. Other researchers reported that the difference in dissolved-oxygen concentration is one of the key factors in controlling the denitrifying community structure (15). In this study, although dissolved oxygen was almost 0 mg liter<sup>-1</sup> in run 2, the microbial community may have more chances to come in contact with oxygen on the water surface of the fluidized bed than on that of the packed bed, suggesting that the difference in the frequency of oxygen contact affects the diversity of nitrite reductase genes. These results indicate that fluidity conditions control denitrifying communities. Comparing the nitrogen removal efficiencies of runs 1 and 2, the removal efficiency of run 1 was found to be more stable than that of run 2, suggesting that a biofilm community is more suitable for MWTs than is a suspended microbial community. Furthermore, the expression of genes associated with denitrification is sensitive to the presence of oxygen (10). Therefore, the expression of the nitrite reductase gene may differ between a biofilm community and a suspended microbial community. More research studies that will introduce some methods of investigating bacterial activity, such as analysis of the expression of *nirK* and *nirS*, combination of fluorescence in situ hybridization and microautoradiography (14), and stable isotope probing analysis (20), will clarify the relationship between the ecology of an active denitrifying community and nitrogen re-

moval efficiency and lead to the improvement of nitrogen removal performance in MWTs.

Microbial community analysis based on nitrite reductase genes may enable the identification of key bacteria concerned with denitrification, which was the most significant process for MWTs. These results show that bacterial community analysis based on functional genes is important for a better understanding of microbial communities in wastewater treatment systems in addition to 16S rRNA gene analysis. Further comprehensive study of the relationships among qualitative and quantitative microbial community compositions, functions, and process stabilities will help in the design of advanced wastewater treatment systems or determination of appropriate operational conditions.

**Nucleotide sequence accession numbers.** The partial *nirK* and *nirS* sequences were submitted to the DDBJ database and assigned accession numbers AB118878 to AB118904.

This study was partly supported by Matsuda Sangyo Co., Ltd., which is thanked for its assistance.

#### REFERENCES

- Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment search tool. *J. Mol. Biol.* **215**:403–410.
- Berendes, F., G. Gottschalk, E. Heine-Dobbernack, E. R. B. Moore, and B. J. Tindall. 1996. *Halomonas desiderata* sp. nov., a new alkalophilic, halotolerant and denitrifying bacterium isolated from a municipal sewage works. *Syst. Appl. Microbiol.* **19**:158–167.
- Braker, G., A. Fesefeldt, and K. Witzel. 1998. Development of PCR primer systems for amplification of nitrite reductase genes (*nirK* and *nirS*) to detect denitrifying bacteria in environmental samples. *Appl. Environ. Microbiol.* **64**:3769–3775.
- Braker, G., H. L. Ayala-del-Río, A. H. Devol, A. Fesefeldt, and J. M. Tiedje. 2001. Community structure of denitrifiers, *Bacteria*, and *Archaea* along redox gradients in Pacific northwest marine sediments by terminal restriction fragment length polymorphism analysis of amplified nitrite reductase (*nirS*) and 16S rRNA genes. *Appl. Environ. Microbiol.* **67**:1893–1901.
- Braker, G., J. Zhou, L. Wu, A. H. Devol, and J. M. Tiedje. 2000. Nitrite reductase genes (*nirK* and *nirS*) as functional markers to investigate diversity of denitrifying bacteria in Pacific northwest marine sediment communities. *Appl. Environ. Microbiol.* **66**:2096–2104.
- Denariy, G., W. J. Payne, and J. Le Gall. 1989. A halophilic denitrifier, *Bacillus halodenitrificans* sp. nov. *Int. J. Syst. Bacteriol.* **39**:145–151.
- Dennis, P. P., and L. C. Shimmin. 1997. Evolutionary divergence and salinity-mediated selection in halophilic archaea. *Microbiol. Mol. Biol. Rev.* **61**:90–104.
- Felsenstein, J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**:783–791.
- Hallin, S., and P.-E. Lindgren. 1999. PCR detection of genes encoding nitrite reductase in denitrifying bacteria. *Appl. Environ. Microbiol.* **65**:1652–1657.
- Härtig, E., and W. G. Zumft. 1999. Kinetics of *nirS* expression (cytochrome *cd*<sub>1</sub> nitrite reductase) in *Pseudomonas stutzeri* during the transition from aerobic respiration to denitrification: evidence for a denitrification-specific nitrate- and nitrite-responsive regulatory system. *J. Bacteriol.* **181**:161–166.
- Hirata, A., Y. Nakamura, and S. Tsuneda. 2001. Nitrogen removal from industrial wastewater discharged from metal recovery processes. *Water Sci. Technol.* **44**:171–180.
- Ichiki, H., Y. Tanaka, K. Mochizuki, K. Yoshimatsu, T. Sakurai, and T. Fujiwara. 2001. Purification, characterization, and genetic analysis of Cu-containing dissimilatory nitrite reductase from a denitrifying halophilic archaeon, *Haloarcula marismortui*. *J. Bacteriol.* **183**:4149–4156.
- Labbe, N., P. Juteau, S. Parent, and R. Villemur. 2003. Bacterial diversity in a marine methanol-fed denitrification reactor at the Montreal biodome, Canada. *Microb. Ecol.* **46**:12–21.
- Lee, N., P. H. Nielsen, K. H. Andreasen, S. Juretschko, J. L. Nielsen, K.-H. Schleifer, and M. Wagner. 1999. Combination of fluorescent in situ hybridization and microautoradiography—a new tool for structure-function analyses in microbial ecology. *Appl. Environ. Microbiol.* **65**:1289–1297.
- Liu, X., S. M. Tiquia, G. Holguin, L. Wu, S. C. Nold, A. H. Devol, K. Luo, A. V. Palumbo, J. M. Tiedje, and J. Zhou. 2003. Molecular diversity of denitrifying genes in continental margin sediments within the oxygen-deficient zone off the Pacific coast of Mexico. *Appl. Environ. Microbiol.* **69**:3549–3560.
- Margalef, R. 1958. Information theory in ecology. *Gen. Syst.* **3**:36–71.
- Nogales, B., K. N. Timmis, D. B. Nedwell, and A. M. Osborn. 2002. Detection and diversity of expressed denitrification genes in estuarine sediment after

- reverse transcription-PCR amplification from mRNA. *Appl. Environ. Microbiol.* **68**:5017–5025.
18. **Page, R. D. M.** 1996. TreeView: an application to display phylogenetic trees on personal computers. *Comput. Appl. Biosci.* **12**:357–358.
  19. **Priemé, A., G. Braker, and J. M. Tiedje.** 2002. Diversity of nitrite reductase (*nirK* and *nirS*) gene fragments in forested upland and wetland soils. *Appl. Environ. Microbiol.* **68**:1893–1900.
  20. **Radajewski, S., P. Ineson, N. R. Parekh, and J. C. Murrell.** 2000. Stable-isotope probing as a tool in microbial ecology. *Nature* **403**:646–649.
  21. **Saitou, N., and M. Nei.** 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**:406–425.
  22. **Scala, D., and L. J. Kerkhof.** 1998. Nitrous oxide reductase (*nosZ*) gene-specific PCR primers for detection of denitrifiers and three *nosZ* genes from marine sediments. *FEMS Microbiol. Lett.* **162**:61–68.
  23. **Scala, D., and L. J. Kerkhof.** 1999. Diversity of nitrous oxide reductase (*nosZ*) genes in continental shelf sediments. *Appl. Environ. Microbiol.* **65**:1681–1687.
  24. **Scala, D., and L. J. Kerkhof.** 2000. Horizontal heterogeneity of denitrifying bacterial communities in marine sediments by terminal restriction fragment length polymorphism analysis. *Appl. Environ. Microbiol.* **66**:1980–1986.
  25. **Shannon, C. E., and W. Weaver.** 1963. The mathematical theory of communication. University of Illinois Press, Urbana.
  26. **Smalla, K.** 1995. Extraction of microbial DNA from sewage and manure slurries, p. 1.1.3. *In* A. D. L. Akkermans, J. D. van Elsas, and F. J. de Bruijn (ed.), *Molecular microbial ecology manual*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
  27. **Song, B., and B. B. Ward.** 2003. Nitrite reductase genes in halobenzoate degrading denitrifying bacteria. *FEMS Microbiol. Ecol.* **43**:349–357.
  28. **Thompson, J. D., D. G. Higgins, and T. J. Gibson.** 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
  29. **Ventosa, A., J. J. Nieto, and A. Oren.** 1998. Biology of moderately halophilic aerobic bacteria. *Microbiol. Mol. Biol. Rev.* **62**:504–544.
  30. **Vreeland, R. H., C. D. Litchfield, E. L. Martin, and E. Elliot.** 1980. *Halomonas elongata*, a new genus and species of extremely salt-tolerant bacteria. *Int. J. Syst. Bacteriol.* **30**:485–495.
  31. **Yoshie, S., N. Noda, T. Miyano, S. Tsuneda, A. Hirata, and Y. Inamori.** 2001. Microbial community analysis in the denitrifying process of saline-wastewater by denaturing gradient gel electrophoresis of PCR-amplified 16S rDNA and the cultivation method. *J. Biosci. Bioeng.* **92**:346–353.
  32. **Yoshie, S., N. Noda, T. Miyano, S. Tsuneda, A. Hirata, and Y. Inamori.** 2002. Characterization of microbial community in nitrogen removal process of metallurgic wastewater by PCR-DGGE. *Water Sci. Technol.* **46**:93–98.
  33. **Zumft, W. G.** 1997. Cell biology and molecular basis of denitrification. *Microbiol. Mol. Biol. Rev.* **61**:533–616.