

Gene Structure and Transcriptional Organization of the *dnaK* Operon of *Bifidobacterium breve* UCC 2003 and Application of the Operon in Bifidobacterial Tracing

Marco Ventura,^{1*} Ralf Zink,² Gerald F. Fitzgerald,¹ and Douwe van Sinderen¹

Alimentary Pharmabiotic Centre and Department of Microbiology, Bioscience Institute, National University of Ireland, Cork, Ireland,¹ and Department of Nutrition and Health, Cognis, Düsseldorf, Germany²

Received 30 June 2004/Accepted 16 August 2004

The incorporation and delivery of bifidobacterial strains as probiotic components in many food preparations expose these microorganisms to a multitude of environmental insults, including heat and osmotic stresses. We characterized the *dnaK* gene region of *Bifidobacterium breve* UCC 2003. Sequence analysis of the *dnaK* locus revealed four genes with the organization *dnaK-grpE-dnaJ*-ORF1, whose deduced protein products display significant similarity to corresponding chaperones found in other bacteria. Northern hybridization and real-time LightCycler PCR analysis revealed that the transcription of the *dnaK* operon was strongly induced by osmotic shock but was not induced significantly by heat stress. A 4.4-kb polycistronic mRNA, which represented the transcript of the complete *dnaK* gene region, was detected. Many other small transcripts, which were assumed to have resulted from intensive processing or degradation of this polycistronic mRNA, were identified. The transcription start site of the *dnaK* operon was determined by primer extension. Phylogenetic analysis of the available bifidobacterial *grpE* and *dnaK* genes suggested that the evolutionary development of these genes has been similar. The phylogeny derived from the various bifidobacterial *grpE* and *dnaK* sequences is consistent with that derived from 16S rRNA. The use of these genes in bifidobacterial species as an alternative or complement to the 16S rRNA gene marker provides sequence signatures that allow a high level of discrimination between closely related species of this genus.

Bifidobacteria are important components of the gastrointestinal microflora and have been the subject of growing scientific interest due to their suspected role in the maintenance of gastrointestinal health (21, 32, 36). The ability of these microorganisms to survive and adapt to changing environmental circumstances depends on their ability to resist harsh conditions encountered in their natural environment or during food manufacture in the case of probiotic strains (35). Under stressful situations, misfolded proteins may accumulate and/or aggregate, a process that may harm bacteria. Bacterial cells respond to stressful conditions by synthesizing molecular chaperones, which promote proper protein folding, thereby preventing the formation of insoluble protein aggregates. The major molecular chaperones include the DnaK complex (composed of DnaK, DnaJ, and GrpE) and the GroESL complexes (composed of GroES and GroEL) (8). Through cycles of protein binding and release, which are fuelled by ATP, the DnaK chaperone complex facilitates substrate folding to the native state (15).

In eubacteria, the *dnaK* operon has been investigated in detail in low-G+C-content bacteria, such as *Escherichia coli* and *Bacillus subtilis*. Cloning, sequencing, and transcriptional analyses of the *dnaK* operon in *B. subtilis* revealed four genes in the order *hrcA-grpE-dnaK-dnaJ*; the products of these genes constitute the *dnaK* chaperone machine (16). In these bacteria transcriptional regulation of the *dnaK* operon involves an in-

verted repeat element termed CIRCE (controlling inverted repeat of chaperone expression) located between the transcription initiation site and the *dnaK* coding region. This organization seems to be the basic and most common organization for *dnaK* operons in low-G+C-content gram-positive bacteria (29). A similar organization, *hrcA-grpE-dnaK* (without the *dnaJ* gene), was described for *Lactococcus lactis* and *Chlamydia* (28). A second variant of this organization, *grpE-dnaK-dnaJ*, was found in spirochetes, whereas in other low-G+C-content gram-negative bacteria (proteobacteria) the genes are organized into two transcriptional units; one operon contains *hrcA-grpE*, while the second operon contains *dnaK-dnaJ*. The first gene of the *B. subtilis* *dnaK* operon, *hrcA*, encodes a repressor that binds to the CIRCE element (30, 45, 46). Basic knowledge of the genetics of the *dnaK* locus in high-G+C-content gram-positive bacteria was obtained through analysis of the *Streptomyces coelicolor* genome sequence. In this microorganism the *dnaK* operon encodes four proteins, DnaK, GrpE, DnaJ, and HspR (5, 6), and the latter protein was shown to function as a repressor of the *S. coelicolor* *dnaK* operon. This gene inventory appears to be unique to *Actinobacteridae* (7). There is no sequence similarity between the Hsp-inverted repeat and HrcA-CIRCE systems, showing that *Streptomyces* has evolved a different strategy for controlling synthesis of the DnaK chaperone machine. Interestingly, a homologue of the *hrcA* gene, which in low-G+C-content bacteria is frequently part of the *dnaK* operon, is not found in this operon but was found to be part of an operon with a second copy of the *dnaJ* gene (12).

The genes constituting the *dnaK* operons are very well conserved across the archaeal, eubacterial, and eukaryal domains of life. Due to their ubiquitous distribution, functional preser-

* Corresponding author. Mailing address: Department of Microbiology, National University of Ireland, Western Road, Cork, Ireland. Phone: 353-21-4901366. Fax: 353-21-4903031. E-mail: m.ventura@ucc.ie.

TABLE 1. Strains used in this study and *grpE* and *dnaK* sequence accession numbers

Species or subspecies	Strain ^a	<i>grpE</i> accession no. ^b	<i>dnaK</i> accession no. ^b
<i>Bifidobacterium animalis</i>	ATCC 25527	AY642878	AY642871
<i>Bifidobacterium animalis</i>	LMG 18900	AY642890	AY642872
<i>Bifidobacterium lactis</i>	LMG 18906	AY642889	AY642873
<i>Bifidobacterium lactis</i>	LMG 11615	AY642887	AY642876
<i>Bifidobacterium lactis</i>	DSM 10140	AY667066	AY642912
<i>Bifidobacterium breve</i>	UCC 2003	AY642911	AY642911
<i>Bifidobacterium adolescentis</i>	JCM 1275	AY642885	AY642864
<i>Bifidobacterium longum</i>	NCC 2705	NC_004307	NC_004307
<i>Bifidobacterium infantis</i>	JCM 1222	AY642882	AY642867
<i>Bifidobacterium suis</i>	JCM 1269	AY642883	AY642869
<i>Bifidobacterium bifidum</i>	JCM 1255	AY642888	AY642868
<i>Bifidobacterium dentium</i>	JCM 1195	AY642886	AY642866
<i>Bifidobacterium catenulatum</i>	JCM 1194	AY642884	AY642875
<i>Bifidobacterium pseudocatenulatum</i>	JCM 1200	AY642881	AY642865
<i>Bifidobacterium angulatum</i>	JCM 7096	AY642880	AY642870
<i>Bifidobacterium magnum</i>	JCM 1218	AY642877	AY642863
<i>Bifidobacterium thermophilum</i>	JCM 1207	AY642879	AY642874
<i>Lactobacillus plantarum</i>	WCFSI	NC_004567	NC_004567
<i>Lactobacillus gasseri</i>	ATCC 33323	NZ_AAA002000001	NZ_AAA002000001
<i>Lactobacillus johnsonii</i>	NCC 533	NC_002662	NC_002662
<i>Lactococcus lactis</i> subsp. <i>lactis</i>	IL1403	NC_002662	NC_002662
<i>Streptococcus pyogenes</i>	MIGAS	NC_002737	NC_002737
<i>Enterococcus faecalis</i>	V583	NC_004668	NC_004668
<i>Clostridium acetobutylicum</i>	ATCC 824	NC_003030	NC_003030
<i>Mycobacterium bovis</i>	AF 2122/97	NC_002945	NC_002945
<i>Streptomyces coelicolor</i>	A3	NC_003888	NC_003888
<i>Corynebacterium glutamicum</i>	ATCC 13032	NC_003450	NC_003450
<i>Listeria monocytogenes</i>	EGD	NC_003210	NC_003210
<i>Escherichia coli</i>	K12	NC_000913	NC_000913

^a ATCC, American Type Culture Collection; DSM, Deutsche Sammlung von Mikroorganismen; JCM, Japanese Collection of Microorganisms; LMG, Bacteria Collection, Universiteit Gent.

^b For the strains whose genome sequences are available, the *grpE* and *dnaK* sequences were retrieved from the complete bacterial genome, and the accession numbers are indicated.

vation, and sequence conservation, the *dnaK* operon genes are considered valuable molecular markers for bacterial phylogenetic investigations. They have been successfully used for identification of *Mycobacterium* (31), *Staphylococcus* (1), *Streptococcus* (1), *Legionella* (22), and *Ruminococcus* (2) species. With the introduction of polyphasic taxonomy (34), many molecular markers that are alternatives to the 16S rRNA gene sequences have been described. Recently, alternative molecules, such as the gene encoding elongation factor Tu (39, 40), ATPase subunits (38), the *recA* gene (19, 40), and *groEL* (37), have been used to examine phylogenetic relationships between bifidobacterial species. The use of these molecular markers corroborates and completes the evolutionary history of bifidobacterial species. However, other molecular markers, which provide finer taxonomic detail to distinguish closely related species (e.g., *Bifidobacterium lactis* and *Bifidobacterium animalis* or *Bifidobacterium catenulatum* and *Bifidobacterium pseudocatenulatum*), are required.

To date, very little is known about the *dnaK* operon of bifidobacteria (26). In this report we describe the *dnaK* operon of *Bifidobacterium breve* UCC 2003. We tested the heat and osmotic inducibility of this operon by performing Northern blot hybridization analyses, primer extension experiments, and real-time PCR. Moreover, we describe below partial sequencing of the genes coding for the GrpE and DnaK chaperones from 14 bifidobacterial species in order to provide new molec-

ular markers for investigation of the phylogeny of the genus *Bifidobacterium*.

MATERIALS AND METHODS

Bacterial strains and culture conditions. All *Bifidobacterium* strains were grown anaerobically in MRS (Difco, Detroit, Mich.) supplemented with 0.05% L-cysteine-HCl and incubated at 37°C for 16 h. The bacterial strains used are summarized in Table 1.

DNA amplification and cloning of the *dnaK* and *grpE* genes. PCR was used to amplify the *dnaK* gene of all *Bifidobacterium* strains investigated. A DNA fragment corresponding to the *dnaK* gene was amplified by using oligonucleotides DNAK-1 (5'-GTCAGGCTGTGACCAAC-3') and DNAK-2 (5'-GGTCATTCCTCGAAGTG-3'), while the *grpE*-containing DNA fragment of *Bifidobacterium* strains was amplified by employing oligonucleotides GRE-UNI (5'-GAGTTCA TCAACTACCGCA-3') and GRE-REV (5'-CTTCCACCAGGTGTC-3'). Each PCR mixture (50 μ l) contained 20 mM Tris-HCl, 50 mM KCl, each deoxynucleoside triphosphate at a concentration of 200 μ M, 50 pmol of each primer, 1.5 mM MgCl₂, and 1 U of *Taq* DNA polymerase (Gibco BRL, Paisley, United Kingdom). Each PCR cycling profile consisted of an initial denaturation step of 3 min at 95°C, followed by amplification for 30 cycles as follows: denaturation for 30 s at 95°C, annealing for 30 s at 50°C, and extension for 1 min at 72°C. The PCR was completed with an elongation phase consisting of 10 min at 72°C. The resulting amplicons were separated on a 1.5% agarose gel, and this was followed by ethidium bromide staining. PCR fragments were purified with a PCR purification spin kit (Genomed, Lohne, Germany) and were subsequently sequenced.

The upstream region of the *dnaK* operon of *B. lactis* DSM 10140 was determined by inverse PCR (24). One microgram of DSM 10140 chromosomal DNA was digested with restriction endonuclease EcoRI or ClaI, and the restriction fragments were self-ligated and amplified by using primers DNAK-1-inv (5'-G TTGGTCACAGCCTGAC-3') and DNAK-2-inv (5'-CACTTCGAGGAAATG

ACC-3') as described by Sambrook and Russell (24). The inverse PCR products obtained were used for direct sequencing with synthetic oligonucleotides.

DNA sequencing and phylogenetic study. Nucleotide sequencing of both strands of PCR amplicons was performed by MWG-Biotech AG (Ebersberg, Germany) with primers DNAK-1, DNAK-2, GRE-UNI, and GRE-REV. The *dnaK* and *grpE* partial gene sequences of all *Bifidobacterium* strains examined in this study were used for comparison. Sequence data assembly and analysis were performed by using the DNASTAR software (version 5.05; DNASTAR, Madison, Wis.). Sequence alignment was performed by using the MultiAlign and Clustal W programs (33). Phylogenetic calculations, including distance calculations and generation of phylogenetic trees, were performed by using the PHYLIP package (9). Trees were calculated by the neighbor-joining method as implemented in the neighbor module of PHYLIP. Also, dendrograms of gene sequences were constructed by using the Clustal X program and were visualized with the TreeView program.

RNA isolation and Northern blot analysis. *B. breve* UCC 2003 cells were grown to an optical density at 600 nm of 0.6. Heat stress was applied by transferring the culture to 43°C, while osmotic stress was applied by addition of prewarmed medium containing 5 M NaCl to obtain a final NaCl concentration of 0.3, 0.5, 0.7, or 1.5 M. At various times, 30 ml of culture was collected and briefly centrifuged to harvest cells. Total RNA was isolated by using macaloid acid (37) and was treated with DNase (Roche, Lewes, East Sussex, United Kingdom). Fifteen micrograms of RNA was subjected to electrophoresis on a 1.5% agarose–3.3% formaldehyde denaturing gel, transferred to a Zeta-Probe blotting membrane (Bio-Rad, Hemel Hempstead, Hertfordshire, United Kingdom) as described by Sambrook and Russell (24), and fixed by UV cross-linking by using a Stratalinker 1800 (Stratagene, La Jolla, Calif.). PCR amplicons obtained by using primer combinations that were designed to target each of the genes of the *B. breve* UCC 2003 *dnaK* operon were radiolabeled (24). Prehybridization and hybridization were carried out at 65°C in 0.5 M NaHPO₄ (pH 7.2)–1.0 mM EDTA–7.0% sodium dodecyl sulfate (SDS). Following 18 h of hybridization, the membrane was rinsed twice for 30 min at 65°C in 0.1 M NaHPO₄ (pH 7.2)–1.0 mM EDTA–1% SDS and twice for 30 min at 65°C in 0.1 M NaHPO₄ (pH 7.2)–1.0 mM EDTA–0.1% SDS and then exposed to X-OMAT autoradiography film (Eastman Kodak, Rochester, N.Y.).

cDNA synthesis and quantification of specific transcripts by LightCycler PCR. Five micrograms of mRNA was treated with DNase (Roche) and used as a template in a 100- μ l reaction mixture containing 20 ng of random primers, each deoxyribonucleoside triphosphate at a concentration of 0.125 mM, and SuperScript enzyme (Invitrogen, Paisley, United Kingdom) used according to the manufacturer's instructions to produce cDNA. The cDNA generated was used as a template for reverse transcription (RT)-PCRs to determine the level of transcription of *grpE* and ORF1 as representative genes of the *dnaK* operon.

In order to accurately determine and compare the transcript levels of any differentially regulated gene, it was necessary to normalize cDNA levels from the individual times. To do this, by using primers 16SRT2 (5'-GTTCACGGGTTCCGTG-3') and 16SRT3 (5'-GAGCTGACGACGACCATG-3') and previously described parameters (23), the constitutively expressed 16S rRNA gene was standardized for each time by adjusting the volume of cDNA added prior to PCR until the crossing point values were identical for all samples. Subsequently, by using the standardized volumes of cDNA, the relative expression of the target genes was determined as outlined below. To prepare quantification standards for the LightCycler PCR, a 285-bp region of the *grpE* gene was amplified by using primers GRE-UNI and GRE-REV, and a 300-bp region of ORF1 was amplified by using primers TRA-1 (5'-GACAAAATGCGTGCGATG-3') and TRA-2 (5'-CAACCAGATGAAGCAGTT-3'). The PCR products were purified with a PCR purification spin kit (Genomed) and were subsequently quantified with a GenQuant spectrophotometer (Pharmacia). The copy number was calculated, and appropriate dilutions containing from 10¹ to 10⁸ copies of the standards were prepared in elution buffer (QIAGEN), divided into aliquots, and stored at 4°C. A Faststart SYBR Green master mixture kit (Roche) was used for all LightCycler PCRs. By using the standards generated as described above, the conditions were optimized for the real-time PCR as recommended by the manufacturer. This included titrating the MgCl₂ and primer concentrations, as well as optimizing the annealing temperatures and extension times. The final optimized conditions used in all subsequent analyses were 2 μ l of either the standard dilution or standardized dilution of each RT reaction mixture as a template, 2 mM MgCl₂, 2 μ l of the Faststart SYBR Green master mixture (Roche), and each relevant primer at a concentration of 0.5 μ M in a 20- μ l (total volume) reaction mixture. The PCR parameters for amplification included denaturation for one cycle at 95°C for 8 min and 40 cycles of 95°C for 5 s, 50°C for 5 s, and 72°C for 8 s. A melting curve analysis was performed at 65 to 90°C. The melting curve for each bacterial sample was generated manually to determine the melting temperature

(T_m). The T_m was the peak of the curve determined by measuring the derivative of fluorescence with respect to temperature ($-dF/dT$). By using the manual T_m function of the LightCycler software, the T_m was defined as the temperature at which the cursor identified the highest point on the melting curve (the $-dF/dT$ curve). If the highest point was a plateau, the midpoint of the plateau was taken to be the T_m .

All LightCycler assay data analyses were carried out by using the fit points option of the Roche LightCycler software (version 3.53). The ability to quantify unknown targets was evaluated by using eight separate DNA samples containing various amounts of template (range, 10¹ to 10⁸ copies) that were analyzed by using RT-PCR in conjunction with known standards in this range. The number of copies of each sample transcript was then determined with the LightCycler software. The standard dilutions were used to generate crossing point values, which were used to quantify the transcript levels of *grpE* and ORF1. By fitting the crossing point values (beginning of the exponential phase) of the known standards to those of dilutions of the unknown real-time reaction mixtures, the number of gene transcripts for each target was calculated. Each experiment was performed in triplicate, and the standard deviations were calculated to ensure statistical integrity.

Primer extension analysis. The 5' ends of the RNA transcripts were determined as described in a previous study (38). The synthetic oligonucleotides used were as follows: dnaK-prom (5'-GTGGTGCGCATGCCCTC-3'), grpE-prom (5'-CTCCTTCATGTGAAGGAGCTG-3'), dnaJ-prom (5'-CTCGGCTTCCTGGTCTTG-3'), and ORF1-prom (5'-GTACTGACGGAGGGTCTG-3'), located at positions 90 to 106, 148 to 162, 125 to 143, and 162 to 179, respectively, in the respective nucleotide sequences.

PCR identification of *B. lactis*. Amplification reactions were performed in 50- μ l (total volume) solutions containing 10 mM Tris-HCl, 50 mM KCl, 1.5 mM MgCl₂, each deoxynucleoside triphosphate (Gibco BRL) at a concentration of 200 μ M, 10 pmol of gro-1 (37), 10 pmol of gro-2 (37), 50 pmol of *B. lactis*-specific primer b-dnaK1 (5'-CACAGATCCTCATGAAGCTG-3'), 50 pmol of *B. lactis*-specific primer b-dnaK2 (5'-CGACGTAGCCTGCACCTGG-3'), 25 ng of template DNA, and 2.5 U of *Taq* DNA polymerase (Gibco BRL). Amplifications were performed with a DNA thermocycler (Primus) with the following temperature profile: one cycle of 95°C for 5 min; 30 cycles of 95°C for 30 s, 50°C for 1 min, and 72°C for 1 min; and one cycle of 72°C for 7 min. For routine identification, cells were lysed by using a rapid DNA extraction protocol (43) and were used as direct PCR templates. PCR amplicons were analyzed by 1.5% (wt/vol) agarose gel electrophoresis in Tris-acetate-EDTA buffer at a constant voltage of 7 V/cm, were visualized with ethidium bromide (0.5 μ g/ml), and were photographed under UV light at 260 nm.

Nucleotide sequence accession numbers. The GenBank accession numbers for partial *grpE* and *dnaK* gene sequences generated in this study are shown in Table 1. The nucleotide sequence data for the *dnaK* operon of *B. breve* UCC 2003 have been deposited in the GenBank database under accession number AY642911.

RESULTS

Genetic organization of the *dnaK* operon. Analysis of the *B. breve* UCC 2003 genome sequence (S. Leahy, J. A. Moreno Munoz, M. O'Connell-Motherway, D. Higgins, G. F. Fitzgerald, and D. van Sinderen, unpublished data) allowed us to identify a 4.4-kb DNA fragment encompassing the *dnaK* operon, which is represented by four open reading frames. The gene order of the *dnaK* operon was *dnaK*-*grpE*-*dnaJ*-ORF1, which is identical to that observed in other high-G+C-content gram-positive bacteria (28). In fact, a comparison with other *Actinobacteridae* (*Bifidobacterium longum*, *S. coelicolor*, *Corynebacterium glutamicum*, and *Mycobacterium bovis*) revealed that the genetic organizations of the *dnaK* operon are identical (Fig. 1). The gene inventory of the *dnaK* operon in unrelated taxa (e.g., *Lactobacillus* and *Escherichia*) was shown to be different, which is in accordance with their distant phylogenetic positions (27).

A protein comparison (Fig. 1) showed that the proteins most similar to the *B. breve* DnaK, GrpE, and DnaJ chaperones were proteins of *B. longum*. Notably, levels of identity of $\geq 24\%$ were observed for these three proteins with the corresponding




FIG. 1. Comparison of the *dnaK* operon in *B. breve* UCC 2003 with corresponding loci in various other bacteria. Each arrow indicates an open reading frame, and the length of the arrow is proportional to the length of the predicted open reading frame. Corresponding genes are indicated by the same color. The putative function of the protein is indicated above each arrow. Genes exhibiting $\geq 55\%$ amino acid similarity are linked by light blue shading, and genes exhibiting $\leq 54\%$ amino acid similarity are linked by light violet shading. Levels of amino acid identity, expressed as percentages, are indicated. Sol. bind. protein, soluble binding protein.

proteins of *Lactobacillus johnsonii*. The DnaK chaperone is the most conserved protein encoded by the operon in high- and low-G+C-content bacteria. Furthermore, a comparison of the *B. breve* *dnaK*, *grpE*, and *dnaJ* products with proteins deposited in the publicly available databases showed highly significant sequence similarities for the complete DnaK, GrpE, chaperones of other high-G+C-content gram-positive bacteria (e.g., *Mycobacterium*, *Streptomyces*, *Corynebacterium*, and *Propionibacterium*). In contrast, the DnaJ chaperone homologue of *B. breve* exhibited considerable similarity to the DnaJ chaperone found in low-G+C-content bacteria (*Bordellia*, *Nostoc*, and *Staphylococcus*).

ORF1, located downstream of *dnaJ*, is a gene which is predicted to encode a protein containing 195 amino acids and having a calculated mass of 21.8 kDa that displays significant similarity to other predicted heat-controlled transcriptional regulators belonging to the Mer family. These regulators include those identified for *Mycobacterium leprae* (level of identity, 43%), *Aquifex aeolicus* (level of identity, 39%), and *Deino-*

coccus radiodurans (level of identity, 32%). Interestingly, the N-terminal regions of these proteins are predicted to contain a classical helix-turn-helix motif, which is assumed to be involved in DNA recognition and DNA binding processes (4). Thus, this finding suggests that the ORF1 product is also involved in such interactions.

The predicted *B. breve* UCC 2003 DnaK protein contains the consensus HSP70 domains, which have been described as being involved in substrate binding and release (pfam0012). The predicted DnaJ chaperone of *B. breve* contains the J domain, which is highly conserved in DnaJ homologs and which is involved in protein interaction and folding (8).

The DNA regions surrounding the *dnaK* operon in *B. breve* do not display homology to the regions flanking the *dnaK* loci of other available bifidobacteria, indicating that the observed conservation of gene order in the two sequenced bifidobacteria appears to be restricted to this operon (data not shown).

Interestingly, the initiation methionine codon of the *grpE* gene overlaps the TGA termination codon of the *dnaK* gene, in

a -1 reading frame, suggesting that the two genes are translationally coupled. The *dnaJ* gene and ORF1, on the other hand, are separated from the surrounding genes by 232- and 159-bp intergenic regions, respectively.

Analysis of the DNA sequences of the *dnaK* operon revealed several inverted repeats which could form potential stem-loop structures and thereby serve as cleavage and/or transcription terminator sites. These regions are shown in Fig. 2a as stem-loop structures above the genes and are designated 1 through 12. Stem-loop structures 1 and 12 may serve as rho-independent transcription terminators (since they occur outside the structural operon), whereas stem-loop structure 11 may represent a potential processing site while stem-loop structures 2 to 10 may be degradation sites (as they are situated within structural genes).

Transcription mapping of the *dnaK* operon. In order to determine transcription patterns for the *dnaK* operon, total RNA extracted from *B. breve* UCC 2003 cells grown under unstressed and stressed conditions was hybridized with DNA probes targeting the four genes of the *dnaK* operon. The results indicated that a long transcript is synthesized, which is subject to extensive degradation, as indicated by the presence of a conspicuous smear in all blots (Fig. 2). A probe encompassing the *dnaK* gene revealed a smear of transcripts, the largest of which had an estimated size of 4.4 kb. This transcription pattern was also observed for all other genes found in the *dnaK* operon, suggesting that the four genes are transcribed as a single polycistronic mRNA (Fig. 2a and b).

Interestingly, when an ORF1-specific probe was used, a smaller hybridization signal corresponding to a 580-bp transcript was observed in addition to the smear of transcripts (Fig. 2b).

Northern blot analysis provided evidence that the ORF1, *dnaJ*, *grpE*, and *dnaK* genes constitute an operon since they appear to be transcribed as a polycistronic mRNA. Further proof for the existence of a polycistronic transcript was provided by RT-PCR experiments (Fig. 2d). cDNA templates were obtained by RT of total *B. breve* UCC 2003 mRNA isolated under heat stress conditions (Fig. 2d, lanes 2, 7, 12, and 15) or osmotic stress conditions (Fig. 2d, lanes 1, 6, 11, and 14). Each cDNA was used as the template in subsequent PCRs with different combinations of primer pairs spanning the *dnaK*, *grpE*, *dnaJ*, and ORF1 genes (Fig. 2a). RNA, which was prepared in the presence of heat and osmotic stresses, produced RT-PCR amplicons that comigrated with the products obtained when chromosomal DNA of *B. breve* UCC 2003 was used as a positive control (Fig. 2d, lanes 3, 8, 13, and 16). The RT-PCR products were also shown to be dependent on RNA in the cDNA preparation; in fact, no products were observed with mock cDNA prepared in the absence of reverse transcriptase (Fig. 2d, lanes 4, 9, and 17). Furthermore, no RT-PCR product was observed in the absence of any template (Fig. 2d, lanes 5, 10, and 18).

Northern blot analysis was performed by using RNA collected from cells following heat shock and osmotic stress. The hybridization signals exhibited the maximum level of intensity for RNA samples collected following exposure for 100 min to a medium containing 0.7 M NaCl (Fig. 2c, lanes 7). On the other hand, after exposure to heat stress, transcription of the

dnaK operon of *B. breve* UCC 2003 appeared to be only weakly activated (Fig. 2b).

Induction of *dnaK* operon mRNA in response to environmental stresses. To further investigate the extent to which expression of the *B. breve* *dnaK* operon is modulated by an osmotic or temperature upshift, quantitative real-time PCR assays were performed. PCR products, obtained by using cDNA samples from different times as templates, were amplified based on the *grpE* and ORF1 sequences. The purified PCR products were used as standards, and dilutions of each standard were used to generate a standard curve. The sensitivity of the LightCycler PCR was estimated to be 10 copies of the *grpE* and ORF1 sequences, which was the cutoff point for accurate detection of DNA. Dilutions corresponding to copy numbers ranging from 10^1 to 10^8 were used in all LightCycler PCRs as standards. Using the LightCycler software, we determined the numbers of copies of the *grpE* and ORF1 transcripts.

In a previous study the heat shock response in *B. breve* UCC 2003 was evaluated (37). A slot blot hybridization procedure was used to test total RNA, which was isolated from *B. breve* UCC 2003 cultures grown for different lengths of time at temperatures ranging from 37 to 50°C. Based on the intensity of the hybridization signal, the most effective temperature for inducing heat shock gene expression was 43°C (37). In contrast to what was previously observed for the *dnaK* gene in *Bifidobacterium adolescentis* and *B. longum* (26), a relatively low level of heat induction of *grpE* and ORF1 gene transcription was observed. A 1.4-fold increase in the copy number of *grpE* transcripts was observed when cultures were shifted from 37 to 43°C for 150 min (Fig. 3a). Similarly, a 1.5-fold upshift of the ORF1 transcripts was observed upon heat stress (data not shown). Conversely, the levels of *grpE* and ORF1 mRNAs dramatically increased following osmotic shock. In fact, an osmotic upshift to 0.7 M NaCl in the media led to 15- and 14-fold increases in the *grpE* and ORF1 mRNAs, respectively (Fig. 3b and data not shown).

Identification of the transcription start site of the *dnaK* operon. To determine the transcriptional start point of the *dnaK* operon, primer extension analyses were performed with RNA extracted from *B. breve* cells grown at 37°C or subjected to heat or osmotic shock (Fig. 4). A major extension product was identified 75 nucleotides 5' to the translational start site of the *dnaK* gene (Fig. 4a). The transcription start sites were in the same positions at 37°C, at 43°C, and under osmotic shock conditions (Fig. 4a). The analysis of the putative promoter region of *dnaK* revealed a potential promoter-like sequence resembling the consensus -10 hexamer and -35 box (Fig. 4c) (36, 37). A specific primer extension product was also mapped within the intercistronic region between ORF1 and *dnaJ* (Fig. 4b). However, screening for promoter consensus sequences of ORF1 did not reveal any typical *E. coli* -10 and -35 sequences, and computer analysis predicted a high degree of RNA secondary structures in this region. Therefore, we cannot rule out the possibility that the mapped 5' end actually represents a cleavage site of the polycistronic mRNA.

Primer extension experiments were also performed to investigate the existence of other transcription initiation sites upstream of the assumed start codons of the *dnaJ* and *grpE* genes, but all these attempts failed to generate extension products. This suggests that these genes do not possess their own pro-




FIG. 2. Transcriptional organization of the *B. breve* UCC 2003 *dnaK* operon. (a) Schematic diagram of the *B. breve* UCC 2003 *dnaK* operon. Hairpin symbols indicate predicted secondary structures. Transcripts are indicated by arrows, which point toward the 3' end of the mRNA. (b) Northern blot analysis of *B. breve* UCC 2003 *dnaK* operon performed by using total mRNA isolated from cultures grown at 37°C (lanes 1 to 5) and 43°C (lanes 6 to 9) for the following lengths of time after a temperature shift: none (beginning of the experiments) (lanes 1), 25 min (lanes 2 and 6), 50 min (lanes 3 and 7), 100 min (lanes 4 and 8), and 150 min (lanes 5 and 9). The probes used are indicated below the panels. (c) Northern blot analysis of *B. breve* UCC 2003 *dnaK* operon performed by using total mRNA isolated from cultures grown in the absence of NaCl (lanes 1)

motor regions, which is consistent with the presence of a polycistronic 4.4-kb transcript.

Comparison of the upstream regulatory regions of *dnaK* operons from *Actinobacteridae*. The upstream regions of the *dnaK* genes of two bifidobacterial strains and one *Streptomyces* strain were aligned in an attempt to identify putative regulated elements (Fig. 4c). For completeness, we determined by inverse PCR the putative promoter region of the *dnaK* gene for a more distantly related *Bifidobacterium* species (*B. lactis*). As Fig. 4c shows, a consensus promoter sequence could be deduced from the three sequences. The putative -10 region was conserved in all bifidobacterial sequences, while the putative -35 box was less conserved. Moreover, a clear consensus inverted repeat sequence (TGAGN₉CTCA), which was designated IR, was identified. Notably, the promoter region of the *dnaK* gene in *B. breve* and *B. longum* contained two copies of this IR sequence (IR1 and IR2), whereas *B. lactis* possessed only a single copy. Interestingly, this inverted repeat resembled a regulatory structure designated HAIR (CTTGAGTN₇ACTCAAG) (5), which has been demonstrated to be the operator sequence of the heat shock protein regulator (HspR) in high-G+C-content gram-positive bacteria (11). Preliminary screening of the publicly available *B. longum* NCC 2705 genome sequence for the consensus sequence (TGAGN₉CTCA) revealed the presence of this inverted repeat in the intergenic regions of a classical stress-induced gene, such as *clpB* (Fig. 4d). Furthermore, this consensus sequence was identified in the promoter regions of the *trmD* and *malQ1* genes, which are predicted to encode a methyltransferase and a glucanotransferase, respectively (25).

Phylogenetic analysis. The *grpE* and *dnaK* DNA sequences of a number of selected bacterial species were aligned and compared. Two highly conserved regions for each gene were identified, and a pair of primers (GRE-UNI and GRE-REV) amplifying a 285-bp region was designed for the *grpE* gene. Furthermore, another primer pair (DNAK-1 and DNAK-2) was designed to amplify a 535-bp region. These primers allowed amplification of *grpE* and *dnaK* sequences from 14 *Bifidobacterium* species. The alignments of *grpE* DNA and *dnaK* sequences were used to generate two phylogenetic trees by the neighbor-joining method (Fig. 5). The data were supported by the bootstrap values obtained. For completeness we included in the analysis the *grpE* DNA and *dnaK* sequences of other strains belonging to different genera representing the lactic acid bacterial group (e.g., *Lactococcus*, *Streptococcus*, *Lactobacillus*, and *Enterococcus*) and other gram-positive and gram-negative bacteria with high and low G+C contents, including some *Actinobacteridae* (e.g., *Corynebacterium*, *Streptomyces*, and *Mycobacterium*). These trees showed two major clusters representing the low-G+C-content gram-positive bacteria (the

genera *Lactobacillus*, *Lactococcus*, *Streptococcus*, and *Enterococcus*) and the high-G+C-content gram-positive bacteria (the genus *Bifidobacterium*).

A comparison of the phylogenetic trees based on the *grpE* and *dnaK* sequences revealed very similar phylogenetic topologies (Fig. 5).

A phylogenetic tree was also constructed on the basis of the 16S rRNA gene sequences available in databases. The phylogenetic positions of bacteria based on 16S rRNA gene sequences were generally in agreement with *grpE*- and *dnaK*-based phylogenies (data not shown). Interestingly, closely related strains which had nearly identical 16S rRNA gene sequences (e.g., *B. animalis* and *B. lactis*, *B. catenulatum* and *B. pseudocatenulatum*, or *B. longum* and *Bifidobacterium infantis*) clearly branched separately in the *grpE* and *dnaK* sequence-based tree (Fig. 5).

The similarity between the partial *grpE* DNA sequences of bifidobacterial species was less than 96.6% in all cases except *B. longum*-*Bifidobacterium suis* and *B. infantis* (99.8%). The sequence divergence in *grpE* was 2.7 times greater than that in the 16S rRNA gene, indicating the greater variance of *grpE*. The *dnaK* sequences were less divergent than those of *grpE* by a factor of 1.4. The relationship between the pairwise dissimilarities of *grpE* and other genes was linear, which suggests that the degree of sequence divergence was proportional to the time needed for speciation. In addition, the pairwise distances between the *dnaK* and 16S rRNA genes exhibited a close linear correlation (data not shown). Unfortunately, the *dnaK* gene failed to act as a discriminating molecular marker for typing purposes. In fact, the phylogenetic distances between *B. lactis* strains, as well as those between *B. animalis* strains, are virtually zero.

Most of the base substitutions in the *grpE* and *dnaK* genes were synonymous; i.e., they did not result in amino acid changes but constituted nucleotide signatures that may be used for tracing bifidobacterial species.

Specific identification of bifidobacteria by melting point analysis and by using species-specific primers. Following amplification of 285-bp *grpE* sequences of the individual bifidobacterial strains, a melting curve of the amplicon was generated by continuously monitoring the fluorescence while the temperature was raised from 82 to 95°C. The T_m s for the partial *grpE* gene sequences remained within a narrow range, but the T_m was different for each of the species examined. The only exceptions were *B. longum* and *B. infantis*, which showed overlapping melting curves. Characteristic melting curves for closely related taxa, like *B. catenulatum* and *B. pseudocatenulatum* or *B. longum* and *B. suis*, as well as for *B. lactis* and *B. animalis*, are shown in Fig. 6a and b. The T_m of *B. lactis* DSM 10140 was 92°C, which was unequivocally higher than that of *B.*

and under hyperosmotic conditions (lanes 2 to 9) with the following NaCl concentrations and for the following times: 0.3 M NaCl for 50 min (lane 2), 0.3 M NaCl for 100 min (lane 3), 0.5 M NaCl for 50 min (lane 4), 0.5 M NaCl for 100 min (lane 5), 0.7 M NaCl for 50 min (lane 6), 0.7 M NaCl for 100 min (lane 7), 1.5 M NaCl for 50 min (lane 8), and 1.5 M NaCl for 100 min (lane 9). The probes used are indicated below the panels. (d) Products generated by RT-PCR. PCR products were obtained with primers spanning the intergenic regions between *dnaK* and *grpE* (primers 1E and 1K) (lanes 1 to 5), between ORF1 and ORFX (primers 1N and 2H) (lanes 6 to 8), between *grpE* and *dnaJ* (primers 1J and 2E) (lanes 9 to 13), and between *dnaJ* and ORF1 (primers 1H and 2J) (lanes 14 to 18). The positions of the primer pairs used are shown in panel a. Products of the PCR were derived under the following conditions: cDNA prepared from *B. breve* RNA (lanes 1, 2, 6, 7, 11, 12, 14, and 15), a negative control in which *B. breve* RNA was used but reverse transcriptase was omitted (lanes 4, 9, and 17), and no template (lanes 5, 10, and 18). A positive control contained *B. breve* DNA (lanes 3, 8, 13, and 16). Lanes MK contained DNA molecular marker X (Roche).




FIG. 3. Osmotic and heat shock induction of the *B. breve* UCC 2003 *gpe* and ORF1 sequences. Total RNA was isolated from *B. breve* UCC 2003 following exposure to various extremes of osmolarity and temperature for a specific length of time and analyzed by quantitative real-time PCR assays. (a) Curves indicating the PCR amplification cycle profiles of the *gpe* sequences obtained by using cDNA samples from different times after heat shock as templates. The histogram indicates the numbers of copies of *gpe* mRNAs for the specific samples. The cDNAs used were synthesized by using RNA collected from cultures grown at 37°C (samples 1 to 2) and 43°C (lanes 3 to 4) for the following lengths of time after a temperature shift: none (beginning of the experiments) (sample 1), 150 min (samples 2 and 4), and 50 min (sample 3). Sample 6 was the negative control (no template). (b) Curves indicating the PCR amplification cycle profiles of the *gpe* sequences obtained by using cDNA samples from different NaCl concentrations as templates. An asterisk indicates the crossing point. The histogram indicates the numbers of copies of *gpe* mRNAs for the specific samples. The cDNAs used were synthesized by using RNA collected from cultures grown in media with the following NaCl concentrations: no NaCl (sample 1), 0.3 M NaCl (sample 2), 0.5 M NaCl (sample 3), 0.7 M NaCl (sample 4), and 1.5 M NaCl (sample 5). Sample 6 was the negative control (no template).

animalis ATCC 25527 (90.8°C) (Fig. 6b). Similar findings were obtained when other *B. lactis* and *B. animalis* strains were used (data not shown). In addition, the assay clearly differentiated *B. catenulatum* from *B. pseudocatenulatum*, which had T_m s of 88.63 and 89.73°C, respectively.

The analysis of the partial *dnaK* gene sequences of two very closely related taxa, *B. lactis* and *B. animalis*, from which the *B. lactis*-specific primers b-dnaK1 and b-dnaK2 were designed, revealed few nucleotide differences. Primers gro-1 and gro-2 (37), which target two conserved regions within the *groES* gene




FIG. 4. Determination of the *dnkA* transcription initiation site of *B. breve* UCC 2003 by primer extension analysis. (a) Primer extension results obtained by using an oligonucleotide targeting the 5' end of *dnkA*. Lane 1, primer extension product obtained by using mRNA extracted from cells grown at 37°C; lane 2, primer extension product obtained by using mRNA extracted from cells grown with 0.7 M NaCl; lane 3, primer extension product obtained by using mRNA extracted from cells grown at 43°C. (b) Putative promoter sequences for the *dnkA* gene. The -10 and -35 putative hexamers are underlined; boldface type with an asterisk indicates the transcription start point; and boldface type without an asterisk indicates the start codon. IR1 and IR2 are putative regulator sequences. (c) Comparison of the promoter regions of four *dnkA* genes from different members of the *Actinobacteridae*. Shaded residues indicate that the level of identity for the sequences is 50% or higher. (d) Alignment of consensus IR sequences from *B. longum* NCC 2705. Identical nucleotides are shaded. The numbers indicate positions relative to the translational start site.




FIG. 5. Phylogenetic trees obtained by using the *grpE* gene sequences (a) and the *dnaK* gene sequences (b). The bars indicate phylogenetic distances. Bootstrap values for a total of 1,000 replicates are indicated at the nodes.




FIG. 6. Determination of the T_m s for closely related bifidobacterial species in *grpE* real-time PCRs. (a and b) $-dF/dT$ (change in fluorescence/change in temperature) plotted against temperature. (c) PCR products of various *Bifidobacterium* species obtained by using a multiplex PCR approach with *B. lactis* species-specific PCR primers and *Bifidobacterium* genus-specific primers. Lane 1, *B. lactis* DSM 10140; lane 2, *B. lactis* LMG 11615; lane 3, *B. lactis* LMG 18906; lane 4, *B. lactis* LMG 11580; lane 5, *B. animalis* LMG 18900; lane 6, *B. animalis* ATCC 25527; lane 7, *B. angulatum* JCM 7096; lane 8, *B. pseudocatenulatum* JCM 1200; lane 9, *B. breve* UCC 2003; lane 10, *B. magnum* JCM 1218; lane 11, *B. thermophilum* JCM 1207; lane 12, *B. pullorum* DSM 20433; lane 13, *B. bifidum* JCM 1255; lane 14, *B. dentium* JCM 1195; lane 15, *B. suis* JCM 1269; lane 16, *B. infantis* JCM 1210; lane 17, *B. adolescentis* JCM 1275; lane 18, *B. catenulatum* JCM 1194; lane 19, negative control. Lane m, 1-kb DNA ladder.

of the genus *Bifidobacterium*, were also included as a positive control in the same PCR. When a multiplex PCR was performed with this mixture of four primers, two amplicons (290 and 200 bp) were detected only in the presence of DNA isolated from *B. lactis*. Amplification of all other *Bifidobacterium* DNAs resulted in only one amplicon (200 bp). The results of these PCR assays, based on the *dnaK* sequences data, are shown in Fig. 6c. The 290-bp amplicon was apparent for three *B. lactis* strains (Fig. 6c, lanes 1, 2, 3, and 4) but not at all for any of the *B. animalis* strains (Fig. 6c, lanes 5 and 6). Furthermore, we confirmed the overall absence of PCR amplicons for all other bifidobacterial type strains (Fig. 6c and data not shown).

DISCUSSION

The *dnaK* operon of *B. breve* UCC 2003 was identified and shows consistent levels of similarity to previously described *dnaK* operons (28). Our results clearly indicate that the *B. breve* *dnaK* operon consists of four genes: *dnaK*, *grpE*, *dnaJ*, and ORF1. This gene organization is identical to that of presumed *dnaK* regions of other sequenced *Bifidobacterium* species and to that of other available sequences of members of the *Actinobacteridae*. In fact, for *S. coelicolor* A3 it has been demonstrated that the *dnaK* operon is tetracistronic and includes the *dnaK*, *grpE*, and *dnaJ* genes and a gene encoding a heat shock regulator (HspR) (7).

Northern blots indicated that a 4.4-kb transcript encompassing the complete operon (*dnaK-grpE-dnaJ-ORF1*) is produced. This primary transcript seems to be very unstable and subject to rapid processing. It is possible that the many stem-loop structures located within the coding regions, as well as between the *dnaJ* and *grpE* genes, may act as processing sites. A similar situation has previously been described in *B. subtilis* (16), where four hairpin-like structures were implicated in the processing of *dnaKJ* mRNAs. In addition, in both *Clostridium acetobutylicum* and *Nitrosomonas europaea* intensive specific mRNA degradation of the polycistronic *dnaK* transcript has been described (3, 17). The significance of this phenomenon is unknown, but cleavage at processing sites may lead to a product with an enhanced half-life (16). Moreover, it has been postulated that the processing of *dnaK* mRNAs may be an important component of regulation of expression of the genes in the *dnaK* operon (3, 20).

Although relatively little is known about the heat and osmotic stress responses, a previous report (26) and the present study demonstrated that the *dnaK* operon is part of the general stress response of bifidobacteria. We are primarily interested in the role of the DnaK chaperone machine in the heat and osmotic stress responses. In fact, identification of the genetic basis of heat and osmotic resistance for industrially applicable bifidobacteria is highly desirable for selecting strains that resist stressful conditions encountered during food manufacture.

Transcription of the *dnaK* operon from *B. breve* occurs at 37°C and in the absence of NaCl in the medium and is increased upon a shift to higher salt concentrations. The mRNA levels for the *dnaK* operon were shown to increase 14- to 15-fold following an osmotic challenge, while the change in transcriptional levels was shown to be not significant when cells were subjected to heat stress (approximately 1.5-fold). Inter-

estingly, in contrast to *B. breve*, the *dnaK* operons of *B. adolescentis* and *B. longum* were shown to be strongly induced upon heat stress (25). These results indicate that in bifidobacteria the amount and type of chaperones necessary for an adaptive response to heat stress are variable. The observed overlap between the heat and osmotic stress responses have also been described for other gram-positive bacteria (e.g., *L. lactis* and *Streptococcus mutans*), in which heat-inducible proteins are also involved in the osmotic stress response, although the induction factors for DnaK chaperones in heat-shocked or osmotically shocked cells vary (4, 18, 35).

In *B. breve* UCC 2003 the level of temperature-induced activation of the *dnaK* operon appeared to be much lower than the levels of temperature-induced activation of the *groEL* and *groES* operons, which are induced approximately 8- and 12-fold (37). The *groESL* and *dnaK* operons of many bacteria, including the *groES* operon of *B. breve* UCC 2003, contain a conserved inverted repeat sequence (CIRCE), which is believed to be involved in the transcriptional regulation of these genetic units (28, 29, 37, 46). The *dnaK* operon of *B. breve* UCC 2003 does not appear to contain such a CIRCE sequence near its transcription start site or upstream of it.

Of course, we cannot rule out the possibility that when applied under different regimens (e.g., different exposure times and/or different severity of stress), some of these stimuli might result in higher expression of *dnaK*. We will address this question in future work and also test the influence of additional stresses.

The comparison of putative regulatory elements for the *dnaK* operon of three bifidobacteria allowed identification of a putative consensus promoter sequence (IR), which exhibited a consistent degree of homology with the HspR binding site (HAIR). The HAIR sequences were shown to be involved in heat shock regulation in *Streptomyces*, in *Mycobacterium*, and in *Helicobacter pylori* (5, 11, 29). Consequently, it is possible that ORF1 acts as a transcriptional regulator of the *dnaK* operon by binding to the putative operator sequence (IR). Future research will be focused on verifying this hypothesis by demonstrating direct protein-DNA interactions by means of gel retardation and DNase footprinting experiments.

Screening of the publicly available *B. longum* NCC 2705 genome sequence revealed that the IR putative operator sequence is located in the promoter region of the *clpB* gene, which has been extensively described previously as being involved in a stress response. Interestingly, the *clpB* gene has been shown to be part of the HspR regulon in *Streptomyces albus* (11). Moreover, the IR putative operator sequence is found in the promoter region of two other genes (*trmD* and *malQ1*), both of which have been described as being associated with the stress response of *L. lactis* (10). This suggests that all these genes may be governed by the same regulatory system.

In this study we investigated the occurrence of the genes encoding the DnaK and GrpE chaperones in different species of the genus *Bifidobacterium*. These genes are ideal target candidates for diagnostic purposes because they are highly conserved and ubiquitous in bacteria (13, 14). They fulfill all of the prerequisites for suitable phylogenetic markers, such as very high genetic stability and a wide distribution. These alternative molecular markers might corroborate and help to complete the evolutionary history of various bifidobacterial species.

Polyphasic taxonomy (35) has been recognized by the International Committee on Systematic Bacteriology as a new tool for describing species and for revising the current nomenclature of some bacterial groups. In view of its demonstrated effectiveness, sequence analysis of protein-encoding genes (e.g., *grpE* and *dnaK* genes) and use of these genes as alternative phylogenetic markers could be added to the arsenal of rRNA sequence databases and to the relatively small *groEL* (37), *recA* (21, 40), *atpD* (38), and *tuf* (39) sequence databases. The data presented here show that the phylogenetic positions of bifidobacteria based on *grpE* sequences are generally in agreement with the *dnaK*-based phylogeny, suggesting that these genes had the same evolutionary development. Moreover, the *grpE* and *dnaK* trees had a topology similar to that obtained with the 16S rRNA genes. Nevertheless, in contrast to the 16S rRNA gene tree, the *grpE* and *dnaK* trees provide higher phylogenetic resolution for closely related taxa.

The *grpE* and *dnaK* gene analysis has many advantages compared to analyses of other genes. In fact, these genes are common in bifidobacteria, are present as single copies on the genome (25), and contain conserved nucleotide signatures that are suitable targets for PCR primers. In spite of the sequence's limited size, *grpE* and *dnaK* sequence analyses may allow resolution of phylogenetic lineages of bifidobacteria at the species and subspecies levels. In fact, they have an advantage over 16S rRNA gene-based differentiation of species in that they allow differentiation of *B. lactis* and *B. animalis*, as well as *B. catenulatum* and *B. pseudocatenulatum*.

In conclusion, the use of several markers, such as *grpE*, in combination with *dnaK*-or 16S rRNA gene-based tracing techniques (41, 42, 43, 44), may be necessary for reliable identification and phylogenetic analysis of *Bifidobacterium* species. Such an approach should increase the accuracy of molecular identification of bifidobacterial species in microbial ecology studies.

Although the work presented here can only be considered a first step toward an understanding of stress responses in bifidobacteria, detailed analysis of the regulatory mechanisms of the *B. breve* UCC 2003 *dnaK* gene and other stress-induced genes represents a suitable model system for the genus *Bifidobacterium*.

ACKNOWLEDGMENTS

This work was financially supported by Enterprise Ireland (grant BR/1998/202), by the Higher Education Authority Programme for Research in Third Level Institutions, by the Science Foundation Ireland Alimentary Pharmabiotic Centre located at University College Cork, and by Marie Curie Development Host Fellowship HPMD-2000-00027 to M.V.

We thank J. O'Mahony and S. Cusack for excellent discussions and technical assistance with the LightCycler PCR technology. We also thank T. Cross and J. Coughlan of the Department of Zoology, Ecology and Plant Science, University College Cork for allowing us to use the LiCor sequencer.

REFERENCES

- Ahmad, S., A. Selvapandian, and R. K. Bratnagar. 2000. Phylogenetic analysis of gram-positive bacteria based on *grpE*, encoded by the *dnaK* operon. *Int. J. Syst. Bacteriol.* **50**:1761–1766.
- Antonopoulos, D. A., W. M. Russel, and B. A. White. 2003. Phylogenetic reconstruction of Gram-positive organisms based on comparative sequence analysis of molecular chaperones from the ruminal microorganism *Ruminococcus flavefaciens* FD-1. *FEMS Microbiol. Lett.* **227**:1–7.
- Bahl, H., H. Muller, S. Behrens, H. Joseph, and F. Narberhaus. 1995.

Expression of heat shock genes in *Clostridium acetobutylicum*. *FEMS Microbiol. Rev.* **17**:341–348.

- Bianchi, A. A., and F. Baneyx. 1999. Hyperosmotic shock induces the σ_{32} and σ_E stress regulon of *Escherichia coli*. *Mol. Microbiol.* **34**:1029–1038.
- Bucca, G., A. M. E. Brassington, H. J. Schonfeld, and C. P. Smith. 2000. The HspR regulon of *Streptomyces coelicolor*: a role for the DnaK chaperone as a transcriptional co-repressor. *Mol. Microbiol.* **38**:1093–1103.
- Bucca, G., G. Ferina, A. M. Puglia, and C. P. Smith. 1995. The *dnaK* operon of *Streptomyces coelicolor* encodes a novel heat-shock protein which binds to the promoter region of the operon. *Mol. Microbiol.* **17**:663–674.
- Bucca, G., Z. Hindle, and C. P. Smith. 1997. Regulation of the *dnaK* operon of *Streptomyces coelicolor* A3(2) is governed by HspR, an autoregulatory repressor. *J. Bacteriol.* **179**:5999–6004.
- Bukau, B., and A. L. Horwich. 1998. The Hsp70 and Hsp60 chaperone machines. *Cell* **92**:351–366.
- Felsenstein, J. 1993. PHYLIP (Phylogeny Inference Package), version 3.5c. Department of Genetics, University of Washington, Seattle.
- Frees, D., P. Varmanen, and H. Igmer. 2001. Stress tolerance and proteolysis in *L. lactis*. *Mol. Microbiol.* **41**:93–103.
- Grandvalet, C., V. Crecy-Lagard, and P. Mazodier. 1999. The ClpB ATPase of *Streptomyces albus* G belongs to the HspR heat shock regulon. *Mol. Microbiol.* **31**:521–532.
- Grandvalet, C., G. Rapoport, and P. Mazodier. 1998. *hrcA*, encoding the repressor of the *groEL* genes in *Streptomyces albus* G, is associated with a second *dnaJ* gene. *J. Bacteriol.* **180**:5129–5134.
- Gupta, R. S. 1995. Evolution of the chaperonin families (Hsp60, Hsp10 and Tcp-1) of proteins and origin of eukaryotic cells. *Mol. Microbiol.* **15**:1–11.
- Gupta, R. S. 2002. Phylogeny of bacteria: are we now close to understanding it? *ASM News* **68**:284–291.
- Hartl, F. U. 1996. Molecular chaperones in cellular protein folding. *Nature* **381**:571–580.
- Homuth, G., S. Masuda, A. Mogk, Y. Kobayashi, and W. Shumann. 1997. The *dnaK* operon of *Bacillus subtilis* is heptacistronic. *J. Bacteriol.* **179**:1153–1164.
- Iizumi, T., and K. Nakamura. 1997. Cloning, nucleotide sequence, and regulatory analysis of the *Nitrosomonas europaea dnaK* gene. *Appl. Environ. Microbiol.* **63**:1777–1784.
- Kilstrup, M., S. Jacobsen, K. Hammer, and F. K. Vogensen. 1997. Induction of heat shock proteins DnaK, GroEL, and GroES by salt stress in *Lactococcus lactis*. *Appl. Environ. Microbiol.* **63**:1826–1837.
- Kullen, M., L. J. Brady, and D. J. O'Sullivan. 1997. Evaluation of using a short region of the *recA* gene for rapid and sensitive speciation of dominant bifidobacteria in the human large intestine. *FEMS Microbiol. Lett.* **154**:377–383.
- Lemos, J. A. C., Y. Y. M. Chien, and R. A. Burne. 2000. Genetic and physiologic analysis of the *groE* operon and role of the HrcA repressor in stress gene regulation and acid tolerance in *Streptococcus mutans*. *J. Bacteriol.* **183**:6074–6084.
- Lievin, V., I. Pfeiffer, S. Hudault, F. Rochat, D. Brassart, J. R. Neeser, and A. L. Servin. 2000. *Bifidobacterium* strains from resident infant human gastrointestinal microflora exert antimicrobial activity. *Gut* **47**:646–652.
- Liu, H., Y. Li, X. Huang, Y. Kawamura, and T. Ezaki. 2003. Use of the *dnaJ* gene for the detection and identification of all *Legionella pneumophila* serogroups and description of the primers used to detect 16S rDNA sequences of major members of the genus *Legionella*. *Microbiol. Immunol.* **47**:859–869.
- O'Mahony, A., T. F. O'Sullivan, Y. Walsh, A. Vaughan, M. Maher, G. F. Fitzgerald, and D. van Sinderen. 2000. Characterization of antimicrobial producing lactic acid bacteria from malted barley. *J. Int. Brew.* **106**:403–410.
- Sambrook, J., and D. W. Russell. 2001. Molecular cloning: a laboratory manual, 3rd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Schell, M. A., M. Karmirantzou, B. Snel, D. Vilanova, G. Pessi, M. C. Zwhalen, F. Desiere, P. Bork, M. Delley, and F. Arigoni. 2002. The genome sequence of *Bifidobacterium longum* reflects its adaptation to the human gastrointestinal tract. *Proc. Natl. Acad. Sci. USA* **99**:14422–14427.
- Schmidt, G., and R. Zink. 2000. Basic features of stress response in three species of bifidobacteria: *B. longum*, *B. adolescentis*, and *B. breve*. *Int. J. Food Microbiol.* **55**:41–45.
- Schmidt, G., C. Hertel, and W. P. Hammes. 1999. Molecular characterisation of the *dnaK* operon of *Lactobacillus sakei* LTH681. *Syst. Appl. Microbiol.* **22**:321–328.
- Segal, G., and E. Z. Ron. 1996. Regulation and organization of the *groE* and *dnaK* operons in Eubacteria. *FEMS Microbiol. Lett.* **138**:1–10.
- Servant, P., and P. Mazodier. 2001. Negative regulation of the heat shock response in *Streptomyces*. *Arch. Microbiol.* **176**:237–242.
- Shultz, A., B. Tzschaschel, and W. Shumann. 1994. Isolation and analysis of mutants of the *dnaK* operon of *Bacillus subtilis*. *Mol. Microbiol.* **15**:421–429.
- Takekaki, S. I., K. Okuzumi, H. Ishiko, K. I. Nakahara, A. Ohkubo, and R. Nagai. 1993. Genus-specific polymerase chain reaction for the mycobacterial *dnaJ* gene and species-specific oligonucleotide probes. *J. Clin. Microbiol.* **31**:446–450.
- Tannock, G. W. 1994. The acquisition of the normal microflora of the

- gastrointestinal tract, p 1–16. In S. A. Gibson (ed.), Human health: the contribution of microorganisms. Springer, London, United Kingdom.
33. **Thompson, J. D., D. G. Higgins, and T. J. Gibson.** 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
 34. **Van de Guchte, M., P. Serror, C. Chervaux, T. Smokvina, S. D. Ehrlich, and E. Manguin.** 2002. Stress responses in lactic acid bacteria. *Antonie Leeuwenhoek* **82**:187–216.
 35. **Vandamme, P., B. Pot, M. Gillis, P. de Vos, K. Kersters, and J. Swings.** 1996. Polyphasic taxonomy, a consensus approach to bacterial systematics. *Microbiol. Rev.* **60**:407–438.
 36. **Ventura, M., D. van Sinderen, G. F. Fitzgerald, and R. Zink.** 2004. Insights into the taxonomy, genetics and physiology of bifidobacteria. *Antonie Leeuwenhoek* **86**:205–223.
 37. **Ventura, M., C. Canchaya, R. Zink, G. F. Fitzgerald, and D. van Sinderen.** 2004. Characterization of the *groEL* and *groES* loci in *Bifidobacterium breve* UCC 2003: genetic, transcriptional and phylogenetic analyses. *Appl. Environ. Microbiol.* **70**:6197–6209.
 38. **Ventura, M., C. Canchaya, D. van Sinderen, G. F. Fitzgerald, and R. Zink.** 2004. *Bifidobacterium lactis* DSM 10140: identification of the *atp* (*atpBEF-HAGDC*) operon and analysis of its genetic structure, characteristics, and phylogeny. *Appl. Environ. Microbiol.* **70**:3110–3121.
 39. **Ventura, M., C. Canchaya, V. Meylan, T. R. Klaenhammer, and R. Zink.** 2003. Analysis, characterization, and loci of the *tuf* genes in *Lactobacillus* and *Bifidobacterium* and their direct application for species identification. *Appl. Environ. Microbiol.* **69**:6908–6922.
 40. **Ventura, M., and R. Zink.** 2003. Comparative sequence analysis of the *tuf* and *recA* genes and restriction fragment length polymorphism of the internal transcribed spacer region sequences supply additional tools for discriminating *Bifidobacterium lactis* from *Bifidobacterium animalis*. *Appl. Environ. Microbiol.* **69**:7517–7522.
 41. **Ventura, M., V. Meylan, and R. Zink.** 2003. Identification and tracing of *Bifidobacterium* species by enterobacterial repetitive intergenic consensus sequences. *Appl. Environ. Microbiol.* **69**:4296–4301.
 42. **Ventura, M., and R. Zink.** 2002. Rapid identification, differentiation, and proposed new taxonomic classification of *Bifidobacterium lactis*. *Appl. Environ. Microbiol.* **68**:6429–6434.
 43. **Ventura, M., M. Elli, R. Reniero, and R. Zink.** 2001. Molecular microbial analysis of *Bifidobacterium* isolates from different environments by the species-specific amplified ribosomal DNA restriction analysis (ARDRA). *FEMS Microbiol. Ecol.* **36**:113–121.
 44. **Ventura, M., R. Reniero, and R. Zink.** 2001. Specific identification and targeted characterization of *Bifidobacterium lactis* from different environmental isolates by a combined multiplex-PCR approach. *Appl. Environ. Microbiol.* **67**:2760–2765.
 45. **Yuan, G., and S. Wong.** 1995. Isolation and characterization of *Bacillus subtilis* *groE* regulatory mutants: evidence for *orf39* in the *dnaK* operon as a repressor gene in regulating the expression of both *groE* and *dnaK*. *J. Bacteriol.* **177**:6462–6468.
 46. **Zuber, U., and W. Schumann.** 1994. CIRCE, a novel heat shock element involved in regulation of heat shock operon *dnaK* of *Bacillus subtilis*. *J. Bacteriol.* **176**:1353–1363.