

## The *Bifidobacterium longum* NCIMB 702259<sup>T</sup> *ctr* Gene Codes for a Novel Chololate Transporter

Claire E. Price,<sup>1</sup> Sharon J. Reid,<sup>1</sup> Arnold J. M. Driessen,<sup>2</sup>  
 and Valerie R. Abratt<sup>1\*</sup>

Department of Molecular and Cell Biology, University of Cape Town, Rondebosch,  
 Cape Town, South Africa,<sup>1</sup> and University of Groningen,  
 Groningen, The Netherlands<sup>2</sup>

Received 13 July 2005/Accepted 2 November 2005

**Preexposure of *Bifidobacterium longum* NCIMB 702259<sup>T</sup> to chololate caused increased resistance to chololate, chloramphenicol, and erythromycin. The *B. longum ctr* gene, encoding a chololate efflux transporter, was transformed into the efflux-negative mutant *Escherichia coli* KAM3, conferring resistance to bile salts and other antimicrobial compounds and causing the efflux of [<sup>14</sup>C]chololate.**

Bifidobacteria are major components of the human intestinal microflora (13) and are widely used as probiotics in food supplements. Probiotic survival depends on resistance to antibiotics and to inhibitory host-produced substances, such as bile salts (9). Bifidobacteria are resistant to a range of antibiotic compounds (5), which could allow them to withstand concurrent antibiotic administration. This study aimed to identify and prove the functionality of a possible efflux system encoded by the *ctr* gene in *Bifidobacterium longum* which may contribute to bile and antibiotic resistance.

**Adaptation to sodium glycocholate and antibiotics.** To determine the intrinsic MICs for *B. longum* NCIMB 702259<sup>T</sup> (NCIMB, United Kingdom) of antimicrobial agents, 10 μl of a standard cell suspension (optical density at 600 nm, 0.5) of a culture grown anaerobically on BYG agar (14) was spotted onto BYG plates containing a twofold dilution range of sodium glycocholate, ampicillin, chloramphenicol, erythromycin, or tetracycline. Adaptation to the antibiotics was tested using a method modified from the work of Carsenti-Etesse et al. (4). Mid-exponential-phase *B. longum* cells grown in BYG broth were streaked for four passages onto sodium glycocholate gradient plates, and the MICs for these cells were tested as described above. Adapted *B. longum* showed an increase in resistance to sodium glycocholate, chloramphenicol, and erythromycin but not to ampicillin and tetracycline (Table 1). This indicated that *B. longum* may possess multidrug transporters, since these are often regulated by the compounds that they transport but may confer resistance to structurally unrelated antimicrobial agents (3).

**Cloning and antimicrobial characterization of the *ctr* gene.** Open reading frame BL1102 (*B. longum* NCC 2705, GenBank accession number AE014295) was identified as a possible sodium-dependent bile acid transporter. The BL1102 orthologue was isolated from *B. longum* NCIMB 702259<sup>T</sup> genomic DNA

TABLE 1. MICs of antimicrobial agents tested against *B. longum* NCIMB 702259<sup>a</sup>

Treatment	Antimicrobial MIC				
	Amp (μg/ml)	Chl (μg/ml)	Chololate (%)	Em (μg/ml)	Tet (μg/ml)
Control	1.6	1.6	0.8	0.4	1.6
Chololate preexposure	1.6	3.2	3.2	0.8	1.6

<sup>a</sup> MICs of antimicrobial agents tested against *B. longum* NCIMB 702259, without (control) and following preexposure to chololate, as determined by plating cells onto BYG plates containing a twofold dilution range of each antibiotic. The antimicrobial agents used were ampicillin (Amp), chloramphenicol (Chl), erythromycin (Em), tetracycline (Tet), and sodium glycocholate (chololate). The experiments were done in triplicate.

(14), using standard PCR protocols and the primers ctrans-F (5'-AGCTGAATTCGCGCAACAGG-3') and ctrans-R (5'-ACGCCCGGTACCTCAATCG-3'). EcoRI and KpnI restriction enzyme sites (underlined) were introduced to ctrans-F and ctrans-R, respectively, to assist subcloning into pBluescriptSK. The nucleotide sequence of the insert in the recombinant plasmid pCtr was determined (14), and nucleotide and amino acid homology searches were performed using the BLAST algorithm and NCBI databases (1). The deduced amino acid sequence of Ctr was 100% identical to that of BL1102. Plasmid pCtr was transformed into competent (2) *Escherichia coli*

TABLE 2. MICs for *E. coli* KAM3 harboring pCtr and the control vector pBluescriptSK as determined by the broth dilution method<sup>a</sup>

Plasmid	MIC of:						
	Acr (μg/ml)	Chl (μg/ml)	Em (μg/ml)	EtBr (μg/ml)	Chololate (%)	SDS (%)	Tet (μg/ml)
pBluescriptSK	6.25	0.4	1.25	12.5	0.5	0.01	0.4
pCtr	25	1.6	2.5	50	8	0.02	1.6

<sup>a</sup> The antimicrobial agents used were acriflavine (Acr), ampicillin (Amp), chloramphenicol (Chl), erythromycin (Em), ethidium bromide (EtBr), sodium dodecyl sulfate (SDS), sodium glycocholate (chololate), and tetracycline (Tet). The experiments were done in triplicate.

\* Corresponding author. Mailing address: Department of Molecular and Cell Biology, University of Cape Town, Rondebosch, 7701 Cape Town, South Africa. Phone: 27 21 6502183. Fax: 27 21 6897573. E-mail: val@science.uct.ac.za.




FIG. 1. Energy-dependent extrusion of [<sup>14</sup>C]cholate in *E. coli* KAM3 harboring (A) pCtr or (B) pBluescriptSK. Cells were preloaded with cholate, and the amount of cell-associated cholate was subsequently measured over time with the addition of glucose (dotted lines) or without glucose (solid lines) Glucose was added to a final concentration of 10 mM at 35 min as indicated by the arrow. During each experiment, samples were taken in triplicate, and the entire experiment was also performed in triplicate. Error bars indicate the deviation from the mean.

KAM3 (11), a K-12 derivative lacking the multidrug transporter AcrAB. The MICs of acriflavine, sodium dodecyl sulfate (Merck), chloramphenicol, erythromycin, ethidium bromide, tetracycline (Sigma), and sodium glycocholate (Difco) were determined using the broth dilution method (8). Plasmid pCtr conferred cholate resistance on *E. coli* KAM3, increasing the

MIC of sodium glycocholate by 16-fold (Table 2). Resistance to the antimicrobial agents was increased by two- to fourfold.

**Efflux of [<sup>14</sup>C]cholate.** To determine whether pCtr conferred resistance to bile through the active efflux of the compound, de-energized washed cell suspensions of *E. coli* KAM3 (pCtr or pBluescriptSK) that had been grown to mid-exponen-




FIG. 2. Multiple sequence alignment of the significant regions of the putative *B. longum* bile transporter Ctr (GenBank accession no. DQ017587) with closely related bacterial sodium/bile acid transporters. Sequences included are from the following organisms: *B. longum* (Ctr) (NP\_696274), *Streptococcus thermophilus* (S.the) (YP\_141686), *Streptococcus mutans* (S.mat) (NP\_721034), *Leuconostoc mesenteroides* (L.mes) (ZP\_00063561), *Oceanobacillus iheyensis* (O.ihe) (NP\_691915), *Bacillus subtilis* (B.sub) (CAB13827.1), *Pseudomonas fluorescens* (P.flu) (ZP\_00262238.1), and *Neisseria meningitidis* (N.men) (NP\_273747). Amino acids conserved in all sequences are shaded in dark gray, and amino acids conserved in over 75% of the sequences are shaded in light gray. The conserved proline residue is indicated by an asterisk, and the SBF signature motif is underlined.

Downloaded from http://aem.asm.org/ on November 26, 2020 by guest




FIG. 3. Phylogenetic relationship analysis of bacterial sodium/bile acid transporters, namely, Ctr from *B. longum*, *Streptococcus thermophilus*, *Streptococcus mutans*, *Leuconostoc mesenteroides*, *Oceanobacillus iheyensis*, *Bacillus subtilis*, *Pseudomonas fluorescens*, and *Neisseria meningitidis*; the plant sodium/bile acid symporter-like proteins AtSbf1 (*Arabidopsis thaliana*) and OsSbf1 (*Oryza sativa*); mammalian ileal sodium/bile acid transporters (IBAT) from *Homo sapiens*, *Mus musculus*, *Oryctolagus cuniculus*, and *Rattus norvegicus*; mammalian hepatic sodium/bile acid transporters (HBAT) from *M. musculus* and *O. cuniculus*; and the arsenate resistance protein (ACR3) from *Saccharomyces cerevisiae*. Database accession numbers are given in parentheses. The tree was constructed using the neighbor-joining method, and bootstrap values (250 replicates) are given at the branch points.

tial phase in Luria-Bertani broth (14) were preloaded with [*carboxyl*-<sup>14</sup>C]cholic acid (New England Nuclear Corp.). The amount of cell-associated radioactivity was monitored with and without the addition of glucose as an energy source. The method of Yokota et al. (17) was used with the following modifications. Washed cells were resuspended to an optical density at 600 nm of 4, aliquots of 1.94 ml were used in the experiment, and all incubation was at 37°C. To preload the cells with cholate, 40 μl of 5.8 mM [<sup>14</sup>C]cholate (16 mCi/mmol) was added (final cholate concentration, 0.116 mM). The amount of radioactivity associated with each aliquot was used to calculate the counts per minute per mmol of cholate. The results were expressed as nmol cholate/μg protein, determined using a DC protein assay kit (Bio-Rad). In the absence of glucose, the external and internal cellular cholate concentrations reached equilibrium within 35 min (Fig. 1A and B). Upon the addition of glucose, cholate was transiently accumulated in the presence and absence of pCtr. In the absence of pCtr, an equilibrium was again reached after 40 min (Fig. 1B). When pCtr was present, however, there was a decrease in the level of cell-associated cholic acid (Fig. 1A), indicating an active efflux of cholate. Glucose metabolism results in a slow generation of a pH gradient, which is likely to drive the initial accumulation of cholate. The subsequent activity of the Na<sup>+</sup>/H<sup>+</sup> antiporter results in the generation of a sodium gradient necessary to drive the cholate transporter. This is evidence that the *ctr* gene

of *B. longum* encodes a cholate efflux transport system that is functional in *E. coli*.

#### Bioinformatic and phylogenetic analysis of the Ctr protein.

Ctr belongs to the sodium/bile acid family (SBF) of transporters (10), showing the signature motif of this family (Fig. 2). Analysis of the predicted membrane topology revealed the presence of nine transmembrane segments as well as a highly conserved proline residue, corresponding to P<sup>290</sup> in the human bile transporter (Fig. 2), which is an essential residue for bile acid transport (15). The phylogenetic relationship of various SBF proteins from different taxa was determined using the neighbor-joining method of ClustalW (Fig. 3). The Ctr protein is closely related to a number of sodium bile acid cotransporter proteins from bacteria, including two *Streptococcus* species and *Leuconostoc mesenteroides*. Prior to this study, the members of the SBF family with proved function were all in eukaryotes, and in mammals, these transmembrane proteins are responsible for the cotransport of sodium and bile acids across the plasma membrane in the liver and ileum (6, 7). SBF transporters from plants, namely, *Arabidopsis thaliana* and *Oryza sativa* (12), form a separate distinct cluster. These are inducible during growth, but neither their efflux function nor their substrates have been established. The ACR3 protein from *Saccharomyces cerevisiae* is an efflux transmembrane protein involved in resistance to arsenic compounds (16).

In this study we have confirmed that the *ctr* gene of *B. longum* encodes a cholate transporter which is responsible for the efflux of cholate from *E. coli* and confers resistance to a number of structurally unrelated antimicrobial compounds. This is the first characterization of a bile acid transporter of the SBF family in bacteria and the first multidrug transporter to be characterized from a *Bifidobacterium* species.

**Nucleotide sequence accession number.** The GenBank accession number for the putative *B. longum* bile transporter Ctr is DQ017587.

We thank the National Research Foundation (NRF) South Africa for financial support of this project. C. E. Price acknowledges an NRF bursary and a University of Cape Town travel scholarship.

#### REFERENCES

- Altschul, S. F., T. L. Madden, A. A. Schaffer, J. Zhang, Z. Zhang, W. Miller, and D. J. Lipman. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**:3389–3402.
- Armitage, P., R. Walden, and J. Draper. 1988. Vectors for the transformation of plant cells using agrobacterium, p. 46–49. *In* J. Draper, R. Scott, P. Armitage, and W. Walden (ed.), *Plant genetic transformation and gene expression, a laboratory manual*. Blackwell Scientific Publications Ltd., Oxford, United Kingdom.
- Baranova, N. N., A. Danchin, and A. A. Neyfakh. 1999. Mta, a global MerR-type regulator of the *Bacillus subtilis* multidrug-efflux transporters. *Mol. Microbiol.* **31**:1549–1559.
- Carsenti-Etesse, H., P.-M. Roger, B. Dunais, S. Durgeat, G. Mancini, M. Bensoussan, and H. Schweizer. 1999. Gradient plate method to induce *Streptococcus pyogenes* resistance. *J. Antimicrob. Chemother.* **44**:439–443.
- Charteris, W. P., P. M. Kelly, L. Morelli, and J. K. Collins. 2000. Effect of conjugated bile salts on antibiotic susceptibility of bile salt tolerant *Lactobacillus* and *Bifidobacterium* isolates. *J. Food Prot.* **63**:1369–1379.
- Hagenbuch, B., and P. Dawson. 2004. The sodium bile salt cotransporter family SLC10. *Pflügers Arch. Eur. J. Physiol.* **447**:566–570.
- Hagenbuch, B., and P. J. Meier. 1994. Molecular cloning, chromosomal localization, and functional characterization of a human liver Na<sup>+</sup> bile salt cotransporter. *J. Clin. Investig.* **93**:1326–1331.
- Koneman, E. W., S. D. Allen, V. R. Dowell, Jr., W. M. Jando, H. M. Sommer, and W. C. Winn, Jr. 1988. Antimicrobial susceptibility testing, p. 473. *In* E. W. Koneman (ed.), *Color atlas and textbook of diagnostic microbiology*, 3rd ed. J. B. Lippincott Co., Philadelphia, Pa.
- Kurdi, P. H. Tanaka, H. van Veen, K. Asano, F. Tomita, A., and Yokota.

2003. Cholic acid accumulation and its diminution by short-chain fatty acids in bifidobacteria. *Microbiology* **149**:2031–2037.
10. **Marchler-Bauer, A., and S. H. Bryant.** 2004. CD-search: protein domain annotations on the fly. *Nucleic Acids Res.* **32**:W327–W331.
  11. **Morita, Y., K. Kodama, S. Shiota, T. Mine, A. Kataoka, T. Mizushima, and T. Tsuchiya.** 1998. NorM, a putative multidrug efflux protein, of *Vibrio parahaemolyticus* and its homolog in *Escherichia coli*. *Antimicrob. Agents Chemother.* **42**:1778–1782.
  12. **Rzewuski, G., and M. Sauter.** 2002. The novel rice (*Oryza sativa* L.) gene *OsSbf1* encodes a putative member of the Na<sup>+</sup>/bile acid symporter family. *J. Exp. Bot.* **53**:1991–1993.
  13. **Simon, G. L., and S. L. Gorbach.** 1984. Intestinal flora in health and disease. *Gastroenterology* **86**:174–193.
  14. **Trindade, M. I., V. R. Abratt, and S. J. Reid.** 2003. Induction of sucrose-utilization genes from *Bifidobacterium lactis* by sucrose and raffinose. *Appl. Environ. Microbiol.* **69**:24–32.
  15. **Wong, M. H., P. Oelkers, and P. A. Dawson.** 1995. Identification of a mutation in the ileal sodium-dependent bile acid transporter gene that abolishes transport activity. *J. Biol. Chem.* **270**:27228–27234.
  16. **Wysocki, R., P. Bobrowicz, and S. Ulaszewski.** 1997. The *Saccharomyces cerevisiae* *ACR3* gene encodes a putative membrane protein involved in arsenite transport. *J. Biol. Chem.* **272**:30061–30066.
  17. **Yokota, A., M. Veenstra, P. Kurdi, H. W. van Veen, and W. N. Konings.** 2000. Chololate resistance in *Lactococcus lactis* is mediated by an ATP-dependent multispecific organic anion transporter. *J. Bacteriol.* **182**:5196–5201.