

Modification of Sexual Development and Carotene Production by Acetate and Other Small Carboxylic Acids in *Blakeslea trispora* and *Phycomyces blakesleeanus*

Vera Kuzina and Enrique Cerdá-Olmedo*

Departamento de Genética, Facultad de Biología, Universidad de Sevilla, E-41080 Sevilla, Spain

Received 2 December 2005/Accepted 11 April 2006

In *Phycomyces blakesleeanus* and *Blakeslea trispora* (order Mucorales, class Zygomycetes), sexual interaction on solid substrates leads to zygospore development and to increased carotene production (sexual carotenogenesis). Addition of small quantities of acetate, propionate, lactate, or leucine to mated cultures on minimal medium stimulated zygospore production and inhibited sexual carotenogenesis in both *Phycomyces* and *Blakeslea*. In *Blakeslea*, the threshold acetate concentration was <1 mmol/liter for both effects, and the concentrations that had one-half of the maximal effect were <2 mmol/liter for carotenogenesis and >7 mmol/liter for zygospore formation. The effects on *Phycomyces* were similar, but the concentrations of acetate had to be multiplied by ca. 3 to obtain the same results. Inhibition of sexual carotenogenesis by acetate occurred normally in *Phycomyces* mutants that cannot use acetate as a carbon source and in mutants whose dormant spores cannot be activated by acetate. Small carboxylic acids may be signals that, independent of their ability to trigger spore germination in *Phycomyces*, modify metabolism and development during the sexual cycle of *Phycomyces* and *Blakeslea*, uncoupling two processes that were thought to be linked and mediated by a common mechanism.

Phycomyces blakesleeanus and *Blakeslea trispora* are actual or potential sources of commercial carotenes used as provitamins, pigments, and antioxidants in the food and feed, pharmaceutical, and cosmetics industries (3, 10, 13, 26). Mutations in various genes (11, 20, 22) result in very high β -carotene contents or interrupt the carotene pathway, allowing production of lycopene and other carotenes. *Phycomyces* stops β -carotene biosynthesis through a feedback mechanism when a certain concentration has been reached and responds to four groups of activators with different modes of action typified by sexual activity, illumination, and addition of retinol or dimethyl phthalate to the medium (5).

Blakeslee (6) classified strains of many Mucorales into two sexes, (+) and (–), by noting that when mycelia of opposite sexes meet on solid media, they become bright yellow and produce a succession of special structures, particularly zygospores. The enhanced coloration is due to increased accumulation of β -carotene, which is termed sexual carotenogenesis to distinguish it from the vegetative carotenogenesis that occurs in single-strain cultures. Sexual carotenogenesis and sexual morphogenesis are induced by trisporic acid C and related trisporoids, which are produced cooperatively by neighboring mycelia of different sexes (8, 34). Sexual carotenogenesis occurs in mated cultures that contain mycelia of the two sexes, in single-strain cultures of either sex exposed to natural or synthetic trisporates (17), in single-strain cultures of intersexual heterokaryons, whose mycelia contain nuclei of both sexes (22, 23), and in single-strain cultures of intersexual diploids (21). In all known mutants the carotene content is increased further by sexual stimulation.

Phycomyces and *Blakeslea* grow on acetate as a sole carbon

source. Acetate must first be converted to acetyl coenzyme A (acetyl-CoA) by an acetyl-CoA synthetase (EC 6.2.1.1). Mutants of *Phycomyces* resistant to fluoroacetate could not utilize acetate because of the loss of the acetyl-CoA synthetase encoded by the *facA* gene (15, 16). *Phycomyces* produces a different acetyl-CoA synthetase in response to carbon starvation (14).

Acetate also has regulatory effects on development. Very few vegetative spores of *Phycomyces* germinate after they are inoculated into a minimal medium in which growth and differentiation occur (28). Spore dormancy can be broken by various treatments, including exposure to heat, acetate, propionate, or other chemicals, some of which cannot be metabolized and others of which are toxic (29, 36, 38). The *ger* mutants of *Phycomyces*, isolated for decreased acetate or propionate spore activation, also are less responsive to activation with heat and do not exhibit the transient increase in the cyclic AMP level that immediately follows such treatments in wild-type spores. In these mutants a protein that detects heat shock, small carboxylic acids, and perhaps other compounds and that transduces the signal for germination may be altered (27, 37).

The objective of this study was to describe novel effects of acetate and other small carboxylic acids on the sexual processes of two zygomycete fungi. We examined whether the morphogenetic and metabolic changes induced by sexual interaction are tightly coupled as a single response. The results are significant for understanding sexuality in the Zygomycetes and for practical application of this sexuality in the carotene industry.

MATERIALS AND METHODS

Strains. NRRL1554 and NRRL1555 are natural (+) and (–) strains of *P. blakesleeanus*, respectively, and were obtained originally from the Northern Regional Research Laboratory (now the National Center for Agricultural Utilization Research, Peoria, IL). Strains F986 and F921 are wild-type (+) and (–) strains of *B. trispora*, respectively, and were obtained from VKM (All-Russian Collection of Microorganisms, Moscow, Russia). *Phycomyces* strains MU136,

* Corresponding author. Mailing address: Departamento de Genética, Facultad de Biología, Apartado 1095, E-41080 Sevilla, Spain. Phone: 0034-954624107. Fax: 0034-954557104. E-mail: eco@us.es.

MU138, and MU141 are *facA* mutants that are resistant to fluoroacetate and are defective for acetyl-CoA synthetase activity (16). Strains S437 and S440 are *ger* mutants whose spores show decreased activation by heat, acetate, and propionate (27). The mutants were isolated after treatment of NRRL1555 spores with *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine.

Culture conditions. Cultures were grown for 4 days (unless indicated otherwise) at 22°C (*Phycomyces*) or at 30°C (*Blakeslea*) in the dark on cellophane disks (Sadipal, Gerona, Spain) placed on the top surface of 25 ml of agar medium in petri dishes (diameter, 85 mm). The minimal medium (9, 33) contained 20 g/liter D-(+)-glucose and 2 g/liter L-asparagine as carbon and nitrogen sources. Glutamate medium contained monosodium L-glutamate (1 g/liter) instead of asparagine. Potato dextrose medium prepared with fresh potatoes (9) gave more reproducible results than the dried commercial form of this medium. *Phycomyces* spores were collected from 4-day-old cultures by rinsing the cultures with sterile distilled water and then were activated by heat shock before plating (9); *Blakeslea* spores were collected from older cultures (1 to 2 weeks) with glycerol (1:3 [vol/vol] in water). Spore stocks were kept refrigerated for no more than 1 week. Each culture was started with 10^4 spores, and half of each sex was used for mated cultures. Extracts obtained by freezing the media (−20°C for at least 2 h), thawing the media (22°C for 1 h), and centrifuging the liquid ($1,000 \times g$, 10 min, 22°C) were used for glucose determination. Similar extracts obtained from media on which mated cultures of *Blakeslea* had grown for 2 days were used instead of water for preparation of minimal medium with trisporic acids.

Quantification of zygospores. Spores were plated directly on a medium without a cellophane disk. The zygospores of *Blakeslea* are distinct and small (diameter, ~50 μm) and were counted with a stereomicroscope. The zygospores of *Phycomyces* are much larger (diameter, ~500 μm), superimposed, and decorated with abundant thorns, which makes them difficult to count; these spores were scraped off the medium with no attempt at purification and weighed.

Chemical analyses. For carotene analyses, mycelia were scraped from the cellophane disks, lyophilized, weighed, and ground with a mortar and pestle in the presence of sand and petroleum ether (boiling point, 40 to 60°C). The extract was centrifuged ($1,000 \times g$, 5 min, 22°C), vacuum dried, and dissolved in *n*-hexane. When possible, the procedures were carried out on ice under dim light, and the extracts were kept under a nitrogen atmosphere. A 10- to 20-μl aliquot of extract was loaded using a G1313A autosampler (Hewlett-Packard, Palo Alto, CA) into a C₁₈ column (4.6 by 100 mm; 5-μm octyldecylsilane particles; Hypersil; Waters, Milford, MA) with a 10-mm refillable guard precolumn filled with the same material (Alltech, Deerfield, IL) in a series 1100 liquid chromatograph (Hewlett-Packard). The column was eluted at room temperature with methanol-acetonitrile-chloroform (47:47:6, vol/vol/vol) at a flow rate of 1 ml/min. The outflow was monitored with a diode array detector at 286, 450, 462, and 473 nm, the absorption maxima of phytoene, β-carotene, γ-carotene, and lycopene, respectively. Concentrations were calculated following calibration with samples of the four carotenes purified from *Phycomyces* cultures. The significance of differences in carotene concentrations was determined by standard tests (Student's *t* test and nonparametric rank test).

Concentrations of trisporic acids were estimated (24) from *A*₃₂₅ values by using an extinction coefficient (1 cm, 1 g/liter) of 57.2. The glucose content was determined with an oxidase-peroxidase kit (Sigma Chemical Co., St. Louis, MO).

RESULTS

Modification of sexual processes by acetate. The presence of small amounts of acetate in the media modified the sexual interaction in mated cultures of both *Phycomyces* and *Blakeslea* (Fig. 1A). Zygospores, which are produced only when mating occurs, were much more abundant and appeared earlier in the presence of acetate than in the absence of acetate (Fig. 1B and similar observations with *Phycomyces*). Single-strain cultures were slightly colored (yellow for *Phycomyces* and orange for *Blakeslea*), and there was some variation among different wild-type strains. To properly identify the colors of mated cultures, zygospores must be scraped off the plates. Mated cultures in the absence of acetate and young mated cultures in the presence of acetate both had enhanced pigmentation due to sexual carotenogenesis. Older mated cultures in the presence of acetate appeared to contain approximately the same amount of pigment as unmated single-strain cultures contained.




FIG. 1. (A) Modification of sexual responses by acetate: mated cultures of *Phycomyces* and *Blakeslea* wild-type strains grown on minimal medium with or without 10 mmol/liter sodium acetate and on potato dextrose medium. (B) Single-strain and mated cultures of *Blakeslea* strains F921 and F986 grown for the numbers of days indicated next to the images on minimal medium (upper rows of plates) or on minimal medium with 10 mmol/liter sodium acetate (lower rows of plates).

Zygospore formation. Few zygospores were produced on minimal medium, but addition of acetate increased the zygospore density about 10-fold in *Phycomyces* cultures and about 25-fold in *Blakeslea* cultures (Table 1). The resulting values were greater than those obtained with glutamate medium, which is recommended (33) for the production of abundant zygospores on thin mycelia with well-separated hyphae. The threshold for a response to sodium acetate was ≤ 1 mmol/liter,

TABLE 1. Zygospor production on different media^a

Medium	<i>Phycomyces</i> (mg [wet mass]/plate)	<i>Blakeslea</i> (zygospores/mm ²)
Minimal (asparagine)	36 ± 9	6 ± 1
Minimal + acetate	380 ± 23	150 ± 17
Glutamate	140 ± 11	83 ± 9
Potato dextrose	670 ± 32	140 ± 19
Potato dextrose + acetate	670 ± 10	140 ± 17
Potato dextrose + asparagine	770 ± 57	100 ± 8
Potato dextrose + asparagine + acetate	510 ± 33	120 ± 23

^a The values are means ± standard errors for 3 to 17 determinations in three or four independent experiments. The compounds added to the standard media were acetate at a concentration of 10 mmol/liter and L-asparagine · H₂O at a concentration of 2 g/liter.

and the response was not saturated with 20 mmol/liter (Fig. 2); >7 mmol/liter was needed for the response to reach one-half the maximal value.

The effects of propionate, leucine, and lactate were qualitatively similar to the effects of acetate except for propionate at a concentration of 10 mmol/liter, in the presence of which there was not sufficient growth of *Blakeslea* for observation of any sexual response. In single-strain cultures these compounds failed to induce the formation not only of zygospores but also of zygothores, the thickened hyphal tips that are the first morphological stage of the sexual response.

Carotene production. The increase in carotene content due to sexual interactions is particularly well known in *Blakeslea*, and mated *Blakeslea* cultures contained 4.2 ± 0.4 mg carotene/g (dry mass) or 13 times the average for the single-strain cultures (Table 2). The amount of carotenoids produced varied with the strain, and F921 contained more carotene than F986 contained. The sexual response of *Phycomyces*, 0.67 ± 0.04 mg carotene/g (dry mass), was only four times the average for single-strain cultures, which were similar in this respect. The increases due to sexual interaction were very significant ($P \ll 0.001$ for both *Blakeslea* and *Phycomyces*). Potato dextrose medium, which is commonly used in sexual cycle research, is not suitable for carotene production or sexual carotenogenesis, with single and mated wild-type strains of *Phycomyces* containing only 30 µg β-carotene per g (dry mass).

Mated cultures grown in the presence of acetate contained much less carotene than mated cultures grown in the absence of acetate contained (Table 2), and the effect was very significant ($P \ll 0.001$ with 10 mmol/liter in both *Blakeslea* and *Phycomyces* cultures). Acetate can be considered an inhibitor of sexual carotenogenesis. Mated cultures with acetate contained only a little more carotene than single-strain cultures without acetate contained; the increase (28% more on average with 10 mmol/liter acetate) was significant ($P < 0.01$) after all the results were pooled. The presence of acetate in single-strain cultures decreased the carotene content (with 10 mmol/liter acetate, to 70% of the content without acetate, on average); the difference was very significant ($P < 0.001$) only after all the results were pooled.

Leucine, DL-lactate, and propionate also were inhibitory, but DL-lactate was less effective with *Blakeslea*. In single-strain cultures, leucine slightly increased the carotene content, consistent with previous observations (12), and the other compounds appeared to be slightly inhibitory in some experiments.

The addition of acetate slightly increased the pH of the medium, but the results did not change when the experiments were repeated with media whose pH was adjusted to 5.4 ± 0.1 before inoculation.

With the exception of *Blakeslea* growing on media containing propionate, there was little or no variation in the overall growth of the cultures, as measured by the amount of dry mycelial mass per plate. For 4-day-old *Phycomyces* single-strain and mated cultures the average dry mycelial mass was 0.22 g per plate. *Blakeslea* strain F986 grew better than F921 (0.20 and 0.16 g/plate, respectively; significantly different at a P value of 0.01), and the average dry mycelial mass for mated cultures was 0.15 g/plate. In cultures of both fungi, glucose was depleted about 2 days after the cultures reached the maximal amount of dry mass. Thus, ~2.5 g/liter glucose remained in the medium of 4-day-old *Phycomyces* cultures.

For *Blakeslea* the threshold acetate concentration for inhibition of sexual carotenogenesis was <1 mmol/liter, and the response was saturated at ~5 mmol/liter. The threshold for *Phycomyces* was similar, but higher acetate concentrations were needed to saturate the response (Fig. 3).

The time courses of sexual carotenogenesis differed for *Phycomyces* and *Blakeslea*. *Phycomyces*, incubated at 22°C, was growing actively at 2 days, and most of the dry mass and most of the carotene in older cultures were produced on day 3 and the following days. In single-strain *Phycomyces* cultures, β-carotene is neither destroyed nor metabolized appreciably (4), but in mated cultures a considerable amount of β-carotene is used for production of trisporates (2). Sexual carotenogenesis increased with age, and the maximum amount of carotene (~0.8 mg total carotene per g [dry mass]) was present in older cultures (Fig. 4).

Blakeslea incubated at 30°C grew most actively on the first day, when it produced one-half of the mature mass, and the remaining mycelial mass was produced by day 3. The maximum sexual carotenogenesis (~8 mg total carotene per g [dry mass]) occurred in




FIG. 2. Dependence of zygospore production on acetate concentration. Mated cultures of *Phycomyces* wild-type strains (○) and *Blakeslea* wild-type strains (▲) were grown on minimal medium with various concentrations of sodium acetate. The values are means for 3 to 17 determinations in one to four independent experiments relative to the values obtained for the controls in the absence of acetate. The average relative standard errors of the means were 12% for *Phycomyces* and 10% for *Blakeslea*. The acetate concentration is plotted on a logarithmic scale.

TABLE 2. Effects of acetate, propionate, lactate, and leucine on the carotene contents of *Phycomyces* and *Blakeslea*^a

Medium	Carotene content (mg/g [dry mass])					
	<i>Blakeslea</i>			<i>Phycomyces</i>		
	F986	F921	F986 + F921	NRRL1554	NRRL1555	NRRL1554 + NRRL1555
Minimal	0.27	0.38	4.22	0.17	0.19	0.67
Minimal + sodium acetate (10 mmol/liter)	0.16	0.34	0.52	0.09	0.14	0.24
Minimal + potassium acetate (10 mmol/liter)	0.22	0.41	0.56	0.07	0.13	0.20
Minimal + L-leucine (10 mmol/liter)	0.45	0.69	0.30	0.22	0.26	0.29
Minimal + DL-lactate (20 mmol/liter)	0.22	0.41	1.80	0.09	0.14	0.23
Minimal + propionate (10 mmol/liter)				0.06	0.11	0.16

^a Single and mated cultures were grown on minimal medium with the compounds indicated. The values are the mean total carotene contents (β -carotene and phytoene in the case of *Phycomyces*; β -carotene, γ -carotene, lycopene, and phytoene in the case of *Blakeslea*) for at least two independent determinations. The average relative standard error of the means was 7%.

young cultures that were 1.5 to 2 days old. Production of β -carotene continued in older cultures, as observed for the phytoene content (Fig. 4) and for the very large amounts of trisporates found in mated cultures of this fungus (34). In the presence of acetate, sexual carotenogenesis occurred in young cultures (up to ~ 1 mg carotene per g [dry mass]) but not in older cultures. Phytoene was abundant, but γ -carotene, lycopene, and other carotenes collectively comprised $<10\%$ of the total carotene in mated cultures without acetate and <0.1 mg/g (dry mass) in the other cultures.

Acetate inhibition of sexual carotenogenesis in mutants. To determine if acetate must be metabolized to be inhibitory, we tested three *facA* mutants which do not utilize acetate as a sole carbon source because they lack the required acetyl-CoA syn-

thetase. To determine if acetate inhibits sexual carotenogenesis by the same mechanism that it uses to activate vegetative spores, we tested two *ger* mutants defective in spore activation by acetate, propionate, and heat (27). The results obtained with the mutants (Table 3) were indistinguishable from the results obtained with their parental strain (only 2 of the 30 pairwise comparisons between the *fac* mutants and the wild-type strains in Table 3 were significant at a *P* value of 0.05, and none was significant at a *P* value of 0.01). Mating increased the carotene content, and the increase was inhibited by acetate; both effects were significant for each of the five mutants and very significant ($P < 0.001$) for the pooled results for the three *facA* mutants and the pooled results for the two *ger* mutants.

These results were independently confirmed with single-strain cultures grown in the presence of trisporic acid extracts (~ 7 μ mol/liter). These conditions resulted in a modest increase in the carotene content (43% on average) that was eliminated by ace-




FIG. 3. Dependence of β -carotene content on acetate concentration. Single-strain and mated cultures of *Phycomyces* wild-type strains (\circ , NRRL1555; \square , NRRL1554; \blacktriangle , mated) and *Blakeslea* wild-type strains (\circ , F921; \square , F986; \blacktriangle , mated) were grown on minimal medium with various concentrations of sodium acetate. The values are means and standard errors for 2 to 15 determinations in one to four independent experiments. The acetate concentration is plotted on a logarithmic scale.




FIG. 4. Time courses for phytoene and β -carotene contents of single-strain and mated cultures of *Phycomyces* wild-type strains (\bullet and \circ , NRRL1555; \blacksquare and \square , NRRL1554; \blacktriangle and \triangle , mated) and *Blakeslea* wild-type strains (\bullet and \circ , F921; \blacksquare and \square , F986; \blacktriangle and \triangle , mated) grown on minimal medium (\circ , \square , and \triangle) or minimal medium with 10 mmol/liter sodium acetate (\bullet , \blacksquare , and \blacktriangle). The values are means and standard errors for 4 to 15 determinations in two to four independent experiments.

TABLE 3. Effect of acetate on the carotene contents of single-strain and mated cultures of various strains of *Phycomyces*^a

Medium	Carotene content (mg/g [dry mass])						
	Wild-type strains		<i>facA</i> mutants			<i>ger</i> mutants	
	NRRL1554	NRRL1555	MU136	MU138	MU141	S437	S440
Single strains							
Minimal	0.17	0.19	0.18	0.17	0.13	0.17	0.19
Minimal + sodium acetate	0.09	0.14	0.12	0.11	0.10	0.12	0.14
Minimal + trisporic acids	0.18	0.22	0.32	0.26	0.19	0.23	0.30
Minimal + trisporic acids + sodium acetate	0.12	0.11	0.18	0.12	0.09	0.11	0.12
Mated with NRRL1554							
Minimal		0.67	0.84	0.65	0.47	0.68	0.67
Minimal + sodium acetate		0.24	0.23	0.22	0.15	0.21	0.25

^a The values are the mean total carotene contents (β -carotene and phytoene) for 2 to 15 determinations in two to four independent experiments. The average relative standard error of the means was 12%. The compounds added to the minimal medium were 10 mmol/liter sodium acetate and trisporic acids from mated *Blakeslea* cultures.

tate; both effects were significant at least at a *P* value of 0.05 for the wild type, for the pooled results for the three *facA* mutants, and for the pooled results for the two *ger* mutants.

DISCUSSION

The increase in carotene biosynthesis and the induction of sexual morphogenesis that leads to zygospore production have been considered two indissoluble aspects of sexuality in the Mucorales, and both of these processes are induced by trisporic acids. Since trisporic acids are metabolites of β -carotene (2), sexual carotenogenesis was seen as part of an autocatalytic cycle for sexual stimulation. Small amounts of trisporic acid precursors produced by the vegetative hyphae reach hyphae of the opposite sex, where the compounds are converted to trisporic acids that trigger sexual morphogenesis and sexual carotenogenesis (34). The additional carotene increases trisporic acid production, and the cycle is repeated. This hypothesis is supported by the observation that mutants unable to produce carotene also do not stimulate carotene production by their partners, and the mated pair of strains does not produce zygospores (33). The link between the two sexual responses has now been broken since we observed that acetate and other chemicals stimulate the morphogenetic response but inhibit the metabolic response. Enhanced sexual morphogenesis occurred in the absence of the high carotene concentrations that usually accompany it. Our results highlight our incomplete knowledge of sexuality in the Mucorales.

Sexual carotenogenesis occurs in both *Blakeslea* and *Phycomyces*, but mated cultures of *Blakeslea* contain more carotene than mated cultures of *Phycomyces* contain. The same sexual behavior need not occur in both organisms, and the organisms differ in morphology, in the regulation of carotenogenesis by light and chemicals (3), and in the production of trisporic acids (34).

Acetate is not a sexual hormone because in single-strain cultures it does not induce the appearance of zygophores, a conspicuous early stage of sexual development. Trisporic acids do induce zygophorogenesis in single-strain cultures of many Mucorales, and this is the basis for simple functional tests for the presence of these compounds in complex media (34).

The sexual and vegetative pathways for carotenogenesis are the same in both fungi and are blocked by the same *car* mu-

tations. Four genes that encode enzymes in the biosynthetic pathway, the early *hmgS* and *hmgR* genes (31) and the late *carRA* and *carB* genes (1, 30, 32, 35), occur only once in the *Phycomyces* genome. Acetate and the other chemicals evaluated in this study cannot block a special pathway, because no such pathway exists. Instead, they inhibit sexual activation of carotenogenesis and have little or no effect on the regulatory mechanisms that are known to operate during vegetative growth (e.g., feedback inhibition).

The threshold for the effect of acetate on sexual carotenogenesis was <1 mmol/liter, a concentration that provided less than 0.3% of the available carbon atoms and should not have caused major changes in metabolism. Under the conditions used in our experiments, acetate was not metabolized to a significant extent, because while glucose is present, acetyl-CoA synthetase is rare (16). Acetate also altered carotene production in mutants that cannot use acetate as a carbon source because they have mutations in the *facA* gene for acetyl-CoA synthetase (15, 16). Thus, acetate does not need be utilized as a carbon source to alter carotene metabolism, but instead it may act as a signal that prevents sexual carotenogenesis while increasing zygospore formation.

There are at least two signal transduction pathways for acetate in *Phycomyces*, based on observations of sexual carotenogenesis and the initiation of zygospore formation following exposure to acetate of *ger* mutants, which were isolated based on their inability to activate spores in response to acetate, propionate, and heat (27). Thus, the receptor thought to be mutated in the *ger* mutants is not used for the modification of sexual responses by acetate, although the two pathways could share later steps.

The two modifications of the sexual processes differ in their dependence on the acetate concentration, and the activation of zygospore formation is less sensitive to acetate than the inhibition of sexual carotenogenesis. Thus, the two effects are mediated by transduction pathways that are at least partially different.

All the compounds that we tested can be considered small carboxylic acids; L-leucine is converted to 2-keto-4-methylpentanoic acid when it enters the cells. The human genome contains a large family of G-protein-coupled receptors (18), two of which specifically bind carboxylic acids with one to six carbon atoms (7, 19, 25).

Glutamate minimal medium is better than the standard min-

imal medium for zygospore production (33), which suggests that sexual development is favored by nitrogen starvation since the amount of nitrogen in glutamate minimal medium is <25% of the amount in the standard minimal medium. Potato dextrose medium also contains a low concentration of nitrogen and is a very good medium for zygospore production, but the low level of nitrogen is not the sole cause of this phenotype since addition of asparagine did not modify the production of zygospores (Table 1). We hypothesized that potatoes contain chemicals that could act like acetate in our experiments. This hypothesis was supported by the lack of synergy between acetate and potato-based media for zygospore production.

Our results have economic implications since fungal carotene production currently relies on sexual carotenogenesis. Media used in commercial carotene production should contain little or no acetate, L-leucine, and lactate. Propionate also should be avoided due to its strong inhibition of *Blakeslea* growth. These chemicals might not be added intentionally to media but could result from the growth of bacterial contaminants in industrial fermentors.

The activation of dormant spores by small carboxylic acids can be easily explained for a saprophyte like *Phycomyces*, which grows too slowly to compete with bacteria but could feed on bacteria. Acetate and other small carboxylic acids are products of bacterial growth on carbohydrates and amino acids and could be suitable signals for breaking the dormancy of spores of a predator of bacteria. *Blakeslea* is an opportunist on plant tissues, and its spores do not need to be activated to germinate. It is not obvious why either fungus has mechanisms to modify sexual processes in response to the same signals.

ACKNOWLEDGMENTS

We thank S. Torres-Martínez and his coworkers at Universidad de Murcia for giving us the MU *Phycomyces* strains.

This work was supported by grant INIA RM2004-12 from the Spanish Government and by grants CVI910 and CV-119 from Junta de Andalucía.

REFERENCES

- Arrach, N., R. Fernández-Martín, E. Cerdá-Olmedo, and J. Avalos. 2001. A single gene for lycopene cyclase, phytoene synthase, and regulation of carotene biosynthesis in *Phycomyces*. *Proc. Natl. Acad. Sci. USA* **98**:1687–1692.
- Austin, D. J., J. D. Bullock, and D. Darke. 1970. The biosynthesis of trisporic acids from β -carotene via retinal and trisporol. *Experientia* **26**:348–349.
- Avalos, J., and E. Cerdá-Olmedo. 2004. Fungal carotenoid production, p. 367–378. *In* D. K. Arora (ed.), *Handbook of fungal biotechnology*. Marcel Dekker, Inc., New York, N.Y.
- Bejarano, E. R., and E. Cerdá-Olmedo. 1992. Independence of the carotene and sterol pathways of *Phycomyces*. *FEBS Lett.* **306**:209–212.
- Bejarano, E. R., F. Parra, F. J. Murillo, and E. Cerdá-Olmedo. 1988. End-product regulation of carotenogenesis in *Phycomyces*. *Arch. Microbiol.* **150**:209–214.
- Blakeslee, A. F. 1904. Sexual reproduction in the Mucorineae. *Proc. Am. Acad. Arts Sci.* **40**:205–319.
- Brown, A. J., S. M. Goldsworthy, A. A. Barnes, M. M. Eilert, L. Tcheang, D. Daniels, A. I. Muir, M. J. Wigglesworth, I. Kinghorn, N. J. Fraser, N. B. Pike, J. C. Strum, K. M. Steplewski, P. R. Murdock, J. C. Holder, F. H. Marshall, P. G. Szekeres, S. Wilson, D. M. Ignar, S. M. Foord, A. Wise, and S. J. Dowell. 2003. The orphan G protein-coupled receptors GPR41 and GPR43 are activated by propionate and other short chain carboxylic acids. *J. Biol. Chem.* **278**:11312–11319.
- Caglioti, L., G. Cainelli, B. Camerino, R. Mondelli, A. Prieto, A. Quilico, T. Salvadori, and A. Selva. 1966. The structure of trisporic-C acid. *Tetrahedron* **7**(Suppl.):175–187.
- Cerdá-Olmedo, E. 1987. Standard growth conditions and variations, p. 337–339. *In* E. Cerdá-Olmedo and E. D. Lipson (ed.), *Phycomyces*. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Cerdá-Olmedo, E. 1989. Production of carotenoids with fungi, p. 27–42. *In* E. J. Vandamme (ed.), *Biotechnology of vitamins, pigments and growth factors*, Elsevier Applied Science, London, United Kingdom.
- Cerdá-Olmedo, E. 2001. *Phycomyces* and the biology of light and color. *FEMS Microbiol. Rev.* **25**:503–512.
- Chichester, C. O., T. Nakayama, G. Mackinney, and T. W. Goodwin. 1955. Incorporation of labeled leucine into carotene by *Phycomyces*. *J. Biol. Chem.* **214**:515–517.
- Ciegler, A. 1965. Microbial carotenogenesis. *Adv. Appl. Microbiol.* **7**:1–34.
- de Cima, S., J. Rúa, E. Perdiguero, P. del Valle, F. Busto, A. Baroja-Mazo, and D. de Arriaga. 2005. An acetyl-CoA synthetase not encoded by the *facA* gene is expressed under carbon starvation in *Phycomyces blakesleeanus*. *Res. Microbiol.* **156**:663–669.
- Garre, V., F. Murillo, and S. Torres-Martínez. 1994. Isolation of the *facA* (acetyl-CoA synthetase) gene of *Phycomyces blakesleeanus*. *Mol. Gen. Genet.* **244**:278–286.
- Garre, V., and S. Torres-Martínez. 1996. Mutants of *Phycomyces blakesleeanus* defective in acetyl-CoA synthetase. *Fungal Genet. Biol.* **20**:70–73.
- Govind, N. S., and E. Cerdá-Olmedo. 1986. Sexual activation of carotenogenesis in *Phycomyces blakesleeanus*. *J. Gen. Microbiol.* **132**:2775–2780.
- Im, D.-S. 2004. Discovery of new G protein-coupled receptors for lipid mediators. *J. Lipid Res.* **45**:410–418.
- Le Poul, E., C. Loison, S. Struyf, J.-Y. Springael, V. Lannoy, M.-E. Decobecq, S. Brezillon, V. Dupriez, G. Vassart, J. Van Damme, M. Parmentier, and M. Detheux. 2003. Functional characterization of human receptors for short chain fatty acids and their role in polymorphonuclear cell activation. *J. Biol. Chem.* **278**:25481–25489.
- Mehta, B. J., and E. Cerdá-Olmedo. 1995. Mutants of carotene production in *Blakeslea trispora*. *Appl. Microbiol. Biotechnol.* **42**:836–838.
- Mehta, B. J., and E. Cerdá-Olmedo. 2001. Intersexual partial diploids of *Phycomyces*. *Genetics* **158**:635–641.
- Mehta, B. J., I. N. Obratsova, and E. Cerdá-Olmedo. 2003. Mutants and intersexual heterokaryons of *Blakeslea trispora* for production of β -carotene and lycopene. *Appl. Environ. Microbiol.* **69**:4043–4048.
- Murillo, F. J., I. L. Calderón, I. López-Díaz, and E. Cerdá-Olmedo. 1978. Carotene-superproducing strains of *Phycomyces*. *Appl. Environ. Microbiol.* **36**:639–642.
- Nieuwenhuis, M., and H. van den Ende. 1975. Sex specificity of hormone synthesis in *Mucor mucedo*. *Arch. Microbiol.* **102**:167–169.
- Nilsson, N. E., K. Kotarsky, C. Owman, and B. Olde. 2003. Identification of a free fatty acid receptor, FFA2R, expressed on leukocytes and activated by short-chain fatty acids. *Biochem. Biophys. Res. Commun.* **303**:1047–1052.
- Ninet, L., and J. Renaut. 1979. Carotenoids, p. 529–544. *In* H. J. Peppler and D. Perlman (ed.), *Microbial technology*, 2nd ed., vol. 1. Academic Press, New York, N.Y.
- Rivero, F., and E. Cerdá-Olmedo. 1987. Spore activation by acetate, propionate and heat in *Phycomyces* mutants. *Mol. Gen. Genet.* **209**:149–153.
- Rivero, F., and E. Cerdá-Olmedo. 1994. Spore germination in *Phycomyces blakesleeanus*. *Mycologia* **86**:781–786.
- Robbins, W. J., V. W. Kavanagh, and F. Kavanagh. 1942. Growth substances and dormancy of spores of *Phycomyces*. *Bot. Gaz.* **104**:224–242.
- Rodríguez-Sáiz, M., B. Paz, J. L. de la Fuente, M. J. López-Nieto, W. Cabri, and J. L. Barredo. 2004. *Blakeslea trispora* genes for carotene biosynthesis. *Appl. Environ. Microbiol.* **70**:5589–5594.
- Ruiz-Albert, J., E. Cerdá-Olmedo, and L. M. Corrochano. 2002. Genes for mevalonate biosynthesis in *Phycomyces*. *Mol. Genet. Genomics* **266**:768–777.
- Ruiz-Hidalgo, M. J., E. P. Benito, G. Sandmann, and A. P. Eslava. 1997. The phytoene dehydrogenase gene of *Phycomyces*: regulation of its expression by blue light and vitamin A. *Mol. Gen. Genet.* **253**:734–744.
- Sutter, R. P. 1975. Mutations affecting sexual development in *Phycomyces blakesleeanus*. *Proc. Natl. Acad. Sci. USA* **72**:127–130.
- Sutter, R. P. 1987. Sexual development, p. 317–336. *In* E. Cerdá-Olmedo and E. D. Lipson (ed.), *Phycomyces*. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Torres-Martínez, S., F. J. Murillo, and E. Cerdá-Olmedo. 1980. Genetics of lycopene cyclization and substrate transfer in β -carotene biosynthesis in *Phycomyces*. *Genet. Res.* **36**:299–309.
- van Laere, A. J., B. Furch, and J. A. Van Assche. 1987. The sporangiospore: dormancy and germination, p. 247–279. *In* E. Cerdá-Olmedo and E. D. Lipson (ed.), *Phycomyces*. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- van Laere, A. J., and F. Rivero. 1986. Properties of a germination mutant of *Phycomyces blakesleeanus*. *Arch. Microbiol.* **145**:290–294.
- van Mulders, R. M., A. J. Van Laere, and M. N. Verbeke. 1986. Effects pH and cations on the germination induction of *Phycomyces* spores with carboxylic acids. *Biochem. Physiol. Pflanz.* **181**:103–115.