

Pathotyping *Escherichia coli* by Using Miniaturized DNA Microarrays^{∇†}

Muna F. Anjum,^{1*} Muriel Mafura,¹ Peter Slickers,² Karin Ballmer,³ Peter Kuhnert,³
 Martin J. Woodward,¹ and Ralf Ehricht²

Department of Food and Environmental Safety, Veterinary Laboratories Agency-Weybridge, New Haw, Addlestone, Surrey KT15 3NB, United Kingdom¹; CLONDIAG Chip Technologies GmbH, Jena, Germany²; and Institute of Veterinary Bacteriology, Vetsuisse Faculty of the University of Bern, Bern, Switzerland³

Received 22 February 2007/Accepted 4 July 2007

The detection of virulence determinants harbored by pathogenic *Escherichia coli* is important for establishing the pathotype responsible for infection. A sensitive and specific miniaturized virulence microarray containing 60 oligonucleotide probes was developed. It detected six *E. coli* pathotypes and will be suitable in the future for high-throughput use.

Pathogenic *Escherichia coli* strains constitute a significant public health problem worldwide (12). In contrast to their nonpathogenic counterparts, these strains have acquired specific virulence attributes that allow them to cause a spectrum of human and animal illnesses (10, 15). Numerous methods exist for the detection of pathogenic *E. coli*, including geno- and phenotypic marker assays for the detection of virulence genes and their products (7, 17, 21, 23). These methods have the common drawback of screening a relatively small number of determinants simultaneously. DNA microarrays offer a viable alternative due to their ability to screen multiple markers simultaneously.

The aim of this work was to develop a simple high-throughput system based in a microtube (details are available from CLONDIAG, Jena, Germany) (13, 20) for pathotyping *E. coli* isolates sent to clinical diagnostic laboratories.

Design and validation of miniaturized virulence arrays. A miniaturized *E. coli* oligonucleotide virulence array was designed containing 39 virulence, 7 bacteriocin, and 15 control (*rhl* and *gad*) gene probes (Table 1). Eighteen genes were specific to a particular *E. coli* pathotype, 13 were common between 2 or more pathotypes, and 7 were unassigned. The design of probes/primers and the specificity were tested as previously described (1, 13).

Control strains were used to validate each probe present on the array (Table 1). PCR amplification and sequencing, using primers given in Appendix 1 of the supplemental material, verified the presence of the probes in control strains. The sequenced genes showed between 92 and 100% sequence identity to the respective target gene and showed 100% sequence identity to the probe and primer regions (data not shown).

Genomic DNA was extracted from cells grown aerobically overnight at 37°C in LB broth, using a DNeasy tissue kit (catalog no. 69504; QIAGEN). One microgram of genomic DNA

from each strain was used as a template in a multiplex linear amplification and labeling reaction with the set of 60 primers (Table 1), as previously described (1). The amplified products were added to ArrayTubes for hybridizations performed according to the method of Ballmer et al. (1, 13).

The sequenced strains EDL933, CFT073, and E2348/69 were used to estimate assay sensitivity to ensure strong signal intensity with minimal nonspecific cross-hybridization. Optimization included varying the concentrations of genomic DNA used for labeling (2 to 0.05 µg), the primers present in the linear multiplex mix (0.135 to 0.810 µM), and the poly-horse-radish peroxidase–streptavidin conjugate used for detecting hybridization (50 to 400 pg/µl). The minimal concentration of genomic DNA found to reliably detect all expected genes was 1.0 µg, while a concentration of 0.135 µM per primer in the stock solution was sufficient for the detection of target DNA (Fig. 1). The optimal concentration of poly-horseradish peroxidase–streptavidin conjugate was found to be 200 pg/µl; concentrations above or below this value resulted in high background or no detectable reaction at all (data not shown).

The spot signal intensity was derived by calculating the quantitative staining value with IconoClust software (version 2; CLONDIAG). The data were normalized using the signal intensity of the *gad* probe, and the normalized signal intensity for genes within positive and negative control strains was used to differentiate between present (signal intensity value above 0.4) and absent (signal intensity value below 0.3) genes. Genes with signal intensity values between 0.3 and 0.4 were considered ambiguous. Two replicate hybridizations were performed for each control strain, and the 95% confidence interval of error across replicate hybridizations was 1.6 to 3% (see Appendix 2 in the supplemental material).

The specificity of each probe was estimated by comparing array data with PCR and sequenced data from control strains. In all cases, the virulence gene(s) known to be present within positive control strains was clearly identified by array, while two negative control strains, including the sequenced strain MG1655, showed the presence of only 23S rRNA and *gad* genes (see Appendix 2 in the supplemental material). For many positive control strains, additional virulence genes were detected (Table 2). Furthermore, PCR amplification in all

* Corresponding author. Mailing address: Department of Food and Environmental Safety, Veterinary Laboratories Agency-Weybridge, New Haw, Addlestone, Surrey KT15 3NB, United Kingdom. Phone: 44 (0)1932 359476. Fax: 44 (0)1932 357268. E-mail: m.anjum@vla.defra.gsi.gov.uk.

† Supplemental material for this article may be found at <http://aem.asm.org/>.

∇ Published ahead of print on 13 July 2007.

TABLE 1. Probes and primers used in the miniaturized microarray^a

Probe/gene	Probe/gene function	Target gene accession no.	Pathotype(s) ^b	Control strain (origin or reference) ^c	Probe sequence (5'-3')	Primer sequence (5'-3')
<i>astA</i> astA_11 astA_21	Heat-stable enterotoxin	AE005345.1	EAEC, ETEC	Abbotstown (22)	TCgTgCATATGGTGCgCA ACAG TCgTgCATATGGTGCgCA ACAG	GACGGCTTTGTAgTCC TTCC TGACGGCTTTGTatTC CTTCC
<i>bfpA</i>	Major subunit of bundle-forming pili	AB024946.1	EPEC	E2348/69 (16)	GGTGTGATGTTTTACTAC CAGTCTGC	CGCTCATTACTTCTGA AATaGCA
<i>cba</i>	Colicin B pore forming	M16816.1	Undesignated	EC2334/03 (VLA)	GGATGGTCTGTcAGTGTG CACG	GCGGAAACTTTCTCGT TTCC
<i>cdtB</i> cdtB_40 cdtB_50	Cytolethal distending toxin B	AJ508930.1	EPEC, STEC, ETEC, ExPEC*	EC934/04 (VLA)	GCTGTGATGCcTtGGT GGAAG GCTGTGATGCcTtGGT GGAAG	GCTAACCAGAGCAAGA TTGAC
<i>celB</i>	Endonuclease colicin E2	X03632.1	Undesignated	EC2334/03 (VLA)	GGACCGTATCTCCGTCAT CAACAG	GCCTTGCTAATCCGG TCAC
<i>cfA</i>	Colonization factor antigen I	M55661.1	ETEC	IMI100 (Bern)	GGAAATAGCCCGCTGGGTA TTACAGA	TCATCCACCAATTTAA GACAGC
<i>cma</i>	Colicin M, resembles beta-lactam antibiotics	M16754.1	Undesignated	EC2334/03 (VLA)	TGTAACGCCGACCGAAAT CTGGT	TCATAAACGCTTATTC CAGGGT
<i>cnf</i>	Cytotoxic necrotizing factor	AF483828.1	ExPEC*	S5 (8)	CTTCCAGTATGGGGATCA GTTTTGATCA	CGACGTTCTCATAAG TATCACC
<i>eae^c</i> eae_10 eae_20 eae_30 eae_40 ^b	Intimin	AJ579371.1	EPEC, EHEC	E2348/69 (25)	GTTACAaCaTTATGGAAC GGCAGAGGT GTTACAaCgTTATGGAAC GGCAGAGG TGGTgAtAATACCCGtTT AGGtATtGGt TGGTgAtAATACCCGcTT AGGtATtGG	CgTCAAAGTTATtACC aCTCTGC AGTcTCGCCAgTAT TCgC
<i>f17A^c</i> f17A_40 ^b f17A_50 ^b f17A_60 ^b	Subunit A of F17 fimbrial protein	AF022140.1	ETEC, ExPEC*	CK210 (6), S5 (8)	ggTAcTAtGCaACgGgtc aGGC CagTAcTAcGCaACgGgt gtGG aCaaTAtTAtGcCaCaGc gccGG	TGATAAgCGATGGTGT AATTCaCaG TGATAAgCGATGGTGT AATTaActG CTGATAAaCGATGGTG TAATTTActG
<i>f17G</i>	Adhesion subunit of F17 fimbrial protein	AF022140.1	ETEC, ExPEC*	CK210 (6)	TGCAATGGATAACCTGCC ATTTGTCT	CCAGACATTTGCATTC TGATATCC
<i>fanA</i>	Involved in biogenesis of K99/F5 fimbriae	X05797.1	ETEC	ETEC562 (VLA)	AGCAAGGTGCTTCCAAT ATTAGTGGG	CGTAAATACCCCTAGA ACTACGT
<i>fasA</i>	Fimbrial 987P/F6 subunit	M35257.1	ETEC	HM1535 (VLA)	GCCAAGTGGATACTTCTA ATCTGTGCG	GAGCAGAAGTAGACAA CTCTCC
<i>fim41a</i>	Mature Fim41a/F41 protein	X14354.1	ETEC	ETEC562 (VLA)	GGCTTGTTPAATCCAGGTC GATTTACTG	GAGAGTCCATTCATT TATAGGCT
<i>gad</i>	Glutamate decarboxylase	M84025.1	All <i>E. coli</i>	All	GATATCGTCTGGGACTTC CGCCT	TGAAGCACTGATCGAT TTCACA
<i>ehx (hlyA)</i>	Hemolysin A	AB011549.2	EPEC, EHEC	EDL933 (19)	TGTAGGATTAActGAAcG TGGTGTTCG	GCAGAAGTTTGTCAAG TTGTGG
<i>hlyE</i>	Avian <i>E. coli</i> hemolysin	AF052225.1	ExPEC*	M1000 (14)	CCAAGATAGACTTTCGA GGCGACAC	TCACTCCACACCATT ATAAACT
<i>ipaH9.8</i>	Invasion plasmid antigen	AF047365.1	<i>Shigella sonnei</i>	NCTC8192 (HPA)**	TCGCCCTCACATGGAACA ATCTC	GCCTGATGGACCAG GAGG
<i>ireA</i>	Siderophore receptor	AF320691.1	ExPEC*	CFT073 (24)	CCACAAATGACTTCTATC TGTcAGGC	CTCCATATAGCTGAAG ACCAAGT
<i>iroN</i>	Enterobactin siderophore receptor protein	AF449498.1	ExPEC*	CFT073 (24)	GCCTGTGAGTAACATGA TCAATGCT	GAGGCTTTGCGAAGT GAGC
<i>iss</i>	Increased serum survival	AF042279.1	ExPEC*	CFT073 (24)	CCGCTCTGGCAATGCTTA TTACAGG	gGTTTGTtTccAACAG TAAACGT
<i>K88ab</i>	K88/F4 protein subunit gene	V00292.1	ETEC	Abbotstown (22)	GCCTGGATGACTGGTGAT TTCAATGG	GTGATACTACCACCGA TATCGAC
<i>lngA</i>	Longus type IV pilus	AF004308.1	ETEC	B1308 (VLA)	CGTCTGGTTCATATGCCA TGACAGC	CCACAGACATATCTAC ACCAAGT
<i>lthA</i>	Heat-labile enterotoxin A subunit	AB011677.1	ETEC	ETEC21d (VLA)	GGTTTCTGCGTTAGGTGG AATACCA	ACCAAATTAACACGA TACCATCC
<i>mchB</i>	Microcin H47 part of colicin H	AJ515252.1	Undesignated	CFT073 (24)	GGTTGTAGTTGAGCCGT ATCTGC	GGTCGAGCCAATTG CTGT
<i>mchC</i>	MchC protein	AJ515252.1	Undesignated	CFT073 (24)	CTGTCCGGTTAGATCTGT GATCCAC	CCGGTGGTACAGGTAG ATATCC
<i>mchF</i>	ABC transporter protein MchF	AJ515251.1	Undesignated	CFT073 (24)	TCCGGTTATTCATCAGAC GGAGACC	CAAAAATGACCGCATAT CATTCG
<i>mcmA</i>	Microcin M part of colicin H	AJ515251.1	Undesignated	CFT073 (24)	CCTCCATGTCTCCCTCAG GTATAGG	GGCACTTGATGTACCT CTGC

Continued on following page

TABLE 1—Continued

Probe/gene	Probe/gene function	Target gene accession no.	Pathotype(s) ^b	Control strain (origin or reference) ^c	Probe sequence (5'–3')	Primer sequence (5'–3')
<i>perA</i> ^c	EPEC adherence factor, transcriptional activator					
perA_10		AF255772.1	EPEC	E2348/69 (16)	TGTTTGGTTGGGTTTAAAT TCCACATCA	TTGGTGTGTGTGTGTA ATATTCTT
perA_20				N1743-95 (Bern)	GCTTGGTTGGTTTAAAT CCACGTC	
<i>pet</i>	Autotransporter enterotoxin	AF056581.1	EAEC	NZ1470-95 (Bern)	GCTGACAAGGATAATCT GCCACAAGA	GCATCGCGAGAGCA AACT
<i>prfB/papB</i>	P-related fimbriae regulatory gene	X76613.1	ExPEC*	CFT073 (24)	GGGAGACTTATACGGCTG AATGCTC	TCATCTGTATAATAAG GTGCAAGC
<i>senB</i>	Plasmid-encoded enterotoxin	Z54195.1	EIEC	NCTC9774 (HPA)	GCTCTATATCGGACACAC CCAGTCAG	GGTGTCAAACATACTG ATACGC
<i>sfaS</i>	S fimbria minor subunit	X16664.4	ExPEC*	E536 (VLA)	CAATGCAGGAAGTGGATC TCCATGG	TCCGGTGAGAGACAG ATCA
<i>sta1A</i> ^c	Heat-stable enterotoxin ST-Ia					
sta1A_111		AJ555214.1	ETEC	ETEC562 (VLA)	ACACATTTTACTGCTGTG AACTTTGTTG	AACATgAGCACAG GCAG
sta1A_121						AACATccAGCACAG GCAG
<i>sta1B</i>	Heat-stable enterotoxin ST-Ib	AY342058	ETEC	IMI100 (Bern)	AGCAATTACTGCTGTGAA TTGTGTTGT	AGCACCCGGTACAA GCAG
<i>stb</i>	Heat-stable enterotoxin II	AJ555214.1	ETEC	Abbotstown (22)	GAGATGGTACTGCTGGAG CATGCT	TTGCTGCAACCATTAT TTGGG
<i>stx1A</i>	Shiga toxin 1 A subunit	AB035142.1	STEC	EDL933 (19)	GTGACAGTAGCTATACCA CGTTACAGC	TCTGCATCCCCGTA CGAC
<i>stx2A</i>	Shiga toxin 2 A subunit	AB035143.1	STEC	EDL933 (19)	GCAGTTATACCACCTCTGC AACGTTGTC	CtgAttTGCATtCCgG aACG
<i>virF</i>	VirF transcriptional activator, <i>ipaBCD</i> -positive regulator	AF386526.1	<i>Shigella flexneri</i>	NCTC8192 (HPA)**	GCCTTTTATCAGCTGTTT CTGATGAGGA	GAGAAGAAGCTATCGA TATCGAAGT
rrl_0101_0177_10	23S rRNA (large rRNA)	M25458.1	All	E2348/69 ^d	GTGTGTTTCGACACACTA TCATTAACCTGA	GGTTCGCCTCATTAAC CTATGG
rrl_0101_0177_20	23S rRNA (large rRNA)	M25458.1	All	E2348/69 ^d	GTGTGATTTCGTCACACTA TCATTAACCTGA	
rrl_0260_0330_10	23S rRNA (large rRNA)	M25458.1	All	E2348/69 ^d	CAGAGCCTGAATCAGTAT GTGTGTTAGT	GCCTTTCCAGACGC TTCC
rrl_0260_0330_20	23S rRNA (large rRNA)	M25458.1	All	E2348/69 ^d	GAGCCTGAATCAGTGTGT GTGTTAGT	
rrl_0260_0330_30	23S rRNA (large rRNA)	M25458.1	All	E2348/69 ^d	AGAGCCTGAATCAGTTTG TGTGTTAGT	
rrl_0520_0580_10	23S rRNA (large rRNA)	M25458.1	All	E2348/69 ^d	GCAGTGGGAGCACGCT TAGG	AAGGTACGCAGTCA CACG
rrl_0520_0580_20	23S rRNA (large rRNA)	M25458.1	All	E2348/69 ^d	AAGCAGTGGGAGCATGCT TAGG	
rrl_1480_1560_coli_10	23S rRNA (large rRNA)	M25458.1	All	E2348/69 ^d	CCGAAAAATCAAGGATGA GGCGTG	CACCGTAGTGCCCTC GTCA
rrl_1480_1560_coli_20	23S rRNA (large rRNA)	M25458.1	All	E2348/69 ^d	CGAAAAATCAAGGCTGAG GCGTG	
rrl_1480_1560_coli_30	23S rRNA (large rRNA)	M25458.1	All	E2348/69 ^d	GGAAAAACAAGGCTGAGG CGTG2	
rrl_1480_1560_shig_40	23S rRNA (large rRNA)	M25458.1	All	E2348/69 ^d	GGAAAAATCAAGCCGAGG CGTG	
rrl_1690_1770_coli_10	23S rRNA (large rRNA)	M25458.1	All	E2348/69 ^d	GCTGATATGTAGGTGAAG CGACTTGC	CGACTGATTTAGCTC CACG
rrl_1690_1770_freu_30	23S rRNA (large rRNA)	M25458.1	All	E2348/69 ^d	CGCTGATATGTAGGTGAA GTGGTTTACT	
rrl_1690_1770_shig_20	23S rRNA (large rRNA)	M25458.1	All	E2348/69 ^d	GCTGATACGTAGGTGAAG CGACTTGC	

^a All probes and primers present in the array representing genes or encompassing allelic variations are listed. The description for each gene, the accession number of the target gene used initially for probe/primer design, the pathotype associated with each gene, and the positive control strain are also given. Probes for the 23S rRNA gene (*rfl*) were included as a species marker, while the *gad* gene was included as an invariant positive control present in low copy number in all *E. coli* strains. *, uropathogenic *E. coli*, avian pathogenic *E. coli*, and neonatal meningitis *E. coli* have been classed together as extraintestinal pathogenic *E. coli* (ExPEC) for this study; **, Health Protection Agency, National Culture Typing Collection. Lowercase letters in sequences indicate sequence variability within the consensus region within which the probe or primer was designed.

^b EAEC, enteroaggregative *E. coli*; ETEC, enterotoxigenic *E. coli*; EPEC, enteropathogenic *E. coli*; STEC, shigatoxigenic *E. coli*; EHEC, enterohemorrhagic *E. coli*; EIEC, enteroinvasive *E. coli*. "All" indicates all *E. coli*.

^c Polymorphic genes where different control strains were found to bind to different probe sets or probes showed different signal intensities reflecting allelic variation that had not been distinguished by PCR (see the supplemental material for details).

^d For details, see www.sanger.ac.uk.

^e HPA, Health Protection Agency; VLA, Veterinary Laboratories Agency.

control strains of five randomly chosen genes (*eae*, *astA*, *ehx* or *hlyA*, *iss*, and *mcmA*), showed 100% correlation between array and PCR data, indicating the probes to be highly specific with minimum cross-reactions (data not shown).

Pathotyping clinical isolates. A panel of 63 *E. coli* human and animal clinical isolates were pathotyped using the virulence miniaturized microarray (see Appendix 3 in the supplemental material). For five strains, two hybridization reactions




FIG. 1. Optimization of the genomic concentration used in this study. The optimal concentration of genomic DNA from EDL933 used for the detection of genes on the virulence oligonucleotide miniaturized microarray chip was assessed using (a) 2 μg , (b) 1 μg , (c) 0.5 μg , (d) 0.1 μg , and (e) 0.05 μg of DNA. The six biotinylated marker spots (C) are visible in all arrays. A concentration of 0.135 μM per primer in the stock solution and 200 $\text{pg}/\mu\text{l}$ of poly-horseradish peroxidase–streptavidin conjugate were used for these assays.

were performed and the 95% confidence interval of error between replicates was 0.9 to 5.0%. Only one hybridization reaction was performed for the remaining test strains.

Fifty-five of the isolates hybridized to more than one virulence determinant and were readily designated within a recognized pathotype, mostly matching the clinical diagnosis where available. Five isolates that harbored only the *iss* gene and/or microcins and three isolates that hybridized to only control genes could not be pathotyped. These isolates may harbor virulence genes not present on our array. Several isolates with novel combinations of genes were detected and included two shigatoxigenic *E. coli* strains, one with *senB*, *iss*, *cma*, *cba*, and *mchBCF* genes and another with *astA*, *cdtB*, and *cnf* genes. The most commonly detected gene was *iss*, which was present in half the strains tested. Other genes which were detected in at least 10 or more isolates included *eae*, *ehx*, *astA*, *iroN*, *mchF*, *mchB*, *mchC*, *fl7A* (three variants combined), *fl7G*, *mcmA*, *cba*, *cma*, and *prfB/papB*. Genes *virF*, *pet*, *hlyE*, *fasA*, and *cfa* were not detected in any test isolate (see Appendix 3 in the supplemental material).

Conclusion. Several *E. coli* virulence arrays for genotyping have been described previously (2–5, 9, 11, 18). These arrays use mostly a glass slide printed with oligonucleotide probes or PCR products for target genes and fluorescent Cy dyes to label DNA used for hybridization. This system is time consuming, with expensive reagents and requires a skilled technician. In contrast, the microtube-based array system used in this study has a short assay time due to an amplification step and inexpensive reagents and requires low technical skills, making it amenable for use in clinical diagnostic laboratories. In the

future, the routine use of virulence microarrays in such laboratories will not only allow rapid detection and designation of the pathotypes of strains sent to diagnostic laboratories but also enable emergent strains harboring novel virulence combinations to be detected before such strains spread to become a health problem.

We are grateful to the Enteric Reference Laboratory at VLA for the provision of *E. coli* strains and in particular to Katherine Spriggs and Louise Finch. We thank Elke Müller and Jana Sachtschal for their assistance.

This project was funded through the VLA seedcorn fund.

REFERENCES

- Ballmer, K., B. M. Korczak, P. Kuhnert, P. Slickers, R. Ehricht, and H. Hachler. 2007. Fast DNA serotyping of *Escherichia coli* by use of an oligonucleotide microarray. *J. Clin. Microbiol.* **45**:370–379.
- Bruant, G., C. Maynard, S. Bekal, I. Gaucher, L. Masson, R. Brousseau, and J. Harel. 2006. Development and validation of an oligonucleotide microarray for detection of multiple virulence and antimicrobial resistance genes in *Escherichia coli*. *Appl. Environ. Microbiol.* **72**:3780–3784.
- Call, D. R., F. J. Brockman, and D. P. Chandler. 2001. Detecting and genotyping *Escherichia coli* O157:H7 using multiplexed PCR and nucleic acid microarrays. *Int. J. Food Microbiol.* **67**:71–80.
- Chen, S., S. Zhao, P. F. McDermott, C. M. Schroeder, D. G. White, and J. Meng. 2005. A DNA microarray for identification of virulence and antimicrobial resistance genes in *Salmonella* serovars and *Escherichia coli*. *Mol. Cell. Probes* **19**:195–201.
- Chizhikov, V., A. Rasooly, K. Chumakov, and D. D. Levy. 2001. Microarray analysis of microbial virulence factors. *Appl. Environ. Microbiol.* **67**:3258–3263.
- Cid, D., R. Sanz, I. Marin, H. de Greve, J. A. Ruiz-Santa-Quiteria, R. Amils, and R. de la Fuente. 1999. Characterization of nonenterotoxigenic *Escherichia coli* strains producing F17 fimbriae isolated from diarrheic lambs and goat kids. *J. Clin. Microbiol.* **37**:1370–1375.
- Clark, C. G., S. T. Johnson, R. H. Easy, J. L. Campbell, and F. G. Rodgers. 2002. PCR for detection of *cdt-III* and the relative frequencies of cytolethal

- distending toxin variant-producing *Escherichia coli* isolates from humans and cattle. *J. Clin. Microbiol.* **40**:2671–2674.
8. **De Rycke, J., E. A. Gonzalez, J. Blanco, E. Oswald, M. Blanco, and R. Boivin.** 1990. Evidence for two types of cytotoxic necrotizing factor in human and animal clinical isolates of *Escherichia coli*. *J. Clin. Microbiol.* **28**:694–699.
 9. **Jenkins, C., C. van Ijperen, E. G. Dudley, H. Chart, G. A. Willshaw, T. Cheasty, H. R. Smith, and J. P. Nataro.** 2005. Use of a microarray to assess the distribution of plasmid and chromosomal virulence genes in strains of enteroaggregative *Escherichia coli*. *FEMS Microbiol. Lett.* **253**:119–124.
 10. **Kaper, J. B., J. P. Nataro, and H. L. Mobley.** 2004. Pathogenic *Escherichia coli*. *Nat. Rev. Microbiol.* **2**:123–140.
 11. **Korczak, B., J. Frey, J. Schrenzel, G. Pluschke, R. Pfister, R. Ehricht, and P. Kuhnert.** 2005. Use of diagnostic microarrays for determination of virulence gene patterns of *Escherichia coli* K1, a major cause of neonatal meningitis. *J. Clin. Microbiol.* **43**:1024–1031.
 12. **Mead, P. S., L. Slutsker, V. Dietz, L. F. McCaig, J. S. Bresee, C. Shapiro, P. M. Griffin, and R. V. Tauxe.** 1999. Food-related illness and death in the United States. *Emerg. Infect. Dis.* **5**:607–625.
 13. **Monecke, S., and R. Ehricht.** 2005. Rapid genotyping of methicillin-resistant *Staphylococcus aureus* (MRSA) isolates using miniaturised oligonucleotide arrays. *Clin. Microbiol. Infect.* **11**:825–833.
 14. **Nagai, S., T. Yagihashi, and A. Ishihama.** 1998. An avian pathogenic *Escherichia coli* strain produces a hemolysin, the expression of which is dependent on cyclic AMP receptor protein gene function. *Vet. Microbiol.* **60**:227–238.
 15. **Nataro, J. P., and J. B. Kaper.** 1998. Diarrheagenic *Escherichia coli*. *Clin. Microbiol. Rev.* **11**:142–201.
 16. **Okeke, I. N., J. A. Borneman, S. Shin, J. L. Mellies, L. E. Quinn, and J. B. Kaper.** 2001. Comparative sequence analysis of the plasmid-encoded regulator of enteropathogenic *Escherichia coli* strains. *Infect. Immun.* **69**:5553–5564.
 17. **Osek, J.** 2003. Detection of the enteroaggregative *Escherichia coli* heat-stable enterotoxin 1 (EAST1) gene and its relationship with fimbrial and enterotoxin markers in *E. coli* isolates from pigs with diarrhoea. *Vet. Microbiol.* **91**:65–72.
 18. **Palaniappan, R. U., Y. Zhang, D. Chiu, A. Torres, C. Debroy, T. S. Whittam, and Y. F. Chang.** 2006. Differentiation of *Escherichia coli* pathotypes by oligonucleotide spotted array. *J. Clin. Microbiol.* **44**:1495–1501.
 19. **Perna, N. T., G. Plunkett III, V. Burland, B. Mau, J. D. Glasner, D. J. Rose, G. F. Mayhew, P. S. Evans, J. Gregor, H. A. Kirkpatrick, G. Posfai, J. Hackett, S. Klink, A. Boutin, Y. Shao, L. Miller, E. J. Grothbeck, N. W. Davis, A. Lim, E. T. Dimalanta, K. D. Potamouis, J. Apodaca, T. S. Anantharaman, J. Lin, G. Yen, D. C. Schwartz, R. A. Welch, and F. R. Blattner.** 2001. Genome sequence of enterohaemorrhagic *Escherichia coli* O157:H7. *Nature* **409**:529–533.
 20. **Perreten, V., L. Vorlet-Fawer, P. Slickers, R. Ehricht, P. Kuhnert, and J. Frey.** 2005. Microarray-based detection of 90 antibiotic resistance genes of gram-positive bacteria. *J. Clin. Microbiol.* **43**:2291–2302.
 21. **Sharma, V. K.** 2002. Detection and quantitation of enterohemorrhagic *Escherichia coli* O157, O111, and O26 in beef and bovine feces by real-time polymerase chain reaction. *J. Food Prot.* **65**:1371–1380.
 22. **Thorns, C. J., C. D. Boarer, and J. A. Morris.** 1987. Production and evaluation of monoclonal antibodies directed against the K88 fimbrial adhesin produced by *Escherichia coli* enterotoxigenic for piglets. *Res. Vet. Sci.* **43**:233–238.
 23. **Wang, G., C. G. Clark, and F. G. Rodgers.** 2002. Detection in *Escherichia coli* of the genes encoding the major virulence factors, the genes defining the O157:H7 serotype, and components of the type 2 Shiga toxin family by multiplex PCR. *J. Clin. Microbiol.* **40**:3613–3619.
 24. **Welch, R. A., V. Burland, G. Plunkett III, P. Redford, P. Roesch, D. Rasko, E. L. Buckles, S. R. Liou, A. Boutin, J. Hackett, D. Stroud, G. F. Mayhew, D. J. Rose, S. Zhou, D. C. Schwartz, N. T. Perna, H. L. Mobley, M. S. Sonnenberg, and F. R. Blattner.** 2002. Extensive mosaic structure revealed by the complete genome sequence of uropathogenic *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **99**:17020–17024.
 25. **Zhu, C., T. S. Agin, S. J. Elliott, L. A. Johnson, T. E. Thate, J. B. Kaper, and E. C. Boedeker.** 2001. Complete nucleotide sequence and analysis of the locus of enterocyte effacement from rabbit diarrheagenic *Escherichia coli* RDEC-1. *Infect. Immun.* **69**:2107–2115.