

Letters to the Editor

Multiply Infected Vectors

Wielinga et al. (10) provide an important reminder that ticks and other insect vectors are sometimes microbial “zoos” rather than single-pathogen vehicles. A single tick may simultaneously transmit *Borrelia* and *Babesia* species (7) or, perhaps, coinfect humans with *Borrelia*, *Babesia*, and *Ehrlichia* species (1, 3).

Wielinga et al. report that the frequency of *Ixodes ricinus* ticks coinfecting with *Borrelia* and *Ehrlichia* or *Anaplasma* species is higher than would be expected on the basis of independence between the pathogens, that this frequency increases with tick age, and that the degree of the statistical surplus of coinfections varies with habitat type. Given that the striking surpluses over the 2 to 3 years in the park, forest, and heather habitats ($P < 0.01$ for each by χ^2 and Fisher’s exact tests) appeared to be absent over the 5 years in the dune habitat ($P > 0.1$), might these differences between the habitats have been mediated by differences between the years in the age structures of the tick populations in the field or the assay samples? Since an individual vector rarely clears an infection, infections accumulate with age, and in a vector population, frequencies of multiply infected vectors typically track seasonal and other environmental influences on age structure (4, 5).

Wielinga et al. also point out that pathogens sharing a vector may interact within that vector. Among mosquito-borne pathogens, for instance, transmission of the Rift Valley fever and eastern equine encephalitis viruses by *Aedes* mosquitoes coinfecting with *Brugia* species is enhanced (8, 9); *Plasmodium* and *Wuchereria* species coinfect *Anopheles punctulatus* at higher-than-expected frequencies (2) but suppress each other’s devel-

opment in *Anopheles gambiae* (6). Wielinga et al. have drawn attention to an important but neglected set of topics for research, the results of which should improve our understanding and control of vector-borne diseases.

REFERENCES

1. **Belongia, E. A.** 2002. Epidemiology and impact of coinfections acquired from *Ixodes* ticks. *Vector Borne Zoonotic Dis.* **2**:265–273.
2. **Burkot, T. R., L. Molineaux, P. M. Graves, R. Paru, D. Battistutta, H. Dagoro, A. Barnes, R. A. Wirtz, and P. Garner.** 1990. The prevalence of naturally acquired multiple infections of *Wuchereria bancrofti* and human malaria in anophelines. *Parasitology* **100**:369–375.
3. **Magnarelli, L. A., J. W. Ijdo, J. F. Anderson, S. J. Padula, R. A. Flavell, and E. Fikrig.** 1998. Human exposure to a granulocytic ehrlichia and other tick-borne agents in Connecticut. *J. Clin. Microbiol.* **36**:2823–2827.
4. **McKenzie, F. E., and W. H. Bossert.** 1997. Mixed-species *Plasmodium* infections of *Anopheles* (Diptera: Culicidae). *J. Med. Entomol.* **34**:417–425.
5. **McKenzie, F. E., G. F. Killeen, J. C. Beier, and W. H. Bossert.** 2001. Seasonality, parasite diversity and local extinctions in *Plasmodium falciparum* malaria. *Ecology* **82**:2673–2681.
6. **Muirhead-Thomson, R. C.** 1953. Inter-relations between filarial and malarial infection in *Anopheles gambiae*. *Nature* **172**:352–353.
7. **Piesman, J., T. C. Hicks, R. J. Sinsky, and G. Obiri.** 1987. Simultaneous transmission of *Borrelia burgdorferi* and *Babesia microti* by individual nymphal *Ixodes dammini* ticks. *J. Clin. Microbiol.* **25**:2012–2013.
8. **Turell, M. J., P. A. Rossignol, A. Spielman, C. A. Rossi, and C. L. Bailey.** 1984. Enhanced arboviral transmission by mosquitoes that concurrently ingested microfilariae. *Science* **225**:1039–1041.
9. **Vaughan, J. A., and M. J. Turell.** 1996. Dual host infections: enhanced infectivity of eastern equine encephalitis virus to *Aedes* mosquitoes mediated by *Brugia* microfilariae. *Am. J. Trop. Med. Hyg.* **54**:105–109.
10. **Wielinga, P. R., C. Gaasenbeek, M. Fonville, A. de Boer, A. de Vries, W. Dimmers, G. A. O. Jagers, L. M. Schouls, F. Borgsteede, and J. W. van der Giessen.** 2006. Longitudinal analysis of tick densities and *Borrelia*, *Anaplasma*, and *Ehrlichia* infections of *Ixodes ricinus* ticks in different habitat areas in The Netherlands. *Appl. Environ. Microbiol.* **72**:7594–7601.

F. E. McKenzie
Fogarty International Center
National Institutes of Health
Bethesda, Maryland

Ed. Note: The authors of the published article declined to respond.