

Transfer of Conjugative Elements from Rumen and Human *Firmicutes* Bacteria to *Roseburia inulinivorans*[∇]

Karen P. Scott,* Jenny C. Martin, Jakub Mrazek,† and Harry J. Flint

Rowett Research Institute, Bucksburn, Aberdeen AB21 9SB, United Kingdom

Received 12 December 2007/Accepted 22 April 2008

Studies on *Firmicutes* bacteria from the gut are hampered by a lack of gene transfer systems. Here the human colonic anaerobe *Roseburia inulinivorans* A2-194 was shown to be a transfer recipient for two conjugative transposons, Tn1545 from *Eubacterium cellulosolvens* and TnK10 from *Clostridium saccharolyticum* K10.

The two most abundant bacterial phyla that inhabit the human colon are the low-G+C-content gram-positive *Firmicutes* and the gram-negative *Bacteroidetes* (13, 25). Recent progress in understanding the ecological roles of gram-positive *Firmicutes* in the human large intestine (4, 11) may soon be supplemented by draft genome sequences. Bacteria related to *Roseburia* spp. and *Eubacterium rectale* are an important group of *Firmicutes* belonging to the *Clostridium coccooides* cluster (also known as clostridial cluster XIVa) that contribute to polysaccharide breakdown in the colon and are major producers of butyrate (1, 10). This group can comprise up to 10% of the total bacterial diversity in the human colon (1). The species *Roseburia inulinivorans* is metabolically versatile, utilizing inulin, starch, and fucose for growth (9, 19, 22). While progress in understanding gene expression responses in *R. inulinivorans* using transcriptomics is being made (22), genetic manipulation methods would clearly enhance the ability to investigate gene function. Some progress has been made with the introduction of transposon and plasmid vectors in the rumen species *Butyrivibrio fibrisolvens* (5, 14, 20), but very little work with related gram-positive strict anaerobes from the human gut has been reported.

Commensal bacteria are known to harbor antibiotic resistance genes and thus contribute to the reservoir of resistance genes in the gut that can potentially be acquired by pathogenic members of the gut microbial community. Several new tetracycline resistance genes have been identified recently in gram-positive anaerobes isolated from the bovine rumen and human colon (3, 16). The transfer of some of these resistance genes was linked to their carriage on conjugative transposons (CTNs), and these elements may offer the potential for genetic manipulation. TnB1230, originally identified in the rumen anaerobe *B. fibrisolvens* (20), is a 45- to 50-kb transposon with some similarity to the *Enterococcus faecalis* transposon Tn1549 (17). Repeated attempts to transfer TnB1230 from *B. fibrisolvens* into *R. inulinivorans* were unsuccessful despite the high fre-

quencies of transfer between *B. fibrisolvens* strains observed previously (20). We have, however, been able to transfer another novel CTN, TnK10, previously identified in a human isolate related to *Clostridium saccharolyticum* (16), into *R. inulinivorans* A2-194, and transfer with Tn1545, originally identified in *Streptococcus pneumoniae*, was also successful (8). Details of these transfer experiments are presented below.

Transfer of the CTN Tn1545. Tn1545, which carries the *tet(M)* gene, was transferable from the rumen *Eubacterium cellulosolvens* strain 5484 to a spontaneous rifampin-resistant mutant of *R. inulinivorans* A2-194 (A2-194^R), at frequencies of 10⁻⁵ transconjugants per donor cell. Anaerobic filter matings followed the method of Hespell and Whitehead (14), modified as described previously (20). Parent cells were grown anaerobically in M2GSC medium lacking antibiotics overnight to stationary phase prior to mating. Potential transconjugants grew on selective M2GSC agar plates containing rifampin (100 μg ml⁻¹) and tetracycline (10 μg ml⁻¹) for 2 days, and their resistance phenotypes were confirmed by replating selected colonies on double-antibiotic plates. Subsequent stocks of transconjugants were prepared from single colonies. The 16S rRNA gene from each transconjugant was amplified by PCR using eubacterial primers fD1 and rP2 (28), and limited sequencing of the amplicons confirmed that transconjugants were *R. inulinivorans* isolates.

Chromosomal DNA isolated from transconjugants was subjected to restriction enzyme digestion, separated by pulsed-field gel electrophoresis (PFGE), Southern blotted, and hybridized to a *tet(M)* probe as described previously (3, 20). Single insertion of Tn1545 was predicted to give one hybridizing fragment following restriction with HindIII [which cleaves 63 nucleotides into *tet(M)* but not within the probe sequence] and two hybridizing fragments following cleavage at the ClaI site, located at 913 nucleotides in *tet(M)*, within the probe sequence. Six different complex hybridization profiles were observed with 12 *R. inulinivorans* transconjugants when the ClaI and HindIII hybridization patterns were compared (Fig. 1, lanes 2, 4, 6, 9, 11, and 12). The lower band (~2.5 kb) present in all transconjugants is due to the presence of an additional ClaI site upstream of *tet(M)*, within Tn1545 (Fig. 1a). Tn1545 was shown previously to insert into different sites in five *E. cellulosolvens* transconjugants (2) and to target AT-rich regions in the genome (15, 27).

DNA sequences flanking the left end of Tn1545 were ob-

* Corresponding author. Mailing address: Rowett Research Institute, Greenburn Road, Bucksburn, Aberdeen, Scotland AB21 9SB, United Kingdom. Phone: 44 (0) 1224 712751. Fax: 44 (0) 1224 716687. E-mail: k.scott@rowett.ac.uk.

† Present address: Institute of Animal Physiology and Genetics, Czech Academy of Sciences, Videnska 1083, 142 20, Prague 4, Czech Republic.

[∇] Published ahead of print on 2 May 2008.




FIG. 1. Southern blot of genomic DNA purified from *R. inulinivorans* A2-194 transconjugants containing Tn1545, digested with *Cla*I (a) or *Hind*III (b). Lanes 1 to 12, selected transconjugants; lane 13, recipient strain. The sizes of key marker bands are indicated on the left. Unique patterns are represented by transconjugants in lanes 2, 4, 6, 9, 11, and 12.

tained for seven transconjugants with six distinct banding patterns. Chromosomal DNA was partially digested with *Sau*3A, and the 0.5- to 1-kb fraction was ligated into a pBluescript vector (Stratagene) and amplified with M13 specific primers in combination with primers located ~100 bp from each end of Tn1545 based on the sequence published by Caillaud and Courvalin (6) (Tn1545left, 5' AGTCATTCATAAGTAGT 3'; Tn1545right, 5' CGTGAAGTATCTTCCTACAG 3'). The resulting amplicons were purified and sequenced directly using primer Tn1545left. All seven sequences were identical up to and following the left end of Tn1545 (6), and the same sequence was also obtained for the donor *E. cellulosolvens* strain. The transferred DNA therefore included DNA in addition to Tn1545 that was presumably derived from *E. cellulosolvens*. The most plausible explanation is that Tn1545 became part of a larger conjugal transfer element upon integration into the *E. cellulosolvens* donor strain and that this larger element is able to insert at more than one site upon transfer to *R. inulinivorans*.

Transfer of the CTn TnK10. The human gut isolate *C. saccharolyticum* K10 harbors the transmissible tetracycline resistance gene *tet*(O/32/O) and also a *tet*(W) gene (16, 24). Transfer of *tet*(O/32/O) only was detected in matings with the rumen recipient strain *B. fibrisolvens* 2221^R (16). *tet*(O/32/O) was therefore postulated to be carried on the CTn TnK10, which had a common insertion site in two transconjugants analyzed (16). Since the two bacteria capable of harboring TnK10 belong to clostridial cluster XIVa, this transposon was considered a promising candidate for further experiments on transfer to *R. inulinivorans* A2-194.

Transfer of tetracycline resistance occurred between *C. saccharolyticum* K10 and the recipient *R. inulinivorans* A2-194^R, at frequencies of 10⁻⁶ transconjugants per donor cell. Gene transfer was confirmed by PCR amplification of the *tet*(O/32/O) gene using a primer pair specific for the internal *tet*(32) region (18) [*tet*(32)for, 5'-AACCGAAGCATACCGCTC-3'; *tet*(32)rev, 5'-CTC




FIG. 2. Southern blot of a representative PFGE gel hybridized to the internal *tet*(32) sequence. R, recipient *R. inulinivorans* A2-194; D, donor *C. saccharolyticum* K10; lanes 1 to 5, representative transconjugants containing TnK10. The sizes of key bands are indicated on the right.

TTTCATAGCCACGCC-3']. We were unable to estimate the size of TnK10 based on PFGE analysis (data not shown). However, sequencing of homologous elements indicated that TnK10 is likely to be larger than 21 kb (EMBL database accession number AM419751; M. Rincon, personal communication). The band profile on a PFGE gel was immobilized by Southern blotting, and the blot was hybridized to a radioactively labeled *tet*(O/32/O) probe. The restriction enzyme *Sal*I does not cut internally in *tet*(O/32/O), and a single insertion of TnK10 should result in a single hybridizing band. A 90-kb band hybridized to the *tet*(O/32/O) probe in all five of the transconjugants analyzed, with an additional 60-kb band present in two transconjugants (Fig. 2, lanes 3 and 5). This implies that TnK10 has a preferred insertion site in *R. inulinivorans* but that insertions into a second site can also occur. A much larger single band of about 200 kb hybridized in the donor genome but was absent in the transconjugants.

Conclusions. The rumen *B. fibrisolvens* and *E. cellulosolvens* and human *R. inulinivorans* isolates studied here all belong to clostridial cluster XIVa (7), the single most abundant phylogenetic group of bacteria resident in both the human colon (25) and the rumen (26). Through this and previous work (21), we have shown that transfer of antibiotic resistance genes occurs in both directions between rumen and human gut representatives of this important group of gut *Firmicutes*. Previous evidence has shown that rumen and human gut representatives of the major gram-negative phylum, the *Bacteroidetes*, can also exchange resistance genes (23). Transfer of TnK10 and Tn1545 between unrelated gut anaerobes was easily detected, but repeated attempts to transfer the rumen CTn TnB1230 to *R. inulinivorans* A2-194 were unsuccessful. Interestingly, TnB1230 transfers at extremely high frequencies between two phylogenetically distinct strains of *B. fibrisolvens* (12, 20), and the lack of transfer to *R. inulinivorans* A2-194 presumably reflects the restricted host range of TnB1230. In fact, the donor, *B. fibrisolvens* 1.230, is phylogenetically more closely related to *Roseburia* species than to the recipient, *B. fibrisolvens* 2221^R (4).

Genetic manipulation of bacteria within clostridial cluster XIVa would enhance the ability to investigate the function of specific genes, and it was hoped that one of the CTns studied here would enable transposon mutagenesis in this group. TnK10 showed preferential insertion in *R. inulinivorans* but

remains of interest as a potential delivery system, once more sequence information is available. Although Tn1545 apparently had multiple insertion sites in the *R. inulinivorans* genome, this was inferred to be due to its transfer as part of a larger element, whose structure and behavior require further investigation. Genomic sequencing of the donor and recipient bacteria may prove the most effective approach to clarifying the transfer mechanisms of Tn1545 and TnK10.

In conclusion, this work demonstrates conjugal gene transfer in laboratory matings with a strictly anaerobic *Firmicutes* bacterium from the human gastrointestinal tract as the recipient, thus hopefully providing a first step toward genetic manipulation in this important group of bacteria. These experiments emphasize the extent of gene flow between predominant groups of gut bacteria.

Nucleotide sequence accession number. The accession number for the 16S rRNA sequence of *C. saccharolyticum* K10 is EU305624.

Jakub Mrazek was supported by a grant from the EU Marie Curie training site (ANAEROBE). The Rowett Research Institute is supported by SG-RERAD (Scottish Government Rural and Environment Research and Analysis Directorate).

We thank Kevin Anderson for kindly supplying *E. cellulosolvans* strains carrying Tn1545.

REFERENCES

- Aminov, R. I., A. W. Walker, S. H. Duncan, H. J. M. Harmsen, G. W. Welling, and H. J. Flint. 2006. Molecular diversity, cultivation, and improved detection by fluorescent in situ hybridization of a dominant group of human gut bacteria related to *Roseburia* spp. or *Eubacterium rectale*. *Appl. Environ. Microbiol.* **72**:6371–6376.
- Anderson, K. L., J. A. Megehee, and V. H. Varel. 1998. Conjugal transfer of transposon Tn1545 into the cellulolytic bacterium *Eubacterium cellulosolvans*. *Lett. Appl. Microbiol.* **26**:35–37.
- Barbosa, T. M., K. P. Scott, and H. J. Flint. 1999. Evidence for recent intergeneric transfer of a new tetracycline resistance gene, *tet(W)*, isolated from *Butyrivibrio fibrisolvans*, and the occurrence of *tet(O)* in ruminant bacteria. *Environ. Microbiol.* **1**:53–64.
- Barcenilla, A., S. E. Pryde, J. C. Martin, S. H. Duncan, C. S. Stewart, C. Henderson, and H. J. Flint. 2000. Phylogenetic relationships of butyrate-producing bacteria from the human gut. *Appl. Environ. Microbiol.* **66**:1654–1661.
- Beard, C. E., M. A. Hefford, R. J. Forster, S. Sontakke, R. M. Teather, and K. Gregg. 1995. A stable and efficient transformation system for *Butyrivibrio fibrisolvans* OB156. *Curr. Microbiol.* **30**:105–109.
- Caillaud, F., and P. Courvalin. 1987. Nucleotide sequence of the ends of the conjugative shuttle transposon Tn1545. *Mol. Gen. Genet.* **209**:110–115.
- Collins, M. D., P. A. Lawson, A. Willems, J. J. Cordoba, J. Fernandez-Garayzabal, P. Garcia, J. Cai, H. Hippe, and J. A. E. Farrow. 1994. The phylogeny of the genus *Clostridium*: proposal of five new genera and eleven new species combinations. *Int. J. Syst. Bacteriol.* **44**:812–826.
- Courvalin, P., and C. Carlier. 1986. Transposable multiple antibiotic resistance in *Streptococcus pneumoniae*. *Mol. Gen. Genet.* **205**:291–297.
- Duncan, S. H., K. P. Scott, A. G. Ramsay, H. J. Harmsen, G. W. Welling, C. S. Stewart, and H. J. Flint. 2003. Effects of alternative dietary substrates on competition between human colonic bacteria in an anaerobic fermentor system. *Appl. Environ. Microbiol.* **69**:1136–1142.
- Duncan, S. H., R. I. Aminov, K. P. Scott, P. Louis, T. B. Stanton, and H. J. Flint. 2006. Proposal of *Roseburia faecis* (sp. nov.), *Roseburia hominis* (sp. nov.), and *Roseburia inulinivorans*, based on isolates from human faeces. *Int. J. Syst. Evol. Microbiol.* **56**:2437–2441.
- Duncan, S. H., P. Louis, and H. J. Flint. 2007. Cultivable bacterial diversity from the human colon. *Lett. Appl. Microbiol.* **44**:343–350.
- Forster, R. J., R. M. Teather, J. Gong, and S.-J. Deng. 1996. 16S rDNA analysis of *Butyrivibrio fibrisolvans*: phylogenetic position and relation to butyrate producing anaerobic bacteria from the rumen of white-tailed deer. *Lett. Appl. Microbiol.* **23**:218–222.
- Franks, A. H., H. J. Harmsen, G. C. Raangs, G. J. Jansen, F. Schut, and G. W. Welling. 1998. Variations of bacterial populations in human feces measured by fluorescent in situ hybridization with group-specific 16S rRNA-targeted oligonucleotide probes. *Appl. Environ. Microbiol.* **64**:3336–3345.
- Hespell, R. B., and T. R. Whitehead. 1991. Conjugal transfer of Tn916, Tn916ΔE, and pAMB1 from *Enterococcus faecalis* to *Butyrivibrio fibrisolvans* strains. *Appl. Environ. Microbiol.* **57**:2703–2709.
- Jaworski, D. D., and D. B. Clewell. 1994. Evidence that coupling sequences play a frequency determining role in conjugative transposition of Tn916 in *Enterococcus faecalis*. *J. Bacteriol.* **176**:3328–3335.
- Melville, C. M., K. P. Scott, D. K. Mercer, and H. J. Flint. 2001. Novel tetracycline resistance gene, *tet(32)*, in the *Clostridium*-related human colonic anaerobe K10 and its transmission in vitro to the rumen anaerobe *Butyrivibrio fibrisolvans*. *Antimicrob. Agents Chemother.* **45**:3246–3249.
- Melville, C. M., R. Brunel, H. J. Flint, and K. P. Scott. 2004. The *Butyrivibrio fibrisolvans tet(W)* gene is carried on the novel conjugative transposon TnB1230, which contains duplicated nitroreductase coding sequences. *J. Bacteriol.* **186**:3656–3659.
- Patterson, A. J., R. Colangeli, P. Spigaglia, and K. P. Scott. 2007. Distribution of specific tetracycline and erythromycin resistance genes in environmental samples assessed by microarray detection. *Environ. Microbiol.* **9**:703–715.
- Ramsay, A. G., K. P. Scott, J. C. Martin, M. T. Rincon, and H. J. Flint. 2006. Cell associated α -amylases of butyrate-producing Firmicute bacteria from the human colon. *Microbiology* **152**:3281–3290.
- Scott, K. P., T. M. Barbosa, K. J. Forbes, and H. J. Flint. 1997. High frequency transfer of a naturally occurring chromosomal tetracycline resistance element in the ruminal anaerobe *Butyrivibrio fibrisolvans*. *Appl. Environ. Microbiol.* **63**:3405–3411.
- Scott, K. P. 2002. The role of conjugative transposons in spreading antibiotic resistance between bacteria that inhabit the gastrointestinal tract. *Cell. Mol. Life Sci.* **59**:2071–2082.
- Scott, K. P., J. C. Martin, G. Campbell, C.-D. Mayer, and H. J. Flint. 2006. Whole-genome transcription profiling reveals genes up-regulated by growth on fucose in the human gut bacterium *Roseburia inulinivorans*. *J. Bacteriol.* **188**:4340–4349.
- Shoemaker, N. B., H. Vlamakis, K. Hayes, and A. A. Salyers. 2001. Evidence for extensive resistance gene transfer among *Bacteroides* spp. and among *Bacteroides* and other genera in the human colon. *Appl. Environ. Microbiol.* **67**:561–568.
- Stanton, T. B., S. B. Humphrey, K. P. Scott, and H. J. Flint. 2005. Hybrid *tet* genes and *tet* gene nomenclature: request for opinions. *Antimicrob. Agents Chemother.* **49**:1265–1266.
- Suau, A., R. Bonnet, M. Sutren, J.-J. Godon, G. R. Gibson, M. D. Collins, and J. Doré. 1999. Direct analysis of genes encoding 16S rRNA from complex communities reveals many molecular species within the human gut. *Appl. Environ. Microbiol.* **65**:4799–4807.
- Tajima, K., R. I. Aminov, T. Nagamine, K. Ogata, M. Nakamura, H. Matsui, and Y. Benno. 1999. Rumen bacterial diversity as determined by sequence analysis of 16S rDNA libraries. *FEMS Microbiol. Ecol.* **29**:159–169.
- Trieu-Cuot, P., C. Poyart-Salmeron, C. Carlier, and P. Courvalin. 1993. Sequence requirements for target activity in site-specific recombination mediated by the Int protein of transposon Tn1545. *Mol. Microbiol.* **8**:179–185.
- Weisburg, W. G., S. M. Barns, D. A. Pelletier, and D. J. Lane. 1991. 16S ribosomal DNA amplification for phylogenetic study. *J. Bacteriol.* **173**:697–703.