

# Fermentation pH Influences the Physiological-State Dynamics of *Lactobacillus bulgaricus* CFL1 during pH-Controlled Culture<sup>∇</sup>

Aline Rault,<sup>1,2</sup> Marielle Bouix,<sup>1</sup> and Catherine Béal<sup>1\*</sup>

AgroParisTech–INRA–UMR 782 Génie et Microbiologie des Procédés Alimentaires, CBAI, 78850 Thiverval-Grignon, France,<sup>1</sup> and ARILAIT Recherches, 42 Rue de Châteaudun, 75314 Paris Cedex 09, France<sup>2</sup>

Received 28 November 2008/Accepted 29 April 2009

**This study aims at better understanding the effects of fermentation pH and harvesting time on *Lactobacillus bulgaricus* CFL1 cellular state in order to improve knowledge of the dynamics of the physiological state and to better manage starter production. The Cinac system and multiparametric flow cytometry were used to characterize and compare the progress of the physiological events that occurred during pH 6 and pH 5 controlled cultures. Acidification activity, membrane damage, enzymatic activity, cellular depolarization, intracellular pH, and pH gradient were determined and compared during growing conditions. Strong differences in the time course of viability, membrane integrity, and acidification activity were displayed between pH 6 and pH 5 cultures. As a main result, the pH 5 control during fermentation allowed the cells to maintain a more robust physiological state, with high viability and stable acidification activity throughout growth, in opposition to a viability decrease and fluctuation of activity at pH 6. This result was mainly explained by differences in lactate concentration in the culture medium and in pH gradient value. The elevated content of the ionic lactate form at high pH values damaged membrane integrity that led to a viability decrease. In contrast, the high pH gradient observed throughout pH 5 cultures was associated with an increased energetic level that helped the cells maintain their physiological state. Such results may benefit industrial starter producers and fermented-product manufacturers by allowing them to better control the quality of their starters, before freezing or before using them for food fermentation.**

Lactic acid bacteria are traditionally used to produce or to preserve various food products such as fermented milks, meats, and vegetables. Their ability to initiate rapid acidification of the raw material is essential to improve the flavor, texture, and safety of these products (11, 14). In order to prevent poor fermentation yields and to improve the quality and reliability of the products, it is important to maintain proper control starter production. This control may be achieved by studying the effects of process parameters on the growth kinetics of the bacteria and on their acidification activity and physiological state in growing conditions. Among all process parameters, pH and harvesting time are key factors that strongly influence the physiological state of lactic acid bacteria after fermentation and stabilization.

Lactic acid starters are currently produced using pH-controlled pure cultures (6), during which pH is generally regulated at an optimal value by continuously adding sodium hydroxide or ammonia in the bioreactor (23). Various growth characteristics such as maximal biomass concentration, specific growth rate, fermentation time, sugar consumption or growth, and product yields are significantly influenced by the pH control value (1, 4). Optimal pH ranges were therefore determined for several lactic acid bacteria, such as *Streptococcus thermophilus* (pH 6.5), *Lactobacillus bulgaricus* (pH 5.8 to 6) (5, 22), or *Lactococcus lactis* subsp. *cremoris* (pH 6.3 to 6.9) (8).

Compared to acidic fermentations, pH-controlled cultures led to higher growth yields and productivity (9, 23) as a result

of the lower level of nondissociated lactic acid in the culture medium (2, 12, 15). The acidification of the cytoplasm induced by the nondissociated form of the weak organic acid leads to the collapse of the proton motive force (13). This phenomenon inhibits nutrient transport and enzymatic reactions and leads to DNA alteration and biomass inactivation (12). Maintaining the extracellular pH (pH<sub>ext</sub>) at a high value helps the cells stabilize their intracellular pH at a sufficiently high value (9), thus decreasing the inhibiting effect of lactic acid.

Fermentation pH also acts on energetic parameters, such as internal pH (pH<sub>i</sub>), pH gradient (dpH), proton motive force, membrane potential, NADH/NAD ratio, ATP level and rate of ATP formation, and lactate dehydrogenase and ATPase activity (1, 9, 17). During batch cultures of *L. lactis* performed with or without pH control, Cachon et al. (9) showed that pH control has a significant influence on the variations of pH<sub>i</sub>, dpH, and NADH/NAD ratio, thus acting on growth parameters. Moreover, in batch cultures, pH<sub>i</sub> is dependent upon both the external pH and the age of culture. Mercade et al. (17) showed that cultures of *L. bulgaricus* at controlled pH 6.4 are inhibited at the level of anabolism but were not energy limited. They are characterized by a high maintenance coefficient in contrast to cultures without pH control which consume intracellular energy for pH<sub>i</sub> regulation.

The effect of pH on cellular physiology is confirmed by other studies which show that it influences acidification activity of lactic acid bacteria (23–25). Whereas Wang et al. (25) indicated that *Lactobacillus acidophilus* cells grown at optimal pH display a higher residual acidification activity than cells grown at lower pH control values, Schepers et al. (24) and Savoie et al. (23) demonstrated that this activity is higher when starters are produced without pH control or at low pH control values. These authors explained that conditions generating high bio-

\* Corresponding author. Mailing address: AgroParisTech, INRA, UMR 782 Génie et Microbiologie des Procédés Alimentaires, CBAI, F-78850 Thiverval-Grignon, France. Phone: 33(0)1 30 81 54 85. Fax: 33(0)1 30 81 55 97. E-mail: beal@grignon.inra.fr.

<sup>∇</sup> Published ahead of print on 8 May 2009.

mass concentrations do not systematically lead to cells with an efficient acidification activity.

From this information, the effect of pH control was elucidated on growth and energetic parameters, whereas its effect on the dynamic of cellular physiology, viability, and acidification activity during growth is still not determined.

A few authors demonstrated that the harvesting time has a strong impact on cellular parameters such as viability and acidification activity (3, 20, 24). Béal et al. (6) specified that there is an optimal range of time during which to harvest cells in a good physiological state, i.e., at a high cellular concentration and a high acidification activity. However, since this optimal range is strongly strain and condition dependent, more information is needed about the influence of harvesting time on physiological parameters.

In order to improve knowledge about the effects of fermentation pH and harvesting time on starter's quality, we sought here to apply some rapid and relevant methods to characterize the dynamic of *L. delbrueckii* subsp. *bulgaricus* CFL1 physiological state throughout pH 6 and pH 5 fermentations. This might allow industrial starter producers to better control their fermentations and to achieve high-quality starters. Among the available methods, the Cinac system and multiparametric flow cytometry, associated with plate counts, made it possible to determine and compare different physiological parameters such as cultivability, acidification activity (Cinac system), membrane damage, enzymatic activity, cell depolarization, intracellular pH, and pH gradient (flow cytometry) (20). Two dynamic schemes of the time course of the physiological state during pH 6 or pH 5 cultures are proposed and discussed.

## MATERIALS AND METHODS

**Bacterial strain and preculture conditions.** *Lactobacillus delbrueckii* subsp. *bulgaricus* CFL1 (INRA, Thiverval-Grignon, France) was used throughout the present study. Frozen cells were stored at  $-80^{\circ}\text{C}$  in MRS broth (AES-Chemunex, Combourg, France) supplemented with 15% glycerol. Before inoculation in the bioreactor, cells were subcultured twice, for 24 h and 8.5 h, at  $42^{\circ}\text{C}$  in 5 ml of MRS broth.

**Fermentation conditions.** The culture medium was composed of 60 g liter $^{-1}$  sweet whey powder (Eurosérum, Port-sur-Saône, France), which was adjusted to pH 5 with 50%  $\text{H}_2\text{SO}_4$ , heat treated at  $110^{\circ}\text{C}$  for 20 min and centrifuged at  $7,000 \times g$  for 30 min at  $4^{\circ}\text{C}$ . After filtration (0.45- $\mu\text{m}$  pore size), it was supplemented with 20 g liter $^{-1}$  yeast extract (Organotechnie, La Courneuve, France) and introduced into a 2-liter bioreactor (Setric Génie Industriel, Toulouse, France). After sterilization at  $110^{\circ}\text{C}$  for 20 min, the initial pH was adjusted at pH 6, and inoculation was done at a low initial level of  $20 \pm 5.5$  CFU ml $^{-1}$  to allow overnight culture. Fermentations were performed at  $42^{\circ}\text{C}$ , 100 rpm, and with pH 6 or the pH 5 control, by automatic adding of a 5 N sodium hydroxide (NaOH) solution in the bioreactor. The NaOH concentration (m, in g liter $^{-1}$ ) and its consumption rate (dm/dt, in g liter $^{-1}$  min $^{-1}$ ) were calculated with WCidus software (INRA, Thiverval-Grignon, France).

The fermentation kinetics were therefore established, and the reproducibility was assessed according to three descriptors: the maximal rate of NaOH consumption ( $V_m$ , in g liter $^{-1}$  min $^{-1}$ ), the time necessary to reach  $V_m$  ( $t_m$ , in min), and the NaOH consumption (dNaOH, in g liter $^{-1}$ ) at  $t_m$ . Samples were taken from the bioreactor at different sampling times, from the beginning of NaOH consumption up to late stationary phase. In order to normalize the data obtained from the different cultures, results were expressed as a function of the time  $t_m$  necessary to reach  $V_m$ , which was considered as a reference time.

**HPLC measurements.** Lactose, glucose, galactose, and lactic acid concentrations were quantified by using high-performance liquid chromatography (HPLC; Waters Associates, Millipore, Molsheim, France). Before the HPLC analyses, each sample was combined with 120 g of trichloroacetic acid (Prolabo, Paris, France) liter $^{-1}$ , centrifuged at  $4,000 \times g$  for 30 min at  $4^{\circ}\text{C}$ , and filtered (0.22- $\mu\text{m}$  pore size). The HPLC analysis was made on a cation-exchange column (Aminex

Ion Exclusion HPX-87H 300\*7.8 mm; Bio-Rad, Richmond, CA) at  $35^{\circ}\text{C}$  with propionic acid (10 g liter $^{-1}$ ) as an internal standard. The mobile phase was 0.005 M  $\text{H}_2\text{SO}_4$ , and the flow rate was set at 0.6 ml min $^{-1}$  (LC-6A pump; Shimadzu, Courtabouef, France).

The dissociated lactate and nondissociated lactic acid concentrations,  $[A^-]$  and  $[AH]$ , respectively, were evaluated as a function of the dissociation constant of lactic acid ( $\text{pK}_a = 3.86$ ), the pH, and total lactic acid concentration,  $[LA]$ , as measured by HPLC. They were calculated by considering the following formulas:

$$[A^-] / [AH] = 10^{(\text{pH} - \text{pK}_a)} \quad (1)$$

$$[A^-] + [AH] = [LA] \quad (2)$$

**Measurement of cultivability.** Cultivability was evaluated by plate counts. After serial dilutions in peptone water (Bacto peptone, 1 mg liter $^{-1}$ ; Difco Laboratories, Le Pont-de-Claix, France), cells were plated onto solid MRS agar (AES-Chemunex) and incubated at  $42^{\circ}\text{C}$  for 48 h under anaerobic conditions (Genbox 96124; bioMérieux, Marcy l'Etoile, France). Each result (N, in CFU ml $^{-1}$ ) was the geometrical mean of at least three counts.

**Fluorescent probes and staining protocols.** (i) **Viability and mortality assessment.** Carboxyfluorescein diacetate (cFDA) was used to assess *L. bulgaricus* CFL1 viability according to the method of Rault et al. (19). The nucleic acid dye propidium iodide (PI) made it possible to quantify damaged and dead cells. Live/dead assays were done by dual staining of each sample to differentiate viable, dead, and damaged cells. Before staining, cell suspensions were diluted in Chemsol B13 buffer (AES-Chemunex) to reach  $10^6$  cells ml $^{-1}$ . A 1-ml portion of the diluted suspension was first supplemented with 10  $\mu\text{l}$  of PI (1.496 mM in distilled water; Sigma-Aldrich, Lyon, France) and incubated for 20 min at  $40^{\circ}\text{C}$ . Then, 10  $\mu\text{l}$  of cFDA (0.217  $\mu\text{M}$  in acetone, Invitrogen-Molecular Probes, Eragny-sur-Oise, France) was added, and incubation took place for 10 min before direct analysis by flow cytometry.

(ii) **Depolarization assessment.** Bis(1,3-dibutylbarbituric acid) trimethine oxonol [DIBAC $_4$ (3)] was used to assess the depolarization state of the cells (7). Dual staining was performed in order to differentiate dead cells with PI and depolarized cells with DIBAC $_4$ (3) in a single analysis. Then, 1 ml of a bacterial suspension was diluted in Chemsol B13 buffer (AES-Chemunex) to reach  $10^6$  cells ml $^{-1}$  and incubated with 10  $\mu\text{l}$  of PI (1.496 mM in distilled water; Sigma-Aldrich) for 10 min at  $40^{\circ}\text{C}$ . Then, 5  $\mu\text{l}$  of DIBAC $_4$ (3) (96.78  $\mu\text{M}$  in dimethyl sulfoxide; Invitrogen-Molecular Probes) was added, and incubation took place for 20 min, before direct analysis by flow cytometry.

(iii) **pHi measurement.** The fluorochrome 5-(6)-carboxyfluorescein diacetate succinimidyl ester (cFDA-SE) was used for pHi measurements (21). The method was first calibrated by using pH-equilibrated cells at known pHi (20). Cell suspensions were diluted in a pH 7.3 Mac Ilvaine buffer (0.1 M citric acid [Fisher Chemical, Elancourt, France]; 0.2 M disodium dihydrogenophosphate [J. T. Baker, Deventer, The Netherlands]) to reach  $10^7$  cells ml $^{-1}$ . Next, 1 ml of the diluted suspension was incubated with 5  $\mu\text{l}$  of cFDA-SE (8.96  $\mu\text{M}$  in acetone; Invitrogen/Molecular Probes) for 10 min at  $40^{\circ}\text{C}$ . After staining, 100  $\mu\text{l}$  of cell suspension were diluted in 1 ml of McIlvaine buffers ranging from pH 4 to pH 7.5. Intracellular pH was equilibrated with buffer pH by addition of valinomycin and nigericin (Sigma Aldrich) to a final concentration of 1  $\mu\text{M}$  each. After 1 min of incubation at  $25^{\circ}\text{C}$ , the fluorescence intensity was measured by flow cytometry.

A calibration curve was established by plotting the logarithm of the mean fluorescence intensity as a function of pH of equilibrated cells, in the range of pH 4 to pH 7.5, as previously described by Rault et al. (20).

Analyses of intracellular pH of samples were carried out according to the same protocol. After staining with cFDA-SE, 100  $\mu\text{l}$  of cell suspension were diluted in 1 ml of a buffer adjusted to the external pH of the cells. The fluorescence intensity was measured with flow cytometry, and the linear calibration curve made it possible to quantify the intracellular pH (20).

(iv) **Flow cytometry and data analyses.** Flow cytometry analyses were performed with a Cyflow cytometer (AES-Chemunex) equipped with a specific volumetric counting system, an air-cooled argon ion laser emitting at 488 nm, and four band-pass filters: a forward-angle light scatter (FSC) combined with a diode collector, a side-angle light scatter (SSC), and two fluorescence signals collected with photomultiplier tubes. A 530-nm band-pass filter (515 to 545 nm) was used to collect the green fluorescence of carboxy fluorescein or DIBAC $_4$ (3) (FL1 channel), and a 630-nm long-pass filter was used to collect the red fluorescence of PI (FL2 channel). The FACSFlow solution (Becton Dickinson, Le Pont-de-Claix, France) was used as sheath fluid. Flow cytometry analyses were performed by using logarithmic gains and specific detector settings adjusted on a sample with unstained cells in order to eliminate cellular autofluorescence. A combination of FSC and SSC was used to discriminate the bacteria from the background.

TABLE 1. Kinetic parameters characterizing fermentations of *L. bulgaricus* CFL1 conducted at controlled pH 6 or pH 5

Parameter <sup>a</sup>	Mean (95% confidence interval) <sup>b</sup>	
	pH 6	pH 5
V <sub>m</sub> (g liter <sup>-1</sup> min <sup>-1</sup> )	-0.32 (0.03)	-0.13 (0.02)
t <sub>m</sub> (min)	806 (23)	1,045 (34)
dNaOH (g liter <sup>-1</sup> )	29.1 (1.5)	23.8 (4.9)
dlact (g liter <sup>-1</sup> )	35.2 (0.8)	32.4 (2.1)
dgal (g liter <sup>-1</sup> )	19.2 (0.7)	16.9 (0.5)
dLA (g liter <sup>-1</sup> )	17.2 (0.4)	16.2 (0.5)

<sup>a</sup> dlact, total lactose consumption; dgal, total galactose production; dLA, total lactic acid production. Other parameters are as defined in the text.

<sup>b</sup> Values are means of at least three measurements, with the corresponding 95% confidence intervals in parentheses.

Data were collected and analyzed with Flowmax software (Partec, Sainte-Geneviève-des-Bois, France). Subpopulations were identified by using dot plots of FSC, SSC, green fluorescence, and red fluorescence, which separated the different events. The data were analyzed with the aid of statistical tables given by the Flowmax software. It indicated numbers (in cells ml<sup>-1</sup>) and percentages of stained cells determined by each detector, along with the mean fluorescence intensity of each fluorescent signal, and the associated coefficients of variation. Each result corresponded to the mean of at least six measurements.

(v) **Specific acidification activity measurements.** The Cinac system (Ysebaert, Frépillon, France) was used to assess the acidification activity of the cells (10). Acidification was measured in triplicate at 42°C in 100 g of dry skim milk (EPI-Ingredient, Ancenis, France) liter<sup>-1</sup>, heat treated at 110°C for 20 min in 150-ml flasks, and stored at 4°C before use. From a previous determination of viable cell concentrations by cFDA staining and flow cytometry, the flasks were

inoculated at an initial concentration of 10<sup>5</sup> viable cells ml<sup>-1</sup>. The pH was continuously measured by the Cinac system and led to the determination of the time necessary to reach pH 5.5 (tpH5.5, in min) as a descriptor to characterize the acidification activity of bacterial suspensions. The higher the tpH5.5 is, the lower the acidification activity.

## RESULTS

**Fermentation kinetics.** Table 1 summarizes the main kinetic parameters characterizing the fermentations performed at pH 6 and pH 5. Since the 95% confidence intervals were low compared to the mean values of the parameters, it was considered that the reproducibility of the cultures was satisfactory, thus making it possible to analyze and compare the physiological characteristics of the cells obtained from the different batch cultures at each controlled pH.

As shown on Table 1, the fermentation kinetic was slower at pH 5 than at pH 6, with a difference of ~4 h for t<sub>m</sub>. This difference was related to the lower maximal rate of NaOH consumption (V<sub>m</sub>) observed at pH 5. If the other kinetic parameters and their confidence intervals are considered, pH 5 cultures do not appear to produce significant differences from pH 6 cultures in terms of lactose consumption, galactose and lactic acid production or NaOH consumption.

In order to normalize the data obtained from the six cultures, results were expressed as a function of the reference time t<sub>m</sub> (in min) necessary to reach the maximal rate of NaOH consumption (Fig. 1a and Fig. 2a). However, since




FIG. 1. Time course of *L. bulgaricus* CFL1 physiological characteristics throughout fermentations performed at pH 6. (a) NaOH consumption (—, g liter<sup>-1</sup>) and rate of NaOH consumption (dNaOH/dt, - - -, g liter<sup>-1</sup> min<sup>-1</sup>); (b) cultivability (Δ, CFU ml<sup>-1</sup>) and relative percentages of viable (●, %), dead (○, %), damaged (◐, %), and depolarized (+, %) cells; (c) specific acidification activity (tpH5.5, ▲, min); (d) intracellular pH (■), pHex (□), and pH gradient (dpH = pHi - pHex) (▣). Results are the means of at least three independent measurements. They are expressed as a function of the time t<sub>m</sub> to reach the maximal rate of NaOH consumption as an adjusted time.




FIG. 2. Time course of *L. bulgaricus* CFL1 physiological characteristics throughout fermentations performed at pH 5. (a) NaOH consumption (—, g liter<sup>-1</sup>) and rate of NaOH consumption (dNaOH/dt, - - -, g liter<sup>-1</sup> min<sup>-1</sup>); (b) cultivability ( $\Delta$ , CFU ml<sup>-1</sup>) and relative percentages of viable ( $\bullet$ , %), dead ( $\circ$ , %), damaged ( $\bullet$ , %); (c) specific acidification activity (tpH5.5,  $\blacktriangle$ , min); (d) intracellular pH ( $\blacksquare$ ), pHext ( $\square$ ), and pH gradient (dpH = pH<sub>i</sub> - pH<sub>ext</sub>) ( $\blacksquare$ ). Results are the means of at least three independent measurements. They are expressed as a function of the time  $t_m$  to reach the maximal rate of NaOH consumption as an adjusted time.

the  $t_m$  was higher at pH 5 than at pH 6, it must be remembered that the resulting adjusted time was different in these two conditions.

**Analysis of *L. bulgaricus* CFL1 physiological characteristics during pH 6 controlled fermentations.** Viability, mortality, injury, depolarization, pH gradient, and intracellular pH of *L. bulgaricus* CFL1 during pH 6 cultures were quantified by using multiparametric flow cytometry. The results were associated with plate counts and acidification activity (Fig. 1) and discussed as a function of the time  $t_m$ , necessary to reach the maximal NaOH consumption rate (Fig. 1a).

As shown in Fig. 1b, whereas a maximal CFU concentration of  $3.8 \times 10^8 \pm 6 \times 10^7$  CFU ml<sup>-1</sup> was reached at  $t_m+100$  min, a significant drop in the percentage of viable cells was observed throughout growth, from  $93\% \pm 3\%$  before  $t_m-200$  min down to  $34\% \pm 2\%$  at  $t_m+800$  min. The percentage of damaged cells reached a maximum of  $19\% \pm 6\%$  just before the maximal acidification point and then decreased and stood quite stable at  $8\% \pm 1\%$ . The viability decrease was inversely correlated to the percentage of dead cells, which rose by  $58\% \pm 2\%$  in the same period. Depolarized cells depicted a slight rise of  $16\% \pm 1\%$  during growth, before stabilizing during stationary phase. By comparing these results with dead cells percentages, it was observed that only a small part of dead cells were depolarized.

During the stationary phase, CFU concentration stood quite steady at a high level up to 500 min after the maximal acidifi-

cation point before decreasing down to  $7 \times 10^7$  CFU ml<sup>-1</sup> (Fig. 1b). Total viable cell counts (V, expressed as viable plus damaged cells ml<sup>-1</sup>) determined by flow cytometry were well correlated with plate counts (N, expressed in CFU ml<sup>-1</sup>) with a significant coefficient of determination of 0.977. Nevertheless, they were slightly lower than plate counts with a mean difference of 5% throughout growth. The corresponding linear relationship was determined as follows:

$$\log V = 0.95 (\pm 0.05) \cdot \log N + 0.08 (\pm 0.04); R^2 = 0.977 \quad (3)$$

From Fig. 1c it can be seen that the specific acidification activity of *L. bulgaricus* CFL1 cells, measured by the Cinac system, fluctuated throughout the culture when the pH was controlled at pH 6. During exponential phase, the cellular activity decreased significantly, as tpH5.5 increased from  $215 \pm 10$  min to reach a maximum of  $461 \pm 39$  min at the end of the log phase ( $t_m+100$  min). During the stationary phase, the cells recovered better acidification activity, as tpH5.5 decreased down to  $316 \pm 25$  min at  $t_m+300$  min. It remained high and stable at  $327 \pm 9$  min up to  $t_m+500$  min. Finally, it declined gradually in late stationary phase, from  $t_m+500$  min until  $t_m+800$  min.

The time course of intracellular pH (pH<sub>i</sub>) and pH gradient (dpH) was followed during pH 6 fermentations and the results

are summarized in Fig. 1d. These two parameters were stable at  $\text{pHi} = 6.34 \pm 0.01$  and  $\text{dpH} = 0.34 \pm 0.01$  during the log phase, until the stationary phase was achieved ( $\text{tm}+100$  min). At the beginning of the stationary phase, these two parameters decreased to  $\text{pHi} = 6.05 \pm 0.01$  and  $\text{dpH} = 0.05 \pm 0.01$  at  $\text{tm}+200$  min and then remained stable until the end of fermentation.

**Analysis of *L. bulgaricus* CFL1 physiological characteristics during pH 5 controlled fermentations.** In order to further analyze the effect of the fermentation pH on physiological characteristics of *L. bulgaricus* CFL1, they were quantified throughout pH 5 controlled cultures. The results are shown in Fig. 2 and discussed as a function of the time  $\text{tm}$ , necessary to reach the maximal NaOH consumption rate (Fig. 2a).

Fermentations performed at controlled pH 5 led to a maximal CFU concentration of  $5.1 \times 10^8 \pm 4 \times 10^7$  CFU  $\text{ml}^{-1}$ . This maximal concentration was reached at the end of log-phase, at  $\text{tm}+60$  min (Fig. 2b). The total viable cell counts (V) determined by flow cytometry were 6% lower than plate counts (N, in CFU  $\text{ml}^{-1}$ ) throughout growth, and these two variables were well correlated. The corresponding linear relationship was determined as follows:

$$\log V = 0.94 (\pm 0.10) \cdot \log N + 0.09 (\pm 0.01); R^2 = 0.967 \quad (4)$$

As shown in Fig. 2b, the percentages of viable, dead, and injured cells remained constant all along the growth curve at  $67\% \pm 5\%$ ,  $20\% \pm 3\%$ , and  $12\% \pm 2\%$ , respectively. As cell depolarization was low throughout pH 5 culture, with  $<0.05\%$  of depolarized cells, results were not significant and therefore not depicted on Fig. 2b.

From Fig. 2c it can be seen that specific acidification activity decreased throughout pH 5 fermentation, as  $\text{tpH}5.5$  increased from  $254 \pm 9$  min until  $423 \pm 11$  min. Fluctuations were less important than at pH 6. After a slight increase during log phase until  $\text{tm}-100$  min, the time to reach pH 5.5 remained stable at  $314 \pm 5$  min up to  $\text{tm}+400$  min. Finally, the acidification activity decreased, i.e.,  $\text{tpH}5.5$  increased up to  $423 \pm 11$  min at  $\text{tm}+800$  min.

Figure 2d shows that, in spite of a constant and low external pH 5, the cells maintained a high intracellular pH ( $\text{pHi} > 6$ ) and a high pH gradient ( $\text{dpH} > 1$ ) until the beginning of the stationary phase. These two characteristics increased from the beginning of the measurements up to a maximum of  $\text{pHi} = 6.5 \pm 0.05$  and  $\text{dpH} = 1.5 \pm 0.05$  at the end of the log phase ( $\text{tm}+60$ ). They then decreased down to  $\text{pHi} = 5.4 \pm 0.08$  and  $\text{dpH} = 0.4 \pm 0.08$  at  $\text{tm}+800$ .

**Relationship between dead cells and lactate concentration in the culture medium.** The final lactate level was fermentation pH dependent. It was higher ( $>15$  g  $\text{liter}^{-1}$ ) at the end of the cultures conducted at pH 6 than at pH 5 ( $15$  g  $\text{liter}^{-1}$ ) and strongly associated to cellular mortality that increased from  $1.9 \times 10^6$  to  $8.2 \times 10^7$  cells  $\text{ml}^{-1}$  with lactate concentration.

If we consider that the fraction of dissociated lactate differed according to the pH, the mortality of *L. bulgaricus* CFL1 was related to these data. In Fig. 3, a linear correlation can be observed between the logarithm of dead cell counts (D, in cells  $\text{ml}^{-1}$ ) measured by flow cytometry and the lactate concentra-




FIG. 3. Relationship between the logarithm of dead cell counts measured by flow cytometry (D, cells  $\text{ml}^{-1}$ ) and dissociated lactate concentration in the culture medium (Lac, g  $\text{liter}^{-1}$ ), during fermentations performed at pH 6 (■) and pH 5 (△).

tion (Lac, in g  $\text{liter}^{-1}$ ) in the culture medium. This relationship was corroborated by the following linear relationship:

$$\log D = 0.08 (\pm 0.01) \cdot \text{Lac} + 6.47 (\pm 0.08); R^2 = 0.882 \quad (5)$$

## DISCUSSION

**Comparison of *L. bulgaricus* CFL1 physiological state during controlled pH 6 and pH 5 cultures.** As shown in our results, even though *L. bulgaricus* CFL1 fermentations at pH 5 were longer than at pH 6, as previously observed by Amrane and Prigent (2) with low pH-controlled cultures of *Lactobacillus helveticus*, they did not lead to significant differences in terms of final lactose consumption, galactose and lactic acid production, or NaOH consumption. The final CFU concentration was slightly higher at pH 5 than at pH 6, but the difference was not significant if standard deviations are considered. This result is in agreement with that obtained by Mercade et al. (18), who depicted a similar final biomass production when *L. lactis* subsp. *cremoris* was cultured at pH 6.6 or pH 5. However, the concentrations obtained in the current study were significantly higher than those obtained by Rault et al. (20) when cultures of *L. bulgaricus* CFL1 were performed without pH control ( $1.3 \times 10^8$  CFU  $\text{ml}^{-1}$ ). This result is consistent with previous studies which compared free and controlled pH cultures of *L. lactis* (9) and *L. bulgaricus* (17).

At both fermentation pH, viable cell counts were well correlated with plate counts, but they remained slightly lower (5 to 6%, respectively). Such a difference was previously observed with other fluorescent probes by Jepras et al. (16), who indicated that the natural clumping of the cells may affect the flow cytometry counts. Compared to flow cytometry samples that need two dilution steps, plate count samples undergo more dilution steps before being plated onto agar, thus facilitating the break-up of cell clumps. Such a difference was not observed during cultures of *L. bulgaricus* CFL1 performed without pH control (20). This situation may be explained by considering that pH affects *L. bulgaricus* chain length, which decreases with pH (22). This hypothesis was corroborated by microscopic analyses which displayed longer cell chains at pH 6 or pH 5 compared to cultures without pH control.

Comparison of pH 6 and pH 5 cultures displayed significant

differences in the dynamics of viability, membrane integrity, acidification activity, intracellular pH, and pH gradient. We first demonstrated that cellular acidification activity, viability, and membrane integrity were maintained at high levels throughout pH 5 cultures compared to pH 6 cultures. Consequently, a lower pH control value allowed the cells to maintain a more robust physiological state. This result is consistent with those of Schepers et al. (24), who demonstrated that viability and activity of *L. helveticus* cultures conducted at pH 4.2 were better than pH 5.5 cultures.

A significantly higher dpH was observed throughout pH 5 cultures (dpH > 1) than with pH 6 cultures (dpH < 0.5). This higher pH gradient at low pH allowed the cells to maintain their pHi at values compatible with growth and cellular enzymatic activities. This result is in agreement with those of Mercade et al. (18), who compared the dpH of *L. lactis* subsp. *cremoris* grown at pH levels varying between pH 6.6 and pH 4.4.

**The dissociated lactate level influenced cellular viability, membrane integrity, and depolarization.** Differences in viability between the fermentations performed at pH 6 or pH 5 may be explained by the lactate concentration in the culture medium. Lactic acid is a weak acid ( $pK_a = 3.86$ ), with a pH-dependent dissociated lactate level that is prevalent and higher at pH 6 than at pH 5. In light of equation 1, it diminished with pH at the expense of the nondissociated form of lactic acid. The concentration of the lactate form was well correlated with cellular mortality and membrane damage. A high level of dissociated lactate led to a high level of dead cells and then to a poor physiological state of the population. This result indicates that the dissociated form of lactic acid strongly affected the physiological state of *L. bulgaricus* CFL1 by damaging membrane integrity. It differed partially from the conclusions of Amrane and Prigent (2) and Even et al. (12), who stated that the nondissociated form of lactic acid was the main inhibitory compound for growth of lactic acid bacteria. Nevertheless, it was in agreement with the findings of Gonçalves et al. (13), who depicted such a negative influence of the ionic lactate form on *Lactobacillus rhamnosus* at high pH values, whereas the nondissociated lactic acid was the inhibiting form at low pH. This observation indicates that the lactic acid inhibition mechanism may be described by combining total lactic acid concentration and pH, thus taking into account the membrane integrity damage induced by the ionic form at high pH values. From a mechanistic point of view, this phenomenon can be explained by considering pH homeostasis in the cells. Lactic acid affects pH homeostasis through a mechanism that depends upon both proton concentration, normally referred to the weak acid inhibition mechanism that inhibits cellular reactions, and lactate concentration, which might affect membrane integrity and cellular viability.

Whereas high levels of dead cells were observed at pH 6 (>50%), most of these cells were not depolarized (<20% of depolarized cells). These high levels were linked to the high dissociated lactate concentrations inside ( $pH_i > 6$ ) and outside the cells when the  $pH_{ext}$  is 6. It induced a high level of negative charges inside and outside the cell, thus leading to damaged or dead cells that were not depolarized. Therefore, an enhanced dissociated lactate concentration in the culture medium was associated to an increased cellular mortality but not to cellular depolarization.




FIG. 4. Schematic representation of physiological events during pH 6 or pH 5 *L. bulgaricus* CFL1 cultures. Bold lines correspond to main changes in the cellular state. Distinct cellular phases are numbered.

**The pH gradient influenced the progress of physiological events during *L. bulgaricus* CFL1 controlled pH cultures.** On the basis of viability, acidification activity, intracellular pH, and pH gradient measurements, significant differences were observed during the fermentations which made it possible to identify distinct physiological states. Two dynamic schemes representing the progress of the cellular state during fermentations performed at pH 6 and pH 5 were proposed in Fig. 4. Four major periods at pH 6 and three major periods at pH 5 were outlined throughout the fermentations and are discussed below.

From cultivability and viability measurements, the first period (period 1) corresponded, in both cases, to the exponential phase. When the pH was regulated at pH 6, the intracellular pH and the pH gradient remained stable at, respectively,  $pH_i$  6.3 and dpH 0.3. The same has previously been observed with *L. bulgaricus* that displayed stable  $pH_i$  6.5 and dpH 0.1 during log phase of pH 6.4 cultures (17) and with *L. lactis* which exhibited a constant dpH of 1.1 during growth at pH 6.5 (9). At the same time, the cultivability increased concomitantly to the decrease of viability and acidification activity. When entering the stationary phase, the cells displayed a very low specific acidification activity, i.e., a high  $tpH_{5.5}$  ( $tpH_{5.5} = 461$  min) and a decreasing percentage of viable cells, thus indicating that they were in a bad physiological state. This observation confirms that the metabolic activity of the cells was mainly devoted to their growth during the pH 6 log phase instead of their physiological state.

During pH 5 cultures, this behavior differed as acidification activity decreased slightly during the log phase but was maintained at a higher level than at pH 6. The pHi increased during this first period and was maintained in the range of 6.2 to 6.5, which was associated to compatible pHi values for *L. bulgaricus* growth (17, 20). Even if this pHi range is equivalent to that observed at pH 6, the pH gradient was much more elevated (dpH 1.2 instead of dpH 0.3). This difference that characterizes a higher energetic level of the whole-cell system at pH 5 may be related to the more stable and higher acidification activity, as well as to the stable viability observed at pH 5. This result is in agreement with the findings of Rault et al. (20) obtained during cultures of *L. bulgaricus* CFL1 without pH control.

A second period (period 2) was observed during pH 6 fermentations. It corresponded to the beginning of the stationary phase and to the deceleration of the NaOH consumption rate. It was characterized by the recovery of a high acidification activity associated with a rapid decrease of intracellular pH and of pH gradient, down to pHi 6.05 and dpH 0.05, respectively. Viability and membrane integrity continued to decrease, while the percentages of injured and depolarized cells remained stable at low levels. A negative relationship was thus demonstrated between the cellular activity and the intracellular pH and pH gradient. These results indicate that the decrease of the pH gradient makes it possible to deliver energy, thus helping the cells to recover their cellular activity. Such an annulment of dpH at the end of growth has previously been described during *L. lactis* batch cultures at pH 6.5 (9).

During the following period (i.e., period 2 for pH 5 cultures and period 3 for pH 6 cultures), cells were in the stationary phase, as shown by cultivability measurements. Their acidification activity was quite high and stable. Differences between the two pH conditions concerned firstly the viability, which decreased at pH 6 but was high and constant at pH 5. The maintenance of such high cellular activity and viability during the stationary phase has previously been observed during *L. helveticus* cultures at low pH (24). Second, a lower lactate level in the culture medium and a higher pH gradient value were observed at pH 5. According to Rault et al. (20), such high dpH values were compatible with high cellular cultivability, viability, and acidification activity. By combining these results, it may be proposed that cellular viability and membrane integrity are maintained at high values only if the dpH remains high, as during pH 5 cultures or during cultures without pH control (20). On the contrary, when a low pH gradient was measured, the cells no longer maintained their viability and membrane integrity.

The last period (period 3 for pH 5 cultures and period 4 for pH 6 cultures) was characterized by the decrease of both cultivability and acidification activity. At pH 6, this decrease was related to a reduction of viability that reached less than 50% of viable cells in the total population, whereas at pH 5, viability and membrane integrity remained high and constant. As previously explained, this difference in cellular viability was associated with the lower lactate level and the significantly higher pH gradient at pH 5 (dpH > 0.42) compared to pH 6 (dpH = 0.05). Despite these high pH gradient and viability values observed during pH 5 cultures, the pHi decreased to pHi = 5.42. This value was lower than 5.8, which was considered by Rault et al. (20) as the limit value compatible with growth and acid-

ification activity of *L. bulgaricus* CFL1. This situation may explain the decrease in cultivability and acidification activity during this last period.

**Conclusion.** As a final conclusion, this study makes it possible to establish the progress of the physiological events that occur during controlled pH cultures of *L. bulgaricus* CFL1. It points out strong differences in the time course of mortality (membrane damage), viability (enzymatic activity), cellular depolarization, intracellular pH and acidification activity between pH 6 and pH 5 controlled cultures. Viability and acidification activity were higher and more stable during growth when cultures were carried out at pH 5, instead of pH 6. These divergences can mainly be explained by differences in dissociated lactate concentration, intracellular pH, and pH gradient during growth. These key factors grant a better understanding of the influence of fermentation pH on *L. bulgaricus* CFL1 cellular state in growing conditions.

At an industrial stage, such results may prove useful to starter producers and fermented-product manufacturers, allowing these producers and manufacturers to better control the quality of their starters, before freezing or before using them for food fermentation. Within this approach, producers and manufacturers may select a well defined physiological state for a given application and apply well defined conditions (harvesting time and pH) to achieve this physiological state. From an economical point of view, the cost of the flow cytometry method remained quite high, thus representing a drawback to its development. Conversely, the Cinac system is still highly used, which confer it a great interest.

In the future, such discrimination between different physiological states in growing conditions may be useful to standardize the cells in a well-defined physiological state during their production. We also aim at explaining the observed differences by analyzing the intracellular proteome and the membrane fatty acid composition of the cells.

#### ACKNOWLEDGMENTS

This study was financially supported by ARILAIT Recherches (Paris, France) and the Genetic Resources Board (BRG, Paris, France).

We thank Donald White for English revision.

#### REFERENCES

1. Adamberg, K., S. Kask, T. M. Laht, and T. Paalme. 2003. The effect of temperature and pH on the growth of lactic acid bacteria: a pH-auxostat study. *Int. J. Food Microbiol.* **85**:171–183.
2. Amrane, A., and Y. Prigent. 1999. Differentiation of pH and free lactic acid effects on the various growth and production phases of *Lactobacillus helveticus*. *J. Chem. Technol. Biot.* **74**:33–40.
3. Barreto, M. T. O., E. P. Melo, J. S. Almeida, A. M. R. B. Xavier, and M. J. T. Carrondo. 1991. A kinetic method for calculating the viability of lactic starter cultures. *Appl. Microbiol. Biotechnol.* **34**:648–652.
4. Béal, C., and G. Corrieu. 1991. Influence of pH, temperature, and inoculum composition on mixed cultures of *Streptococcus thermophilus* 404 and *Lactobacillus bulgaricus* 398. *Biotechnol. Bioeng.* **38**:90–98.
5. Béal, C., P. Louvet, and G. Corrieu. 1989. Influence of controlled pH and temperature on the growth and acidification of pure cultures of *Streptococcus thermophilus* 404 and *Lactobacillus bulgaricus* 398. *Appl. Microbiol. Biotechnol.* **32**:148–154.
6. Béal, C., M. Marin, E. Fontaine, F. Fonseca, and J. P. Obert. 2008. Production et conservation des ferments lactiques et probiotiques, p. 661–785. In G. Corrieu and F.-M. Luquet (ed.), *Bactéries lactiques, de la génétique aux ferments*. Tec&Doc Lavoisier, Paris, France.
7. Ben Amor, K., P. Breeuwer, P. Verbaarschot, F. M. Rombouts, A. D. L.

- Akkermans, W. M. De Vos, and T. Abee. 2002. Multiparametric flow cytometry and cell sorting for the assessment of viable, injured, and dead *Bifidobacterium* cells during bile salt stress. *Appl. Environ. Microbiol.* **68**:5209–5216.
8. Bibal, B., G. Goma, Y. Vayssier, and A. Pareilleux. 1988. Influence of pH, lactose and lactic acid on the growth of *Streptococcus cremoris*: a kinetic study. *Appl. Microbiol. Biotechnol.* **28**:340–344.
9. Cachon, R., P. Antérieux, and C. Diviès. 1998. The comparative behavior of *Lactococcus lactis* in free and immobilized culture processes. *J. Biotechnol.* **63**:211–218.
10. Corrieu, G., H. E. Spinnler, Y. Jomier, and D. Picque. 1988. Automated system to follow up and control the acidification activity of lactic acid starters. French patent FR 2 629 612.
11. Daly, C., G. F. Fitzgerald, L. O'Connor, and R. Davis. 1998. Technological and health benefits of dairy starter cultures. *Int. Dairy J.* **8**:195–205.
12. Even, S., N. D. Lindley, P. Loubiere, and M. Coccain-Bousquet. 2002. Dynamic response of catabolic pathways to autoacidification in *Lactococcus lactis*: transcript profiling and stability in relation to metabolic and energetic constraints. *Mol. Microbiol.* **45**:1143–1152.
13. Gonçalves, L. M. D., A. Ramos, J. S. Almeida, A. M. R. B. Xavier, and M. J. T. Carrondo. 1997. Elucidation of the mechanism of lactic acid growth inhibition and production in batch cultures of *Lactobacillus rhamnosus*. *Appl. Microbiol. Biotechnol.* **48**:346–350.
14. Grattepanche, F., S. Miescher-Schwenninger, L. Meile, and C. Lacroix. 2008. Recent developments in cheese cultures with protective and probiotic functionalities. *Dairy Sci. Technol.* **88**:421–444.
15. Hutkins, R. W., and N. L. Nannen. 1993. pH homeostasis in lactic acid bacteria. *J. Dairy Sci.* **76**:2354–2365.
16. Jepras, R. I., J. Carter, S. C. Pearson, F. E. Paul, and M. J. Wilkinson. 1995. Development of a robust flow cytometric assay for determining numbers of viable bacteria. *Appl. Environ. Microbiol.* **61**:2696–2701.
17. Mercade, M., F. Duperray, and P. Loubière. 2004. Energetic analysis of cultures of *Lactobacillus delbrueckii* subsp. *bulgaricus*: identification of the type of control between catabolism and anabolism. *Lait* **84**:39–47.
18. Mercade, M., N. D. Lindley, and P. Loubière. 2000. Metabolism of *Lactococcus lactis* subsp. *cremoris* MG 1363 in acid stress conditions. *Int. J. Food Microbiol.* **55**:161–165.
19. Rault, A., C. Béal, S. Ghorbal, J.-C. Ogier, and M. Bouix. 2007. Multiparametric flow cytometry allows rapid assessment and comparison of lactic acid bacteria viability after freezing and during frozen storage. *Cryobiology* **55**:35–43.
20. Rault, A., M. Bouix, and C. Béal. 2008. Dynamic analysis of *Lactobacillus delbrueckii* subsp. *bulgaricus* CFL1 physiological characteristics during fermentation. *Appl. Microbiol. Biotechnol.* **81**:559–570.
21. Rechinger, K. B., and H. Siegmund. 2002. Rapid assessment of cell viability of *Lactobacillus delbrueckii* subsp. *bulgaricus* by measurement of intracellular pH in individual cells using fluorescence ratio imaging microscopy. *Int. J. Food Microbiol.* **75**:53–60.
22. Rhee, S. K., and M. Y. Pack. 1980. Effect of environmental pH on chain length of *Lactobacillus bulgaricus*. *J. Bacteriol.* **144**:865–868.
23. Savoie, S., C. P. Champagne, S. Chiasson, and P. Audet. 2007. Media and process parameters affecting the growth, strain ratios and specific acidifying activities of a mixed lactic starter containing aroma-producing and probiotic strains. *J. Appl. Microbiol.* **103**:163–174.
24. Schepers, A. W., J. Thibault, and C. Lacroix. 2002. *Lactobacillus helveticus* growth and lactic acid production during pH-controlled batch cultures in whey permeate/yeast extract medium. Part I. Multiple factor kinetic analysis. *Enzyme Microb. Technol.* **30**:176–186.
25. Wang, Y., G. Corrieu, and C. Béal. 2005. Fermentation pH and temperature influence the cryotolerance of *Lactobacillus acidophilus* RD758. *J. Dairy Sci.* **88**:21–29.