

# Homo-D-Lactic Acid Fermentation from Arabinose by Redirection of the Phosphoketolase Pathway to the Pentose Phosphate Pathway in L-Lactate Dehydrogenase Gene-Deficient *Lactobacillus plantarum*<sup>∇</sup>

Kenji Okano,<sup>1</sup> Shogo Yoshida,<sup>2</sup> Tsutomu Tanaka,<sup>3</sup> Chiaki Ogino,<sup>2</sup>  
Hideki Fukuda,<sup>3</sup> and Akihiko Kondo<sup>2\*</sup>

Department of Molecular Science and Material Engineering, Graduate School of Science and Technology,<sup>1</sup> Department of Chemical Science and Engineering, Graduate School of Engineering,<sup>2</sup> and Organization of Advanced Science and Technology,<sup>3</sup> Kobe University, 1-1 Rokkodaicho, Nada-ku, Kobe 657-8501, Japan

Received 9 March 2009/Accepted 29 May 2009

**Optically pure D-lactic acid fermentation from arabinose was achieved by using the *Lactobacillus plantarum* NCIMB 8826 strain whose L-lactate dehydrogenase gene was deficient and whose phosphoketolase gene was substituted with a heterologous transketolase gene. After 27 h of fermentation, 38.6 g/liter of D-lactic acid was produced from 50 g/liter of arabinose.**

Poly(lactic acid) (PLA) can be produced from biomass and stereocomplex PLA, which is composed of both poly-L- and poly-D-lactic acid, has recently attracted attention due to its high thermostability. Therefore, D-lactic acid production, in addition to L-lactic acid production, is of significant importance.

Lignocellulose is inedible and represents one of the most abundant biomass sources in the world, consisting of three major components: cellulose, hemicellulose, and lignin (13). Cellulose consists of glucose that is available for microbial lactic acid fermentation, while hemicellulose consists primarily of pentose sugars such as xylose and arabinose, which are unavailable for most microorganisms (5). Therefore, efficient utilization of pentose sugars should be also developed.

Some lactic acid bacteria, such as *Lactobacillus pentosus* (2), *Lactobacillus brevis* (3), *Lactobacillus plantarum* (4), and *Leuconostoc lactis* (11), are known to ferment either or both arabinose and xylose. In these microorganisms, arabinose is converted to xylulose-5-phosphate (X5P) by arabinose isomerase (AraA), ribulokinase (AraB), and ribulose 5-phosphate 4-epimerase (AraD) (4) (Fig. 1). X5P is further converted to equimolar amounts of lactic acid and acetic acid (14). This low yield of lactic acid by heterolactic acid fermentation is an undesirable feature for industrial mass production of lactic acid as a raw material for production of PLA.

In this study, we successfully demonstrated homo-D-lactic acid fermentation from arabinose by engineering a metabolic pathway in L-lactate dehydrogenase gene-deficient *Lactobacillus plantarum* NCIMB 8826 (*L. plantarum*  $\Delta dhL1$ ). *L. plantarum*  $\Delta dhL1$  strain can produce optically pure D-lactic acid (12) from arabinose. However, this strain produces both D-lactic acid and acetic acid from arabinose. As shown in Fig. 1, the phosphoketolase (PK) pathway produces both lactic acid and

acetic acid, but the pentose phosphate (PP) pathway produces only lactic acid as a final product. Therefore, we promoted redirection of the PK pathway to the PP pathway in *L. plantarum*  $\Delta dhL1$  by substituting an endogenous PK gene (*xpk1*) with the heterologous transketolase gene (*tkt*) from *Lactococcus lactis* IL 1403 (1), thereby shifting heterolactic acid fermentation to homolactic acid fermentation.

The bacterial strains, oligonucleotides, and plasmids used in this study are listed in Table 1. The plasmid for disruption of the PK gene (*xpk1*) was constructed as follows. The 520-bp upstream region from the start codon of *xpk1* and the 1,000-bp downstream region from the stop codon of *xpk1* were amplified by PCR from the genome of *L. plantarum* NCIMB 8826 using oligonucleotide primers *xpk1*-up\_F plus *xpk1*-up\_R and *xpk1*-down\_F plus *xpk1*-down\_R, respectively. The resulting fragments were digested with *Sal*I and ligated. Using the ligated fragment (1,520 bp) as a template, the same fragment was amplified by PCR using oligonucleotide primers *xpk1*-up\_F and *xpk1*-down\_R. The amplified fragment was digested with *Xho*I and *Not*I and was subsequently inserted into the *Xho*I and *Not*I sites of the plasmid pG<sup>+</sup>host9 (8). The resulting plasmid was designated pGh9- $\Delta xpk1$ . The plasmid to substitute *xpk1* with the heterologous transketolase gene from *L. lactis* IL 1403 (*tkt*) was constructed as follows. The open reading frame of *tkt* was amplified by PCR from the genome of *L. lactis* IL 1403 using oligonucleotide primers *tkt*\_F and *tkt*\_R. The amplified fragment was digested with *Bgl*II and *Sal*I and was subsequently inserted into the *Bgl*II and *Sal*I sites of plasmid pGh9- $\Delta xpk1$ . The resulting plasmid was designated pGh9-*xpk1*::*tkt*. Disruption and substitution of the *xpk1* gene of *L. plantarum*  $\Delta dhL1$  using pGh9- $\Delta xpk1$  and pGh9-*xpk1*::*tkt* were carried out by pG<sup>+</sup>host plasmid-based double-crossover homologous integration, as described previously (12). The resulting *xpk1*-disrupted or *xpk1*-substituted strains of *L. plantarum*  $\Delta dhL1$  were designated *L. plantarum*  $\Delta dhL1$ - $\Delta xpk1$  and *L. plantarum*  $\Delta dhL1$ -*xpk1*::*tkt*, respectively. Deletion and substitution of *xpk1* were confirmed by PCR, and the stability of a deleted or substituted *xpk1* gene is maintained after several times of passage cultivation.

\* Corresponding author. Mailing address: Department of Chemical Science and Engineering, Graduate School of Engineering, Kobe University, 1-1 Rokkodaicho, Nada-ku, Kobe 657-8501, Japan. Phone and fax: 81-78-803-6196. E-mail: akondo@kobe-u.ac.jp.

<sup>∇</sup> Published ahead of print on 5 June 2009.




FIG. 1. Predicted pathway for arabinose metabolism in *L. plantarum* mutant strains. The key enzymes phosphoketolase and transketolase are underlined, and the key products lactic acid and acetic acid are double-underlined. P, phosphate; FBP, fructose 1,6-bisphosphate; DHAP, dihydroxyacetone phosphate; GAP, glyceraldehyde 3-phosphate.

Cell growth with arabinose was evaluated using modified MRS medium (10 g/liter of peptone, 5 g/liter of yeast extract, 2 g/liter of  $K_2HPO_4$ , 2 g/liter of ammonium citrate, 0.58 g/liter of  $MgSO_4 \cdot 7H_2O$ , and 0.28 g/liter of  $MnSO_4 \cdot 5H_2O$ ; all reagents are purchased from Nacalai Tesque, Inc., Kyoto, Japan) containing 2.0% (wt/vol) arabinose and lacking sugar at 37°C. Figure 2 shows the optical density at 600 nm ( $OD_{600}$ ) of each

strain after 12 h of cultivation. The *L. plantarum*  $\Delta dhL1$  strain was efficiently grown using arabinose as the sole carbon source ( $OD_{600}$  of 1.48). In the case of control cultivation (without arabinose), growth decreased markedly ( $OD_{600}$  of 0.59). Alternatively, *L. plantarum*  $\Delta dhL1-\Delta xpk1$  showed poor growth with arabinose ( $OD_{600}$  of 0.26), even compared with control cultivation ( $OD_{600}$  of 0.45). This lack of growth was restored by

TABLE 1. Strains, plasmids, and oligonucleotide primers used in this study

Strain, plasmid, or primer	Relevant phenotype and description or sequence (5' to 3') <sup>a</sup>	Source or reference
<b>Strains</b>		
<i>Escherichia coli</i>		
VE7108	Kanamycin resistance marker; containing the wild-type <i>repA</i> plasmid gene (not thermosensitive); host for pG <sup>+</sup> host9 DNA manipulation	9
VE6838	Kanamycin resistance marker; VE7108 carrying pG <sup>+</sup> host9	9
<i>Lactobacillus plantarum</i>		
WT	NCIMB 8826 wild-type strain	NCIMB
$\Delta dhL1$	NCIMB 8826 $\Delta dhL1$ strain	12
$\Delta dhL1-\Delta xpk1$	NCIMB 8826 $\Delta dhL1 \Delta xpk1$ strain	This study
$\Delta dhL1-xpk1::tkl$	NCIMB 8826 $\Delta dhL1$ strain with <i>xpk1</i> gene replaced with <i>tkl</i>	This study
<b>Plasmids</b>		
pG <sup>+</sup> host9	Thermosensitive replicon; erythromycin resistance marker	8
pGh9- $\Delta xpk1$	Vector for deletion of <i>xpk1</i> gene of $\Delta dhL1$ ; erythromycin resistance marker	This study
pGh9- <i>xpk1::tkl</i>	Vector for replacement of <i>xpk1</i> gene of $\Delta dhL1$ with <i>tkl</i> ; erythromycin resistance marker	This study
<b>Oligonucleotide primers</b>		
<i>xpk1</i> -up_F	CCGCTCGAGGCTGCTGGTCATTCGGAATT	
<i>xpk1</i> -up_R	AACGAGCTCGTCGACAGATCTTTCAAAAACACTCCTTAATTTGCTTTTC	
<i>xpk1</i> -down_F	ACGCGTCGACGAGCTCGCAGTTAAAATCTTAATTTGAAAATAGCGT	
<i>xpk1</i> -down_R	GTGGCGGCCGCATGAGCTTGCATGGTCCG	
<i>xpk1</i> -up_seq	AAGCACTGAACTGGTTGCC	
<i>xpk1</i> -down_seq	ATAATTA AAAACCAATTA AAAATATGCTCACAATTC	
<i>tkl</i> _F	GGAAGATCTATGTTTGATACTACTGATCAATTGGCTG	
<i>tkl</i> _R	ACGCGTCGACTTAGTCAAGCTCTTTGTATGCTTTAACTA	

<sup>a</sup> For primers, restriction enzyme sites are underlined.




FIG. 2. Growth assay with arabinose using *L. plantarum* NCIMB 8826  $\Delta ldhL1$ , *L. plantarum*  $\Delta ldhL1 - \Delta xpk1$ , and *L. plantarum*  $\Delta ldhL1 - xpk1::tkl$  strains. +, growth in medium with arabinose; -, growth in medium without arabinose. Each bar represents the mean and standard deviations for three independent experiments.

introducing *tkl* into the *xpk1* locus, and the resulting strain *L. plantarum*  $\Delta ldhL1 - xpk1::tkl$  showed a higher  $OD_{600}$  value, 1.73, than that for control cultivation ( $OD_{600}$  of 0.56). These results suggest that the PK pathway is the sole pathway responsible for arabinose assimilation in *L. plantarum*  $\Delta ldhL1$  and that introduction of transketolase enables assimilation of pentose through the PP pathway. Although *L. plantarum*  $\Delta ldhL1$  has another gene, *xpk2*, that encodes a putative PK (7), our results suggest that *xpk2* is silent or its gene product was inactivated in some way under the present cultivation conditions. And although there are several genes that encode presumed transketolases (*tkl2*, *tkl3*, and *tkl4*) and transaldolases (*tal1*, *tal2*, and *tal3*) in the genome of *L. plantarum* (7), these enzymes do not appear to be expressed in their active form, as *L. plantarum*  $\Delta ldhL1 - \Delta xpk1$  could not utilize arabinose (Fig. 2).

Encouraged by these findings, we carried out lactic acid fermentation from 50 g/liter of arabinose using *L. plantarum*  $\Delta ldhL1$  and *L. plantarum*  $\Delta ldhL1 - xpk1::tkl$  strains in a 2.0-liter bioreactor. Heat-sterilized, liquid-modified MRS medium and arabinose solution were added to the fermentor, 10 M  $H_2SO_4$  was added to the medium in order to adjust the pH to 6.0, and 7 ml of inoculum (adjusted to an  $OD_{600}$  of 10 with sterile distilled water) was added. Prior to this addition, inoculum was grown in MRS medium and subcultured at regular intervals

(12 h) to stabilize the growth rate. Temperature was maintained at 36°C, agitation speed was maintained at 100 rpm, and pH was kept at approximately 6.0 ( $\pm 0.03$ ) by the automatic addition of 10 M  $NH_3$  solution. Arabinose and lactic acid concentrations were measured by high-performance liquid chromatography as described previously (6, 10). The optical purity of lactic acid was measured using a BF-5 biosensor (Oji Scientific Instruments, Hyogo, Japan) as described previously (12). Optical purity of lactic acid was defined as follows: optical purity (%) = (D-lactic acid concentration - L-lactic acid concentration)/(D-lactic acid concentration + L-lactic acid concentration)  $\times$  100.

As shown in Fig. 3a and b, *L. plantarum*  $\Delta ldhL1 - xpk1::tkl$  showed significantly higher growth, sugar consumption, and acid production rates than *L. plantarum*  $\Delta ldhL1$ . The  $OD_{600}$  value for *L. plantarum*  $\Delta ldhL1 - xpk1::tkl$  increased with fermentation time and reached 14.9 after 24 h of fermentation, while that of *L. plantarum*  $\Delta ldhL1$  was significantly lower, even after 36 h of fermentation (Fig. 3a). Differences in sugar consumption and acid production were also observed between the strains; *L. plantarum*  $\Delta ldhL1 - xpk1::tkl$  consumed all sugar within 27 h of fermentation and produced 39.2 g/liter of organic acid (Fig. 3b), while fermentation continued after 36 h with *L. plantarum*  $\Delta ldhL1$ , with 19.6 g/liter of arabinose being consumed and 15.0 g/liter of organic acid being produced (Fig. 3b).

As shown in Fig. 3c and Table 2, *L. plantarum*  $\Delta ldhL1$  produced both lactic acid and acetic acid (8.7 and 6.3 g/liter, respectively), while *L. plantarum*  $\Delta ldhL1 - xpk1::tkl$  produced predominantly lactic acid over acetic acid (38.6 and 0.4 g/liter, respectively). As a result, *L. plantarum*  $\Delta ldhL1 - xpk1::tkl$  achieved a high lactic acid yield of 0.82 (gram per gram of consumed arabinose) compared with that of *L. plantarum*  $\Delta ldhL1$  (0.44) (Table 2). Moreover, the optical purity of lactic acid was very high at 99.9% (Table 2). These results strongly indicate that introduction of the PP pathway by substitution of *xpk1* with *tkl* was useful for homo-D-lactic acid fermentation from arabinose.

Surprisingly, substitution of *xpk1* with *tkl* provoked not only a shift to homolactic acid fermentation but also a significant improvement in cell growth and fermentation rate (Fig. 3) (Table 2). The high maximum volumetric productivity of lactic acid by *L. plantarum*  $\Delta ldhL1 - xpk1::tkl$  from arabinose (3.78 g/liter/h) is comparable to that from glucose (4.54 g/liter/h)




FIG. 3. Fermentation experiments with arabinose using *L. plantarum* NCIMB 8826  $\Delta ldhL1$  (closed symbols) and *L. plantarum*  $\Delta ldhL1 - xpk1::tkl$  (open symbols) strains. (a)  $OD_{600}$  of culture (diamonds). (b) Arabinose (squares) and organic acid (as sum of lactic acid and acetic acid) (circles) concentrations. (c) Lactic acid (gray bars) and acetic acid (white bars) concentrations. Data points and bars represent means and standard deviations for three independent experiments.

TABLE 2. Various parameters in arabinose fermentation<sup>a</sup>

Strain	Lactic acid produced (g/liter)	Acetic acid produced (g/liter)	Arabinose consumed (g/liter)	Yield (g lactic acid produced per g arabinose consumed)	Max lactic acid productivity (g/liter/h)	Optical purity of lactic acid (%)
$\Delta ldhL1^b$	8.7 ± 0.6	6.3 ± 0.5	19.6 ± 0.2	0.44 ± 0.03	0.37 ± 0.03	99.8 ± 0.1
$\Delta ldhL1-xpk1::tkf^c$	38.6 ± 1.8	0.4 ± 0.0	47.4 ± 1.7	0.82 ± 0.05	3.78 ± 0.12	99.9 ± 0.0

<sup>a</sup> Values represent means ± standard deviations for three independent experiments.

<sup>b</sup> Values after 36 h of cultivation.

<sup>c</sup> Values after 27 h of cultivation.

(12). As introduction of multicopy plasmids for overexpression of *xpk1* in *L. plantarum*  $\Delta ldhL1$  did not lead to an increased cell growth or fermentation rate (data not shown), conversion of X5P by Xpk1 does not appear to be a rate-limiting step.

In conclusion, we successfully demonstrated homo-D-lactic acid production from arabinose using *ldhL1*-deficient and *xpk1*-substituted *L. plantarum*. Application of the findings obtained in this study to D- and L-lactic acid production from not only arabinose but also xylose, or even a mix of pentose and hexose sugars, is thus expected for lignocellulose utilization. This is the first report of homo-D-lactic acid fermentation from arabinose.

We are grateful to Emmanuelle Maguin for supplying the *E. coli* VE7108 and VE6838 (VE7108 containing pG<sup>+</sup>host9 plasmid) strains.

This work was supported in part by a grant-in-aid for JSPS Fellows (20000860), Tokyo, and Special Coordination Funds for Promoting Science and Technology, Creation of Innovation Centers for Advanced Interdisciplinary Research Areas (Innovative Bioproduction Kobe), MEXT, Japan.

#### REFERENCES

- Bolotin, A., P. Wincker, S. Mauger, O. Jaillon, K. Malarme, J. Weissenbach, S. D. Ehrlich, and A. Sorokin. 2001. The complete genome sequence of the lactic acid bacterium *Lactococcus lactis* ssp. *lactis* IL 1403. *Genome Res.* **11**:731–753.
- Bustos, G., A. B. Moldes, J. M. Cruz, and J. M. Domínguez. 2005. Influence of the metabolism pathway on lactic acid production from hemicellulosic trimming vine shoots hydrolyzates using *Lactobacillus pentosus*. *Biotechnol. Prog.* **21**:793–798.
- Chaillou, S., Y.-C. Bor, C. A. Batt, P. W. Postma, and P. H. Pouwels. 1998. Molecular cloning and functional expression in *Lactobacillus plantarum* 80 of *xyt*, encoding the D-xylose-H<sup>+</sup> symporter of *Lactobacillus brevis*. *Appl. Environ. Microbiol.* **64**:4720–4728.
- Helanto, M., K. Kiviharju, M. Leisola, and A. Nyssölä. 2007. Metabolic engineering of *Lactobacillus plantarum* for production of L-ribulose. *Appl. Environ. Microbiol.* **73**:7083–7091.
- Hofvendahl, K., and B. Hahn-Hägerdal. 2000. Factors affecting the fermentative lactic acid production from renewable resources. *Enzyme Microb. Technol.* **26**:87–107.
- Katahira, S., A. Mizuike, H. Fukuda, and A. Kondo. 2006. Ethanol fermentation from lignocellulosic hydrolysate by a recombinant xylose- and cellobiosaccharide-assimilating yeast strain. *Appl. Microbiol. Biotechnol.* **72**:1136–1143.
- Kleerebezem, M., J. Boekhorst, R. van Kranenburg, D. Molenaar, O. P. Kuipers, R. Leer, R. Turchini, S. A. Peters, H. M. Sandbrink, M. W. E. J. Fiers, W. Stiekema, R. M. K. Lankhorst, P. A. Bron, S. M. Hoffer, M. N. N. Groot, R. Kerkhoven, M. de Vries, B. Ursing, W. M. de Vos, and R. J. Siezen. 2003. Complete genome sequence of *Lactobacillus plantarum* WCFS1. *Proc. Natl. Acad. Sci. USA* **100**:1990–1995.
- Maguin, E., H. Prévost, S. D. Ehrlich, and A. Gruss. 1996. Efficient insertional mutagenesis in lactococci and other gram-positive bacteria. *J. Bacteriol.* **178**:931–935.
- Mora, D., E. Maguin, M. Masiero, C. Parini, G. Ricci, P. L. Manachini, and D. Daffonchio. 2004. Characterization of urease genes cluster of *Streptococcus thermophilus*. *J. Appl. Microbiol.* **96**:209–219.
- Narita, J., K. Okano, T. Kitao, S. Ishida, T. Sewaki, M.-H. Sung, H. Fukuda, and A. Kondo. 2006. Display of  $\alpha$ -amylase on the surface of *Lactobacillus casei* cells by use of the PgsA anchor protein, and production of lactic acid from starch. *Appl. Environ. Microbiol.* **72**:269–275.
- Ohara, H., M. Owaki, and K. Sonomoto. 2006. Xylooligosaccharide fermentation with *Leuconostoc lactis*. *J. Biosci. Bioeng.* **101**:415–420.
- Okano, K., Q. Zhang, S. Shinkawa, S. Yoshida, T. Tanaka, H. Fukuda, and A. Kondo. 2009. Efficient production of optically pure D-lactic acid from raw corn starch by using genetically modified L-lactate dehydrogenase gene-deficient and  $\alpha$ -amylase-secreting *Lactobacillus plantarum* strain. *Appl. Environ. Microbiol.* **75**:462–467.
- Stöcker, M. 2008. Biofuels and biomass-to-liquid fuels in the biorefinery: catalytic conversion of lignocellulosic biomass using porous materials. *Angewandte Chemie* **47**:9200–9211.
- Tanaka, K., A. Komiyama, K. Sonomoto, A. Ishizaki, S. J. Hall, and P. F. Stanbury. 2002. Two different pathways for D-xylose metabolism and the effect of xylose concentration on the yield coefficient of L-lactate in mixed-acid fermentation by the lactic acid bacterium *Lactococcus lactis* IO-1. *Appl. Microbiol. Biotechnol.* **60**:160–167.