

Modified *Bacillus thuringiensis* Toxins and a Hybrid *B. thuringiensis* Strain Counter Greenhouse-Selected Resistance in *Trichoplusia ni*[∇]

Michelle T. Franklin,^{1*} Christal L. Nieman,¹ Alida F. Janmaat,² Mario Soberón,³ Alejandra Bravo,³ Bruce E. Tabashnik,⁴ and Judith H. Myers¹

Department of Zoology, University of British Columbia, 2370-6270 University Boulevard, Vancouver, British Columbia, Canada V6T 1Z4¹; Department of Biology, University of the Fraser Valley, 33844 King Road, Abbotsford, British Columbia, Canada V2S 7M8²; Instituto de Biotecnología, Universidad Nacional Autónoma de México, Apdo. Postal 510-3, Cuernavaca 62250, Morelos, Mexico³; and Department of Entomology, University of Arizona, Tucson, Arizona 85721⁴

Received 20 March 2009/Accepted 6 July 2009

Resistance of greenhouse-selected strains of the cabbage looper, *Trichoplusia ni*, to *Bacillus thuringiensis* subsp. *kurstaki* was countered by a hybrid strain of *B. thuringiensis* and genetically modified toxins Cry1AbMod and Cry1AcMod, which lack helix α -1. Resistance to Cry1AbMod and Cry1AcMod was >100-fold less than resistance to native toxins Cry1Ab and Cry1Ac.

Insecticidal proteins from *Bacillus thuringiensis* are used widely for pest control, but evolution of resistance by pests can reduce their efficacy (3, 4, 6, 14). Resistance to *B. thuringiensis* toxins has been reported in field populations of four species of Lepidoptera, one species in response to sprays (3, 14) and three species in response to transgenic crops (10, 15, 16). Here, we focus on understanding and countering resistance to sprays of *Bacillus thuringiensis* subsp. *kurstaki* that evolved in commercial greenhouse populations of the cabbage looper, *Trichoplusia ni* (7, 17).

We compared responses to single toxins and formulations of *B. thuringiensis* by two resistant strains (GipBtR and GlenBtR) and two related susceptible strains (GipS and GlenS) of *T. ni*. All four strains were started by the collection of larvae in 2001 from commercial greenhouses near Vancouver in British Columbia, Canada (7). Resistance evolved in the greenhouses in response to repeated sprays of DiPel (7), a formulation of *B. thuringiensis* subsp. *kurstaki* strain HD1 containing Cry1Aa, Cry1Ab, Cry1Ac, and Cry2Aa (9). Previously reported concentrations required to kill 50% of larvae (LC₅₀s) indicated that, relative to a susceptible laboratory strain, initial resistance to DiPel was 113-fold in the Gip population (labeled T2c in reference 7) and 24-fold in the Glen population (labeled P5 in reference 7).

We reared larvae on a wheat germ diet (5) at 26°C on a light-to-dark schedule of 16 h:8 h. GipS and GlenS were reared on diet without *B. thuringiensis* toxins, which allowed resistance to decline (7). To maintain resistance, GipBtR and GlenBtR were reared each generation on a diet treated with 5 or 10 mg of DiPel WP (Abbott Laboratories, Ontario, Canada) per milliliter of diet (7). In bioassays, groups of five third-instar larvae were put in 60-ml plastic cups containing diet, and

mortality was assessed after 3 days by gently probing larvae for movement.

We used diet overlay bioassays to evaluate the toxicity to GipBtR and GipS of the protoxin forms of Cry1Ab, Cry1Ac, Cry1AbMod, and Cry1AcMod produced in *B. thuringiensis* strains (12). Cry1AbMod and Cry1AcMod are genetically engineered variants of Cry1Ab and Cry1Ac, respectively, each lacking 56 amino acids from the amino-terminal region, including helix α -1 (12). An 80- μ l aliquot containing distilled water and toxin was dispensed evenly over the surfaces of 2 ml of diet (a mean surface area of 7.1 cm²) and allowed to dry. Fifty to 200 larvae from each strain were tested at five to eight concentrations of each toxin.

We used diet incorporation bioassays (7) to evaluate the toxicities of DiPel and Agree WG (Certis, Columbia, MD) to GipS, GipBtR, GlenS, and GlenBtR. Agree is a formulation of hybrid strain GC91, which was created from the conjugation-like transfer of a plasmid from *B. thuringiensis* subsp. *kurstaki* strain HD191 into *B. thuringiensis* subsp. *aizawai* strain HD135, and it contains Cry1Ac, Cry1C, and Cry1D (1, 8). DiPel and Agree were diluted in distilled water and mixed into diet (7). Twenty-five to 50 larvae from each strain were tested at six to seven concentrations of DiPel and Agree.

We used probit analysis (13) to estimate the LC₅₀s and their 95% fiducial limits (FL), as well as the slopes of concentration-mortality lines and their standard errors. The mortality of larvae fed treated diet was not adjusted for the mortality of control larvae on untreated diet, because the control mortality was low (mean, 3.6%; range, 0 to 16%). LC₅₀s with nonoverlapping 95% FL are significantly different. Resistance ratios were calculated as the LC₅₀ of a resistant strain (GipBtR or GlenBtR) divided by the LC₅₀ of its susceptible counterpart (GipS or GlenS).

The genetically modified toxins Cry1AbMod and Cry1AcMod were much more effective than the native toxins Cry1Ab and Cry1Ac against larvae of *T. ni* from the resistant GipBtR strain (Table 1). Resistance ratios of GipBtR were 580 for Cry1Ab and 1,400 for Cry1Ac but only 5.5 for Cry1AbMod and 9.3 for Cry1AcMod (Table 1). Against GipBtR, the LC₅₀ was 53-fold

* Corresponding author. Mailing address: Department of Zoology, University of British Columbia, 2370-6270 University Boulevard, Vancouver, British Columbia, Canada V6T 1Z4. Phone: (604) 822-5979. Fax: (604) 822-2416. E-mail: franklin@zoology.ubc.ca.

[∇] Published ahead of print on 10 July 2009.

TABLE 1. Responses of resistant (GipBtR and GlenBtR) and susceptible (GipS and GlenS) strains of *T. ni* to native toxins (Cry1Ab and Cry1Ac), modified toxins (Cry1AbMod and Cry1AcMod), and formulations (DiPel and Agree)

Toxin or formulation	Strain	No. of larvae	LC ₅₀ (95% FL) ^a	Slope ± SE	Resistance ratio ^b
Cry1Ab	GipBtR	400	180 (59–2,900) ^c	0.41 ± 0.09	580
	GipS	376	0.30 (0.21–0.41)	0.56 ± 0.06	
Cry1AbMod	GipBtR	400	3.4 (2.6–4.6)	0.52 ± 0.05	5.5
	GipS	375	0.62 (0.51–0.75)	0.99 ± 0.09	
Cry1Ac	GipBtR	600	54 (35–110) ^d	0.50 ± 0.07	1,400
	GipS	1,450	0.038 (0.031–0.046)	0.44 ± 0.02	
Cry1AcMod	GipBtR	600	5.1 (4.4–5.8)	0.85 ± 0.06	9.3
	GipS	1,145	0.55 (0.47–0.64)	0.60 ± 0.03	
DiPel	GipBtR	125	66 (21–420,000) ^e	0.43 ± 0.17	370
	GipS	125	0.18 (0.08–0.27)	0.73 ± 0.16	
Agree	GipBtR	300	4.9 (3.6–7.7)	0.81 ± 0.12	9.9
	GipS	300	0.49 (0.42–0.57)	1.4 ± 0.14	
DiPel	GlenBtR	150	3.2 (2.7–3.9)	1.9 ± 0.27	26
	GlenS	125	0.13 (0.05–0.17)	1.5 ± 0.44	
Agree	GlenBtR	300	2.0 (1.7–2.4)	1.2 ± 0.12	5.9
	GlenS	295	0.34 (0.29–0.39)	1.4 ± 0.17	

^a Concentration that killed 50% and its 95% FL in mg protoxin per cm² diet for toxins and mg formulation per ml of diet for DiPel and Agree.

^b LC₅₀ of the resistant strain divided by the LC₅₀ of the related susceptible strain for each toxin or formulation.

^c Total of 17% mortality at the highest toxin concentration tested (17 mg protoxin/cm² diet).

^d Total of 35% mortality at the highest toxin concentration tested (23 mg protoxin/cm² diet).

^e Total of 24% mortality at the highest toxin concentration tested (15 mg DiPel/ml diet).

higher for Cry1Ab than for Cry1AbMod and 11-fold higher for Cry1Ac than for Cry1AcMod (Table 1). Against GipS, however, the LC₅₀ was 2-fold higher for Cry1AbMod than for Cry1Ab and 14-fold higher for Cry1AcMod than for Cry1Ac (Table 1).

Agree was more effective than DiPel against the two resistant strains GipBtR and GlenBtR (Table 1). Resistance ratios for DiPel were 370 for GipBtR and 26 for GlenBtR compared to resistance ratios for Agree, which were 9.9 for GipBtR and 5.9 for GlenBtR (Table 1). For the two resistant strains, LC₅₀s were higher for DiPel than for Agree (13-fold higher against GipBtR and 1.6-fold higher against GlenBtR) (Table 1). Conversely, against the two susceptible strains, the LC₅₀s were higher for Agree than for DiPel (2.7-fold higher against GipBtR and 2.6-fold higher against GlenBtR).

The resistant GipBtR strain examined here (Table 1) and the resistant GLEN-Cry1Ac-BCS strain of *T. ni* studied by Wang et al. (17) had >500-fold resistance to Cry1Ab and Cry1Ac. Both GipBtR and GLEN-Cry1Ac-BCS were derived from greenhouse populations of *T. ni* that had been sprayed repeatedly with DiPel (7, 17), which contains Cry1Ab and Cry1Ac but not Cry1C or Cry1D (9). The GLEN-Cry1Ac-BCS strain had cross-resistance of only 2.5-fold to Cry1C and 2.4-fold to Cry1D (17). Agree contains Cry1C and Cry1D (8), which probably boosted its efficacy against GipBtR and GlenBtR (Table 1).

The results here with Cry1AbMod and Cry1AcMod extend those of previous work indicating that modified toxins killed larvae of *Manduca sexta* in which susceptibility to Cry1Ab was decreased via RNA interference and also killed larvae of *Pectinophora gossypiella* that had laboratory-selected, genetically based resistance to Cry1Ab and Cry1Ac (12). The efficacy of Cry1AbMod and Cry1AcMod against greenhouse-selected *T. ni* suggests that the modified toxins may be useful against resistance that evolves in commercial agricultural settings. The results here also increase the number of lepidopteran species

against which the modified toxins were effective to three, with each species representing a different family (Sphingidae, Gelechiidae, and Noctuidae). In the two other species, decreased susceptibility to native Cry1A toxins was mediated by alterations in a cadherin protein that binds Cry1Ac (2, 11, 12), whereas the role of cadherin in *T. ni* resistance has not been demonstrated or excluded.

Similar to patterns observed with *P. gossypiella* (12), modified toxins were more effective than native toxins against resistant *T. ni* larvae, but native toxins were more effective than modified toxins against susceptible *T. ni* larvae (Table 1). This raises the intriguing possibility that combinations of native and modified toxins might be especially effective against populations with a mixture of susceptible and resistant individuals. In any case, the Cry1AMod toxins and hybrid *B. thuringiensis* products applied either jointly or separately may be useful for countering or delaying evolution of resistance in *T. ni*. However, further work is needed to determine how native and modified toxins interact when used in combination and how modified toxins perform in the greenhouse and field.

We thank Jing Niu for her technical assistance in the laboratory. We also thank a vegetable grower in British Columbia for providing DiPel, and we thank Certis USA for providing Agree.

Support was provided by the British Columbia Vegetable Greenhouse Growers' Research Council, an NSERC discovery grant, and a grant from USDA/CSREES/NRI.

REFERENCES

- Burges, D. H., and P. Jarrett. November 1991. Preparation of strains of *Bacillus thuringiensis* having an improved activity against certain lepidopteran pests and novel strain produced thereby. U.S. patent 5,063,055.
- Fabrick, J. A., and B. E. Tabashnik. 2007. Binding of *Bacillus thuringiensis* toxin Cry1Ac to multiple sites of cadherin in pink bollworm. *Insect Biochem. Mol. Biol.* 37:97–106.
- Ferré, J., and J. Van Rie. 2002. Biochemistry and genetics of insect resistance to *Bacillus thuringiensis*. *Annu. Rev. Entomol.* 47:501–533.
- Gassmann, A. J., Y. Carrière, and B. E. Tabashnik. 2009. Fitness costs of insect resistance to *Bacillus thuringiensis*. *Annu. Rev. Entomol.* 54:147–163.

5. **Ignoffo, C. M.** 1963. A successful technique for mass-rearing cabbage loopers on a semisynthetic diet. *Ann. Entomol. Soc. Am.* **56**:178–182.
6. **James, C.** 2008. Global status of commercialized biotech/GM crops 2008, brief no. 39. International Service for the Acquisition of Agri-biotech Applications, Ithaca, NY.
7. **Janmaat, A. F., and J. H. Myers.** 2003. Rapid evolution and the cost of resistance to *Bacillus thuringiensis* in greenhouse populations of cabbage looper, *Trichoplusia ni*. *Proc. R. Soc. Lond. B* **270**:2263–2270.
8. **Liu, K., B. Zheng, H. Hong, C. Jiang, R. Peng, J. Peng, Z. Yu, J. Zheng, and H. Yang.** 2004. Characterization of cultured insect cells selected by *Bacillus thuringiensis* crystal toxin. *In Vitro Cell. Dev. Biol. Anim.* **40**:312–317.
9. **Liu, Y.-B., B. E. Tabashnik, and M. Puszta-Carey.** 1996. Field-evolved resistance to *Bacillus thuringiensis* toxin Cry1C in diamondback moth (Lepidoptera: Plutellidae). *J. Econ. Entomol.* **89**:798–804.
10. **Matten, S. R., G. P. Head, and H. D. Quemada.** 2008. How governmental regulation can help or hinder the integration of *Bt* crops into IPM programs, p. 27–39. *In* J. Romeis, A. M. Shelton, and G. G. Kennedy (ed.), *Integration of insect-resistant genetically modified crops within IPM programs*. Springer, New York, NY.
11. **Morin, S., R. W. Biggs, M. S. Sisterson, L. Shriver, C. Ellers-Kirk, D. Higginson, D. Holley, L. J. Gahan, D. G. Heckel, Y. Carrière, T. J. Dennehy, J. K. Brown, and B. E. Tabashnik.** 2003. Three cadherin alleles associated with resistance to *Bacillus thuringiensis* in pink bollworm. *Proc. Natl. Acad. Sci. USA* **100**:5004–5009.
12. **Soberón, M., L. Pardo-López, L. López, I. Gómez, B. E. Tabashnik, and A. Bravo.** 2007. Engineering modified *Bt* toxins to counter insect resistance. *Science* **318**:1640–1642.
13. **Station, R. E.** 1998. Genstat 5, release 4.1. Lawes Agricultural Trust, Rothamsted, United Kingdom.
14. **Tabashnik, B. E.** 1994. Evolution of resistance to *Bacillus thuringiensis*. *Annu. Rev. Entomol.* **39**:47–79.
15. **Tabashnik, B. E., A. J. Gassmann, D. W. Crowder, and Y. Carrière.** 2008. Insect resistance to *Bt* crops: evidence versus theory. *Nat. Biotechnol.* **26**:199–202.
16. **Van Rensburg, J. B. J.** 2007. First report of field resistance by the stem borer, *Busseola fusca* (Fuller) to *Bt*-transgenic maize. *S. Afr. J. Plant Soil* **24**:147–151.
17. **Wang, P., J. Zhao, A. Rodrigo-Simón, W. Kain, A. F. Janmaat, A. M. Shelton, J. Ferré, and J. Myers.** 2007. Mechanism of resistance to *Bacillus thuringiensis* toxin Cry1Ac in a greenhouse population of the cabbage looper, *Trichoplusia ni*. *Appl. Environ. Microbiol.* **73**:1199–1207.