

ATP-Dependent but Proton Gradient-Independent Polyphosphate-Synthesizing Activity in Extraradical Hyphae of an Arbuscular Mycorrhizal Fungus^{∇†}

Chiharu Tani,¹ Ryo Ohtomo,² Mitsuru Osaki,¹ Yukari Kuga,³ and Tatsuhiro Ezawa^{1*}

Research Faculty of Agriculture, Hokkaido University, Sapporo, Hokkaido 060-8589, Japan¹; NARO-NILGS, Nasushiobara, Tochigi 329-2793, Japan²; and Graduate School of Integrated Arts and Sciences, Hiroshima University, Higashi Hiroshima, Hiroshima 739-8521, Japan³

Received 29 June 2009/Accepted 10 September 2009

Arbuscular mycorrhizal (AM) fungi benefit their host plants by supplying phosphate obtained from the soil. Polyphosphate is thought to act as the key intermediate in this process, but little is currently understood about how polyphosphate is synthesized or translocated within arbuscular mycorrhizas. *Glomus* sp. strain HR1 was grown with marigold in a mesh bag compartment system, and extraradical hyphae were harvested and fractionated by density gradient centrifugation. Using this approach, three distinct layers were obtained: layers 1 and 2 were composed of amorphous and membranous materials, together with mitochondria, lipid bodies, and electron-opaque bodies, and layer 3 was composed mainly of partially broken hyphae and fragmented cell walls. The polyphosphate kinase/luciferase system, a highly sensitive polyphosphate detection method, enabled the detection of polyphosphate-synthesizing activity in layer 2 in the presence of ATP. This activity was inhibited by vanadate but not by bafilomycin A₁ or a protonophore, suggesting that ATP may not energize the reaction through H⁺-ATPase but may act as a direct substrate in the reaction. This report represents the first demonstration that AM fungi possess polyphosphate-synthesizing activity that is localized in the organelle fraction and not in the cytosol or at the plasma membrane.

Arbuscular mycorrhizal (AM) fungi are obligate biotrophs that form symbiotic associations with most land plants (29). These fungi promote the growth of host plants via enhanced uptake of phosphate (P_i) and thus play important roles in the terrestrial phosphorus cycle. In the symbiotic phase, AM fungi take up P_i from soil through an extensive network of extraradical hyphae and rapidly accumulate inorganic polyphosphate (polyP). This accumulation was as rapid as that for a polyP-hyperaccumulating bacterium found in activated sludge (6). PolyP is a linear polymer of three to hundreds of molecules of P_i linked by high-energy phosphoanhydride bonds and has been found across all classes of organisms (19). Although polyP is considered to play a central role in long-distance translocation of P_i in AM fungal associations (4, 10, 30, 31), the translocation mechanism, metabolism, and dynamics in the fungi have not been elucidated due to the difficulty in obtaining sufficient fungal material for analysis.

Many enzymes/genes involved in polyP synthesis/metabolism have been identified and characterized in prokaryotes (19). For instance, exopolyphosphatase hydrolyzes the terminal high-energy bonds of polyP, and polyphosphate glucokinase (PPGK) transfers the terminal P_i residue to glucose. Polyphosphate kinase 1 (PPK1) is responsible both for polyP synthesis, using ATP as a phosphoryl donor, and for the reverse ATP-

generating reaction. This enzyme is bound to the plasma membrane (18) and has been found in a wide range of bacteria (17). Unlike the case for prokaryotes, knowledge of polyP synthesis/metabolism in eukaryotes remains limited. The first eukaryotic PPK genes, DdPPK1 (32) and DdPPK2 (14), were identified from the social slime mold *Dictyostelium discoideum*. The products of these genes, as known for bacterial PPK1s, are responsible both for polyP synthesis and for the ATP-generating reaction and have been suggested to be associated with vacuoles or small vesicles (14, 32). Although several homologues of bacterial PPK1 genes have now been found in the genomes of eukaryotic microorganisms (17), yeast *Candida humicola* is the only organism apart from *D. discoideum* for which PPK-like activity has been confirmed (22). The model organism *Saccharomyces cerevisiae* is known to accumulate polyP, to up to 10% of its dry weight (19). A unique polyP synthetic pathway different from those of PPK1 has been proposed for *S. cerevisiae* based on the observation that vacuolar-type H⁺-ATPase (V-ATPase)-defective mutants could not accumulate polyP (23). In this hypothetical pathway, P_i would be polymerized by an analogous system (enzyme) of mitochondrial F₁-ATPase on the vacuolar membrane, using the proton motive force created by V-ATPase (23). On the other hand, Hothorn et al. (16) demonstrated very recently that vacuolar transporter chaperone 4 (VTC4), a small transmembrane protein associated with the membrane, polymerizes P_i by using the γ-P_i residue of ATP as a phosphoryl donor in *S. cerevisiae*.

More than 2 decades ago, Capaccio and Callow (3) reported the presence of polyP-hydrolyzing, -metabolizing (PPGK), and -synthesizing (PPK-like) activities in the soluble (cytosolic)

* Corresponding author. Mailing address: Research Faculty of Agriculture, Hokkaido University, Sapporo, Hokkaido 060-8589, Japan. Phone and fax: 81-11-857-9732. E-mail: tatsu@res.agr.hokudai.ac.jp.

† Supplemental material for this article may be found at <http://aem.asm.org/>.

∇ Published ahead of print on 18 September 2009.

fractions of the hyphae of the AM fungus *Glomus mosseae*. Recently, polyP-hydrolyzing activity was found in both the cytosolic and insoluble (membrane) fractions and then characterized (8). PPGK activity has also been confirmed in the cytosolic fraction, although the activity was quite low and hexokinase (ATP-hexose phosphotransferase) activity appeared to dominate in the glucose phosphorylation process (9). PPK-like activity, however, could not be detected in the same fraction (10), and this seems likely because all other prokaryotic (reviewed in reference 17) and eukaryotic (14, 16, 22, 32) polyP-synthesizing enzymes, so far, are associated with membranes. These observations suggest that AM fungi possess a polyP-synthesizing enzyme that is probably associated with membranes and that ATP may be essential in the synthesis as a phosphoryl donor or via H^+ -ATPase, as suggested by Ogawa et al. (23). In this study, a cell fractionation technique was applied to demonstrate polyP-synthesizing activity in an AM fungus, and then the role of ATP in the synthesis was investigated.

MATERIALS AND METHODS

Fungal material. *Glomus* sp. strain HR1 (MAFF520076) was isolated from the rhizosphere soil of *Lespedeza* sp. grown in acidic soil and deposited in the NIAS Genebank (http://www.gene.affrc.go.jp/about_en.php). The small-subunit rRNA gene sequence (GenBank accession number AB220171) showed high similarity to those of *Glomus manihotis* and *Glomus clarum*, which belong to *Glomus* group A in the Glomeromycota (28). Dwarf marigold (*Tagetes patula* L. cv. Bonanza Orange; Murakami Seed, Ibaraki, Japan) was inoculated with 1,000 spores of *Glomus* sp. strain HR1 and grown in a mesh bag compartment system in which the root/hyphal (R+H) compartment and the hyphal (H) compartment were separated by a mesh bag (37- μ m nylon mesh; 40-ml volume) in a plastic pot (5.5 cm in diameter; 90-ml volume) (8). The R+H compartment was filled with a 1:2 autoclaved washed pumice (4 to 10 mm in diameter)-river sand mixture, while the H compartment was filled with autoclaved river sand. The plants (one batch consisted of 60 to 70 pots) were grown in growth chambers (16-h photoperiod, 25°C, relative humidity of 60%) and received distilled water for the first week, followed by Peters Professional 25-5-20 (N-P₂O₅-K₂O) liquid fertilizer (Scotts-Sierra Horticultural Products, OH) at 50 mg N liter⁻¹ for the second week and then low-P liquid fertilizer (4 mM NH₄NO₃, 1 mM K₂SO₄, 0.75 mM MgSO₄, 2 mM CaCl₂, 0.5 mM Fe-EDTA, and 50 μ M KH₂PO₄) for the third to sixth week every other day in sufficient amounts until the solutions flowed out from the drain holes. At the end of the sixth week, 1 mM KH₂PO₄ solution was applied to the plants in sufficient amounts 4 to 9 h prior to harvest, and extraradical hyphae were collected from the H compartment for all pots of the batch by wet sieving, combined, cleaned under a dissecting microscope as quickly as possible, and placed on ice.

Cell fractionation. All of the following experiments were done under ice-cooled conditions. Approximately 0.5- to 1.5-g hyphal samples were homogenized immediately after the harvest by mortar and pestle with a fivefold volume (wt/vol) of buffer A (1 mM Na₂ATP and 1.2 M sorbitol in 10 mM HEPES-KOH, pH 7.4) with protease inhibitor cocktail for use with fungal and yeast extracts (Sigma-Aldrich, Tokyo, Japan). The slurry was transferred to a 15-ml plastic tube, the mortar and pestle were washed with the same volume of buffer A, and then the solutions were combined. After centrifugation at 160 \times g for 5 min at 4°C, the upper layer was transferred to a new tube, and the pellet was resuspended in a fivefold volume (wt/vol) of buffer A and centrifuged under the same conditions. The upper layers were combined, layered on a continuous density gradient that was prepared by centrifugation of 50% (vol/vol) Percoll (GE Healthcare, Tokyo, Japan) in buffer A at 20,000 \times g for 5 h at 4°C, and centrifuged at 2,500 \times g for 2 h at 4°C. The resultant fractions, with densities of 1.02 to 1.05, 1.06 to 1.09, and >1.15 g ml⁻¹, were designated layers 1, 2, and 3, respectively (Fig. 1). These fractions were collected separately, mixed with a two-fold volume of buffer A, and centrifuged at 18,000 \times g for 15 min at 4°C. The pellets were washed twice with the same buffer, resuspended in a minimum volume of the buffer, and stored on ice.

For transmission electron microscopic (TEM) observation of the fractions, the pellets of layers 1 to 3 were fixed with 4% (wt/vol) paraformaldehyde-1% (wt/vol) glutaraldehyde in buffer A for 16 h at 4°C. After the fixation, the pellets were




FIG. 1. Schematic diagram of fractionation of extraradical hyphae of *Glomus* sp. strain HR1. The densities of the fractions were as follows: layer 1, 1.02 to 1.05 g ml⁻¹; layer 2, 1.06 to 1.09 g ml⁻¹; and layer 3, >1.15 g ml⁻¹.

centrifuged and rinsed three times with 10 mM HEPES-KOH at pH 7.4 for 10 min, embedded in a drop of 1% (wt/vol) low-melting-temperature agarose (Sigma-Aldrich), and processed by a standard procedure for TEM. Briefly, the materials were dehydrated with an ethanol series, substituted with propylene oxide, and infiltrated in Supper's resin (Nisshin EM, Tokyo, Japan) that was polymerized at 70°C overnight. Ultrathin sections were cut with glass knives or a diamond knife, put on copper grids, stained with tannic acid followed by lead citrate, and observed using a JEM-1200EX TEM (JEOL, Tokyo, Japan).

Quantification of polyP. Each fraction was mixed with a ninefold volume of polyP extraction buffer (8 M urea in 50 mM Tris-HCl, pH 8.0), vortexed for 30 s, and centrifuged at 18,000 \times g for 15 min at 4°C. The supernatant was collected, and urea was eliminated using a Micro Bio-Spin P-6 gel filtration column (Bio-Rad Laboratories, Tokyo, Japan) pretreated with TE buffer (1 mM EDTA in 10 mM Tris-HCl, pH 8.0) according to the manufacturer's instructions. PolyP content was determined by the *Escherichia coli* PPK/luciferase method (2), with some modifications (6). Total protein in the fractions was precipitated with trichloroacetic acid and reextracted with NaOH (7), and the concentrations were determined by the modified Lowry method, using a DC protein assay kit (Bio-Rad Laboratories, Tokyo, Japan) according to the manufacturer's instructions.

Enzyme assay conditions. PolyP-synthesizing activity was assessed based on an increase in polyP content after incubation in the presence of ATP. Fifteen microliters of each fraction was mixed with an equal volume of buffer A on ice, and two 10- μ l aliquots of the mixture were transferred to a new tube. One was mixed with 90 μ l of polyP extraction buffer and left on ice as the time zero control, while the other was incubated for 30 min at 30°C. After incubation, the solutions were mixed with 90 μ l of polyP extraction buffer. PolyP concentrations in the mixtures before (time zero control) and after the incubation were determined by the PPK/luciferase method. To examine the involvement of proton motive force in polyP synthesis, the effects of 500 μ M vanadate (plasma membrane-type H⁺-ATPase [P-ATPase] inhibitor), 100 nM bafilomycin A₁ (V-ATPase inhibitor), and 50 μ M carbonyl cyanide *m*-chlorophenylhydrazone (CCCP [a protonophore]) on the synthesizing activity were assessed. In this assessment, 0.5% (vol/vol) dimethyl sulfoxide (DMSO) was added to all reaction mixtures, as the stock solutions of bafilomycin A₁ (20 μ M) and CCCP (10 mM) were prepared with DMSO. PolyP-synthesizing activity was defined based on an increase in P_i residues of polyP per min per unit of protein under the specified conditions.

P- and V-ATPases and cytochrome *c* oxidase (COX) were employed as marker enzymes for the plasma membrane, tonoplast, and mitochondria, respectively, to characterize the fractions. The fractions used for the assessment of P- and V-ATPase activities were prepared from hyphae to which 1 mM P_i solution was not applied prior to harvest, as fractions prepared from P_i-treated hyphae contained a large amount of polyP that was hydrolyzed during incubation and interfered with the assay. P- and V-ATPase activities were determined as "specific inhibitor-sensitive activities," as follows: inhibitor-sensitive activity = total ATPase activity - inhibitor-insensitive activity (total ATPase activity = ATP

TABLE 1. Fractionation of extraradical hyphae of *Glomus* sp. strain HR1

Fraction ^a	PolyP			Protein	
	Total amt (nmol)	Concn (nmol mg ⁻¹ protein)	Recovery (%)	Total amt (mg)	Concn (mg ml ⁻¹)
Homogenate	14,448	504.6	100.0	28.6	3.58
Centrifugation at 160 × g					
Upper layer	4,464	244.8	30.9	15.2	1.69
Pellet	6,382	386.1	44.2	16.5	2.07
Density gradient centrifugation					
Supernatant	1,409	123.4	9.8	11.4	1.27
Layer 1 (1.02 to 1.05 g ml ⁻¹)	75	124.7	0.5	0.6	3.34
Layer 2 (1.06 to 1.09 g ml ⁻¹)	74	45.7	0.5	0.6	2.46
Layer 3 (>1.15 g ml ⁻¹)	42	59.2	0.3	0.7	2.36

^a The detailed procedure for fractionation is described in Materials and Methods. The experiment was conducted several times, using independent batches of fungal material, and one set of results is shown.

hydrolyzing activity – nonspecific phosphatase activity, which was determined using ADP as a substrate).

Twenty-five microliters of each fraction was mixed with an equal volume of reaction mixture consisting of 100 mM KCl, 20 mM MgCl₂, 2.4 M sorbitol, 0.2% DMSO, 2 mM substrate (Na₂ATP or Na₂ADP), and 80 mM HEPES-KOH at pH 7.5 in the presence or absence of inhibitors on ice, and two 20- μ l aliquots of the mixture were transferred to new tubes. One was mixed with an equal volume of 10% (wt/vol) sodium dodecyl sulfate and left on ice as a time zero control, and the other was incubated for 30 min at 30°C, and then an equal volume of 10% sodium dodecyl sulfate was mixed with the solution. The levels of released P_i in the solutions before (time zero control) and after the incubation were determined as previously described (24). The activity was expressed as the amount of P_i released per min per unit of protein under the specified conditions. COX activity was measured as previously described (15). Prior to the assessment, 1 ml of 2 mg ml⁻¹ horse heart cytochrome *c* (Wako Pure Chemicals, Osaka, Japan) in 100 mM potassium phosphate buffer (pH 7.5) was reduced by being mixed with 2 to 3 mg sodium dithionite and passed through a PD-10 Sephadex G-25 M column (GE Healthcare, Tokyo, Japan) equilibrated with the phosphate buffer to remove excess sodium dithionite. Ten microliters of each fraction was mixed with an equal volume of 1% (vol/vol) Triton X-100 in the phosphate buffer and 180 μ l of 1 mg ml⁻¹ reduced cytochrome *c* in the phosphate buffer, and decreases in absorbance at 550 nm were monitored for 3 min at room temperature. The concentration of oxidized cytochrome *c* was calculated based on an extinction coefficient of 18.5 mM⁻¹ cm⁻¹, and the activity was expressed as an increase in oxidized cytochrome *c* per min per unit of protein under the specified conditions. PolyP-hydrolyzing activity was measured based on the liberation of P_i at pH 7.5, using 1 mM polyP type 75+ (average chain length, 79; Sigma-Aldrich) as a substrate (8).

Each enzyme assay was performed in triplicate, using the same batch of material ($n = 3$). The treatments that showed zero or negative values were excluded from subsequent statistical analysis and expressed as “not detected.” Analysis of variance followed by Fisher’s protected least significant difference test or Student’s *t* test was performed by StatView software (SAS Institute Inc., NC) for tests of significance.

RESULTS

Cell fractionation and polyP-synthesizing activity. Cell fractionation was carried out several times using different batches of fungal material (Table 1; see Table S1 in the supplemental material), and representative results are shown in Table 1. After centrifugation of the hyphal homogenate at 160 × g, 20 to 45% of the polyP was recovered in the upper layer. Layers 1, 2, and 3, obtained from the 160 × g upper layer by Percoll density gradient centrifugation, retained only 0.1 to 0.5% of total polyP, whereas 30 to 70% of the polyP in the 160 × g upper layer was recovered from the supernatant of the density gradient centrifugation. PolyP concentrations per unit of protein in layers 1 to 3 were lower than that of the homogenate.

Ultrastructural observations of the fractions revealed that layers 1 and 2 were composed of membranous and amorphous materials, together with organelles that were identified as mitochondria, lipid bodies, and electron-opaque bodies (Fig. 2a to c). A few bacteria were observed in these layers (data not shown). No conspicuous difference in composition between layers 1 and 2 was found. Layer 3 was composed of mainly partially broken hyphae and fragmented cell walls (Fig. 2d and e). Amorphous material associating with the cell wall fragments was also observed.

PolyP-synthesizing activity was consistently detected in layer 2 of all batches in the presence of ATP (Table 2; see Table S2 in the supplemental material). Although layer 1 showed polyP-synthesizing activity in some batches, the detection of the activity in layer 1 was poorly reproducible (data not shown). Layer 2 was thus used for subsequent characterization of the polyP-synthesizing activity. P-ATPase activity was enriched in layer 3, while V-ATPase activity could not be detected in any of the layers after density gradient centrifugation (Table 3). COX activity was enriched in layers 1 and 2. PolyP-hydrolyzing activity was diluted in all layers after density gradient centrifugation.

Substrate for polyP synthesis. To examine whether ATP was used as a direct substrate or as an energy source to create the proton gradients for polyP synthesis (P_i would be the substrate in this case), layer 2 was incubated with ATP, P_i, or both, using the fraction prepared in the presence (original method) or absence (ATP was withheld from buffer A used for the washing step after density gradient centrifugation) of ATP. The activity of the fraction prepared with ATP and incubated with ATP was regarded as a positive control. No polyP-synthesizing activity was detected in the fraction prepared without ATP and incubated with no substrate or with 1 mM P_i (Fig. 3). The fraction prepared without ATP and incubated in the presence of ATP showed polyP-synthesizing activity, although the activity was lower than that of the positive control. No activity was observed in the fraction prepared in the presence of ATP and incubated with both ATP and P_i. The effects of P- and V-ATPase inhibitors and a protonophore were examined for further characterization. Vanadate showed an inhibitory effect on polyP-synthesizing activity, whereas bafilomycin A₁ did not (Fig. 4a). CCCP had no inhibitory effect on the activity (Fig. 4b).




FIG. 2. (a to e) Ultrastructure of layers 2 and 3 of *Glomus* sp. strain HR1 extraradical hyphae. (a to c) Layer 2, composed of amorphous and membranous materials (black arrowheads), with mitochondria (black arrows), lipid body (L), and electron-opaque body (white arrows). (d and e) Layer 3, composed of partially broken hyphae and fragmented hyphal cell wall (asterisks), with which amorphous material was associated (double arrowhead). Bars: 1 μm (a), 0.2 μm (b), 0.5 μm (c), 2 μm (d), and 1 μm (e).

DISCUSSION

Given the fact that AM fungal associations play a key role in phosphorus acquisition of the majority of land plants, it is of significance to clarify the polyP synthetic pathway, the first step of polyP metabolism and translocation, in the biotrophic fungi. PolyP-synthesizing activity in an AM fungus, *Glomus* sp. strain HR1, was successfully demonstrated in combination with cell fractionation. PolyP-synthesizing activity could not be detected

TABLE 2. PolyP-synthesizing activity in fractions prepared from extraradical hyphae of *Glomus* sp. strain HR1

Fraction ^a	Activity (nmol P _i mg ⁻¹ protein min ⁻¹) ^b
Centrifugation at 160 × g (upper layer)	ND
Density gradient centrifugation	
Supernatant	ND
Layer 1	ND
Layer 2	1.01 ± 0.31
Layer 3	ND

^a The detailed procedure for fractionation is described in Materials and Methods. The experiment was conducted several times, using independent batches of fungal material, and one set of results is shown.

^b Rate of increase in polyP during 30 min of incubation at 30°C in the presence of ATP. The amount of polyP is expressed as nanomoles of P_i residues. The values are the mean ± standard error (n = 3). ND, not detected.

TABLE 3. Characterization of layers obtained by cell fractionation of *Glomus* sp. strain HR1 extraradical hyphae, based on the activities of P-ATPase, V-ATPase, COX, and polyphosphatase^a

Fraction ^b	Activity (nmol mg ⁻¹ protein min ⁻¹)			
	P-ATPase ^c	V-ATPase ^c	COX ^d	Polyphosphatase ^e
Centrifugation at 160 × g (upper layer)	5.44	0.95	2.55 ± 0.07	45.6 ± 3.8
Density gradient centrifugation				
Supernatant	ND	ND	ND	30.5 ± 3.0
Layer 1	ND	ND	5.04 ± 0.16	15.2 ± 0.5
Layer 2	ND	ND	6.55 ± 0.04	13.1 ± 0.4
Layer 3	7.68	ND	1.99 ± 0.03	23.0 ± 2.8

^a Each enzyme activity was measured using different batches of fungal material.

^b The detailed procedure for fractionation is described in Materials and Methods. The fractions for the assessment of P- and V-ATPase activities were prepared from hyphae to which 1 mM P_i solution was not applied prior to harvest.

^c Rate of specific inhibitor-sensitive and ATP-specific hydrolysis during 30 min of incubation at 30°C. Vanadate and bafilomycin A₁ were used as specific inhibitors of P- and V-ATPases, respectively. Only the mean values (n = 3) are indicated for P- and V-ATPase activities, because these activities were determined by subtracting the inhibitor-sensitive activity from total ATP-specific activity. ND, not detected.

^d Rate of cytochrome c oxidation during 3 min of incubation at room temperature. The values are means ± standard errors (n = 3).

^e Rate of polyphosphate hydrolysis during 30 min of incubation at 30°C. The values are means ± standard errors (n = 3).




FIG. 3. Effect of ATP and phosphate (P_i) on polyP-synthesizing activity in layer 2 prepared from the extraradical hyphae of *Glomus* sp. strain HR1 in symbiosis with *T. patula*. The fraction was washed with (ATP) or without (Non) 1 mM ATP and incubated in the absence (NS [no substrate]) or presence of 1 mM ATP, P_i , or both. Error bars indicate standard errors ($n = 3$). nd, not detected. Different letters indicate a significant difference ($P < 0.01$; Student's t test between the two treatments).

in the soluble (cytosolic) fractions but was associated with the insoluble cellular components (membranous and amorphous material and organelles), whose densities were within the range of 1.06 to 1.09 g ml⁻¹. It has been shown that bacterial PPKs bind peripherally to the inner plasma membrane (19). The association of the polyP-synthesizing activity with the plasma membrane, however, could be excluded in the AM fungus, as the activity of the P-ATPase, a plasma membrane marker, was not concurrent with the polyP-synthesizing activity. The association of the activity with mitochondria may also be unlikely, because polyP-synthesizing activity was consistently detected only in layer 2, but the mitochondrial marker enzyme COX was enriched in both layers 1 and 2 after fractionation, to the same extent. The amorphous material was not specific to layer 2 and was observed in all fractions, and thus the association of polyP-synthesizing activity with the material is unlikely. The membranous material observed in layer 2 is likely to be vacuolar membrane due to the fact that AM fungal cells contain numerous vacuoles, although the vacuolar marker enzyme V-ATPase could not be detected in any fraction. Possibly, V-ATPase of the fungus is susceptible to physical disruption/fractionation or the activity is intrinsically very low in the fungus. It seems likely that polyP-synthesizing activity is associated with the vacuolar membrane due to the following two reasons. First, all other eukaryotic polyP-synthesizing enzymes, DdPPK1 (32) and DdPPK2 (14) in the slime mold and VTC4 in yeast (16), are associated with vacuoles or small vesicles. Second, AM fungi accumulate polyP in vacuoles, as observed in the germ tubes of *Gigaspora margarita* (20) and in the extra- and intraradical hyphae in *Glomus* sp. strain HR1 (Y. Kuga and T. Ezawa, unpublished observations). Further characterization is required to localize the activity.

Our results are not consistent with those reported by Caccacio and Callow (3), in which polyP-synthesizing activity was detected in the extract (soluble fraction) of an AM fungus. They incubated radioactive ATP with the extract and identified polyP as a trichloroacetic acid-precipitated radioactive compound. This approach was the most reliable for detection of polyP at that time but may not be specific to polyP. One




FIG. 4. Effects of H⁺-ATPase inhibitors (a) and a protonophore (b) on polyP-synthesizing activity in layer 2 prepared from extraradical hyphae of *Glomus* sp. strain HR1 in symbiosis with *T. patula*. (a) Layer 2 was incubated with 1 mM ATP in the presence or absence of 500 μM vanadate (plasma membrane-type H⁺-ATPase inhibitor) or 100 nM bafilomycin A₁ (vacuolar-type H⁺-ATPase inhibitor), nd, not detected. No significant difference in activity between the control and bafilomycin A₁ treatments was observed ($P > 0.05$; Student's t test). (b) Layer 2 was incubated with 1 mM ATP in the presence or absence of 50 μM CCCP. No significant difference in activity between the control and CCCP treatments was observed ($P > 0.05$; Student's t test). Error bars indicate standard errors ($n = 3$).

possibility, therefore, is that the polyP-synthesizing activity reported previously represented an activity of another phosphotransferase-type enzyme that was localized in the cytosol and used ATP as a phosphoryl donor. It is also likely that the extract was contaminated with the membrane fraction that retained polyP-synthesizing activity. The radioactive compound-based method is highly sensitive, such that contamination by trace amounts of membrane may also result in the detection of activity.

The following three technical breakthroughs were indispensable for the detection of polyP-synthesizing activity in the fungus. (i) A small-scale and highly sensitive polyP assay system, the PPK/luciferase-polyP assay system (2), has been applied to AM fungi recently (6, 25). Although the method is relatively less sensitive to polyP species shorter than 20 P_i residues (26), it was essential for the measurement of picomole levels of polyP in the present study. (ii) We selected a fungal species that produces few spores and a large amount of extraradical hyphae. *Glomus* sp. strain HR1 produced fewer spores and a greater hyphal mass over a 6- to 8-week culture than did other species examined (data not shown), although *Glomus* sp. strain HR1 produced a large amount of spores after a 4-month culture. This is a quite important characteristic for cell fractionation, because AM fungal spores are, in general, filled with lipids that aggregate with organelles and interfere with cell fractionation. (iii) We enriched the polyP-synthesizing activity by cell fractionation. The activities of exopolyphosphatase (3, 8) and acid (8, 12) and alkaline (1, 7, 11, 13) phosphatases have been detected in AM fungi, and these enzymes are likely to be involved in polyP hydrolysis. In fact, polyP-hydrolyzing activity in the hyphal homogenate was quite high in the present study. Therefore, cell fractionation that could enrich polyP-synthesizing activity and dilute polyP-hydrolyzing activity was necessary for detection. It should be noted, in addition, that the

polyP-synthesizing activity estimated in our study might have been underestimated due to the concurrence of polyP-hydrolyzing activity in the fraction.

ATP was essential for polyP synthesis in the fungus, and the results strongly suggest that ATP did not energize the reaction via P- or V-ATPase but acted as a direct substrate in the reaction due to the following reasons. First, the V-ATPase inhibitor and protonophore did not inhibit the activity. Second, P-ATPase activity was enriched in layer 3, which showed no polyP-synthesizing activity. Although vanadate, the P-ATPase inhibitor, inhibited the polyP-synthesizing activity, the involvement of P-ATPase, i.e., the proton gradient across the plasma membrane, could be validated only if the activity was inhibited by both vanadate and the protonophoric reagent, because vanadate has been known to inhibit many phosphoryl transfer enzymes as a structural and chemical mimic of phosphate (5). The fact that all polyP-synthesizing enzymes found in prokaryotes (19) and eukaryotes (14, 16, 22, 32) use ATP as a direct phosphoryl donor also supports our observations. Catalysis of the reverse ATP-regenerating reaction is a typical feature of PPK-type enzymes in both prokaryotes (19) and eukaryotes (14, 32) but is unlikely in the case of VTC4 found in yeast (16). In the present study, it was difficult to examine whether layer 2 catalyzed the reverse reaction due to the high background of ATP that was essential for the protection/stabilization of the polyP-synthesizing activity during fractionation. The polyP-synthesizing activity was inhibited by the coexistence of ATP and P_i . It is postulated that P_i interfered with the activity through binding to the catalytic center or that P_i might act as an allosteric effector that regulates the balance between ATP consumption and polyP synthesis in the cell. However, given the fact that cytoplasmic P_i homeostasis is strictly maintained at millimolar levels, e.g., 5 to 10 mM in the case of plants (27), it is unlikely that cytoplasmic P_i is directly involved in the regulation of polyP synthesis in AM fungi. The underlying mechanism will be clarified if the enzyme is purified and localized at the subcellular level.

It is important to identify enzymes/genes involved in polyP metabolism for a clear understanding of the role of polyP in P_i translocation in AM fungi. The present study suggests that purification/localization of the enzyme will be possible using polyP-synthesizing activity as a marker. Application of forthcoming genomic information (21) in conjunction with biochemical analysis will be one promising approach.

ACKNOWLEDGMENTS

The PPK-overexpressing *E. coli* strain was a kind gift from A. Kornberg (Stanford University, CA). We thank Derek B. Goto (Hokkaido University) for a critical reading of the manuscript.

This work was supported by a grant-in-aid for scientific research (17380043) from the Japan Society for the Promotion of Science (T.E. and Y.K.).

REFERENCES

- Aono, T., I. E. Maldonado-Mendoza, G. R. Dewbre, M. J. Harrison, and M. Saito. 2004. Expression of alkaline phosphatase genes in arbuscular mycorrhizas. *New Phytol.* **162**:525–534.
- Ault-Riché, D., C. D. Fraley, C. M. Tzeng, and A. Kornberg. 1998. Novel assay reveals multiple pathways regulating stress-induced accumulations of inorganic polyphosphate in *Escherichia coli*. *J. Bacteriol.* **180**:1841–1847.
- Capaccio, L. C. M., and J. A. Callow. 1982. The enzymes of polyphosphate metabolism in vesicular-arbuscular mycorrhizas. *New Phytol.* **91**:81–91.
- Cox, G., K. J. Moran, F. Sanders, C. Nockolds, and P. B. Tinker. 1980. Translocation and transfer of nutrients in vesicular-arbuscular mycorrhizas. III. Polyphosphate granules and phosphorus translocation. *New Phytol.* **84**:649–659.
- Davies, D. R., and W. G. J. Hol. 2004. The power of vanadate in crystallographic investigations of phosphoryl transfer enzymes. *FEBS Lett.* **577**:315–321.
- Ezawa, T., T. R. Cavagnaro, S. E. Smith, F. A. Smith, and R. Ohtomo. 2004. Rapid accumulation of polyphosphate in extraradical hyphae of an arbuscular mycorrhizal fungus as revealed by histochemistry and a polyphosphate kinase/luciferase system. *New Phytol.* **161**:387–392.
- Ezawa, T., S. Kuwahara, K. Sakamoto, T. Yoshida, and M. Saito. 1999. Specific inhibitor and substrate specificity of alkaline phosphatase expressed in the symbiotic phase of the arbuscular mycorrhizal fungus, *Glomus etunicatum*. *Mycologia* **91**:636–641.
- Ezawa, T., S. E. Smith, and F. A. Smith. 2001. Differentiation of polyphosphate metabolism between the extra- and intraradical hyphae of arbuscular mycorrhizal fungi. *New Phytol.* **149**:555–563.
- Ezawa, T., S. E. Smith, and F. A. Smith. 2001. Enzyme activity involved in glucose phosphorylation in two arbuscular mycorrhizal fungi: indication that polyP is not the main phosphagen. *Soil Biol. Biochem.* **33**:1279–1281.
- Ezawa, T., S. E. Smith, and F. A. Smith. 2002. P metabolism and transport in AM fungi. *Plant Soil* **244**:221–230.
- Funamoto, R., K. Saito, H. Oyaizu, M. Saito, and T. Aono. 2007. Simultaneous in situ detection of alkaline phosphatase activity and polyphosphate in arbuscules within arbuscular mycorrhizal roots. *Funct. Plant Biol.* **34**:803–810.
- Gianinazzi, S., V. Gianinazzi-Pearson, and J. Dexheimer. 1979. Enzymatic studies on the metabolism of vesicular-arbuscular mycorrhiza. III. Ultrastructural localization of acid and alkaline phosphatase in onion roots infected by *Glomus mosseae* (Nicol. & Gerd.). *New Phytol.* **82**:127–132.
- Gianinazzi-Pearson, V., and S. Gianinazzi. 1978. Enzymatic studies on the metabolism of vesicular-arbuscular mycorrhiza. II. Soluble alkaline phosphatase specific to mycorrhizal infection in onion roots. *Physiol. Plant Pathol.* **12**:45–53.
- Gomez-Garcia, M. R., and A. Kornberg. 2004. Formation of an actin-like filament concurrent with the enzymatic synthesis of inorganic polyphosphate. *Proc. Natl. Acad. Sci. USA* **101**:15876–15880.
- Horie, S., and M. Morrison. 1963. Cytochrome c oxidase components. I. Purification and properties. *J. Biol. Chem.* **238**:1855–1860.
- Hothorn, M., H. Neumann, E. D. Lenherr, M. Wehner, V. Rybin, P. O. Hassa, A. Uttenweiler, M. Reinhardt, A. Schmidt, J. Seiler, A. G. Ladurner, C. Herrmann, K. Scheffzek, and A. Mayer. 2009. Catalytic core of a membrane-associated eukaryotic polyphosphate polymerase. *Science* **324**:513–516.
- Kornberg, A. 2008. Abundant microbial inorganic polyphosphate, poly P kinase are underappreciated. *Microbe* **3**:119–123.
- Kornberg, A. 1995. Inorganic polyphosphate: toward making a forgotten polymer unforgettable. *J. Bacteriol.* **177**:491–496.
- Kornberg, A., N. N. Rao, and D. Ault-Riche. 1999. Inorganic polyphosphate: a molecule of many functions. *Annu. Rev. Biochem.* **68**:89–125.
- Kuga, Y., K. Saito, K. Nayuki, R. L. Peterson, and M. Saito. 2008. Ultrastructure of rapidly frozen and freeze-substituted germ tubes of an arbuscular mycorrhizal fungus and localization of polyphosphate. *New Phytol.* **178**:189–200.
- Martin, F., V. Gianinazzi-Pearson, M. Hijri, P. Lammers, N. Requena, I. R. Sanders, Y. Shachar-Hill, H. Shapiro, G. A. Tuskan, and J. P. W. Young. 2008. The long hard road to a completed *Glomus intraradices* genome. *New Phytol.* **180**:747–750.
- McGrath, J. W., A. N. Kulakova, L. A. Kulakov, and J. P. Quinn. 2005. In vitro detection and characterisation of a polyphosphate synthesising activity in the yeast *Candida humicola* G-1. *Res. Microbiol.* **156**:485–491.
- Ogawa, N., J. DeRisi, and P. O. Brown. 2000. New components of a system for phosphate accumulation and polyphosphate metabolism in *Saccharomyces cerevisiae* revealed by genomic expression analysis. *Mol. Biol. Cell* **11**:4309–4321.
- Ohnishi, T., R. S. Gall, and M. L. Mayer. 1975. An improved assay of inorganic phosphate in the presence of extralabile phosphate compounds: application to the ATPase assay in the presence of phosphocreatine. *Anal. Biochem.* **69**:261–267.
- Ohtomo, R., and M. Saito. 2005. Polyphosphate dynamics in mycorrhizal roots during colonization of an arbuscular mycorrhizal fungus. *New Phytol.* **167**:571–578.
- Ohtomo, R., Y. Sekiguchi, T. Mimura, M. Saito, and T. Ezawa. 2004. Quantification of polyphosphate: different sensitivities to short-chain polyphosphate using enzymatic and colorimetric methods as revealed by ion chromatography. *Anal. Biochem.* **328**:139–146.
- Schachtman, D. P., R. J. Reid, and S. M. Ayling. 1998. Phosphorus uptake by plants: from soil to cell. *Plant Physiol.* **116**:447–453.
- Schwarzott, D., C. Walker, and A. Schüssler. 2001. *Glomus*, the largest genus of the arbuscular mycorrhizal fungi (Glomales), is nonmonophyletic. *Mol. Phylogenet. Evol.* **21**:190–197.

29. **Smith, S. E., and D. J. Read.** 2008. Mycorrhizal symbiosis, 3rd ed. Academic Press, San Diego, CA.
30. **Solaiman, M., T. Ezawa, T. Kojima, and M. Saito.** 1999. Polyphosphates in intraradical and extraradical hyphae of an arbuscular mycorrhizal fungus, *Gigaspora margarita*. *Appl. Environ. Microbiol.* **65**:5604–5606.
31. **Viereck, N., P. E. Hansen, and I. Jakobsen.** 2004. Phosphate pool dynamics in the arbuscular mycorrhizal fungus *Glomus intraradices* studied by *in vivo* ³¹P NMR spectroscopy. *New Phytol.* **162**:783–794.
32. **Zhang, H., M. R. Gomez-Garcia, X. Shi, N. N. Rao, and A. Kornberg.** 2007. Polyphosphate kinase 1, a conserved bacterial enzyme, in a eukaryote, *Dicystelium discoideum*, with a role in cytokinesis. *Proc. Natl. Acad. Sci. USA* **104**:16486–16491.