

Chromosome Stability and Gene Loss in Cockroach Endosymbionts^{∇†}

Zakee L. Sabree,^{1,2} Patrick H. Degnan,² and Nancy A. Moran^{2*}

Center for Insect Science¹ and Department of Ecology and Evolutionary Biology,² University of Arizona, Tucson, Arizona

Received 3 February 2010/Accepted 18 April 2010

Insect endosymbiont genomes reflect massive gene loss. Two *Blattabacterium* genomes display colinearity and similar gene contents, despite high orthologous gene divergence, reflecting over 140 million years of independent evolution in separate cockroach lineages. We speculate that distant homologs may replace the functions of some eliminated genes through broadened substrate specificity.

Obligate symbionts of insects exhibit extreme patterns of genome evolution and include the smallest known bacterial genomes (10, 11, 14). Two recently published sequences of *Blattabacterium*, the obligate symbiont of cockroaches (7, 16), present the opportunity to analyze genome evolution in an additional symbiont lineage with extreme genome reduction.

General features. Genomes of the *Blattabacterium* species (*Bacteroidetes*) of *Periplaneta americana* (the BPLAN bacte-

rium) and *Blattella germanica* (the BGE bacterium) are similarly sized at 637 kb and are nearly identical in numbers of protein-encoding sequences (BPLAN, 582; BGE, 586) and RNAs (BPLAN, 37; BGE, 40). Despite differential gene loss, >96% of the genes in each genome are shared (BPLAN, 609/619; BGE, 607/627). Neither the 3.4-kb plasmid harbored by BPLAN nor the genes encoded on the plasmid were reported for the BGE sequenced strain (7). However, we de-




FIG. 1. Gene order comparison of shared *Blattabacterium* orthologs. BGE gene coordinates are plotted on the BPLAN genome. Locus names for orthologs present in one genome and absent in the other are indicated in open boxes, and the arrows point to their approximate locations. Locus names for *Blattabacterium* sp. BGE have been shortened from “BLBBGE” to “BGE.” Arrows that cross the slope indicate the transition from an intact open reading frame (ORF) to a pseudogenized ORF. Annotated pseudogenes are shown as red boxes.

* Corresponding author. Mailing address: Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ 85721. Phone: (520) 626-8355. Fax: (520) 621-9190. E-mail: nmoran@email.arizona.edu.

† Supplemental material for this article may be found at <http://aem.asm.org/>.

∇ Published ahead of print on 23 April 2010.




FIG. 2. Core gene phylogeny of the *Bacteroidetes*. Maximum likelihood phylogeny of the *Bacteroidetes* was estimated using a concatenation of 38 conserved proteins identified in 41 complete and in-progress genomes (see Table S1 in the supplemental material). Consistent with previous phylogenies, *Blattabacterium* (in bold) is sister to “*Ca. Sulcia muelleri*,” endosymbiont of cicadas, leafhoppers, and spittlebugs, and falls just inside the *Flavobacteriales*. Both insect endosymbionts exhibit long branches, indicative of genome-wide elevation in the rate of fixation of slightly deleterious mutations (including nonsynonymous mutations), which is common among obligate endosymbionts. Genome sizes in megabase pairs are indicated in parentheses. Size estimates of unfinished genomes marked with asterisks are based on total base pairs in finished contigs. Support values from RAxML (above) and PhyML (below) were generated from 100 bootstrap replicates; nodes with support of <70% by both methods are not shown.

tected a plasmid bearing at least one of these genes (*nrdF*) in BGE residing within *B. germanica* specimens obtained from North Carolina and maintained in Arizona, indicating either that the plasmid was overlooked or that the strains differ in its presence (for details, see the supplemental material). Nevertheless, the main chromosomes show considerable conservation of genome colinearity with the exception of a single ~19-kb inversion (Fig. 1). This overall structural stability resembles that observed in three other obligate endosymbiont clades for which multiple genomes are available, namely, *Buch-*

nera aphidicola, “*Candidatus Blochmannia*” sp., and “*Candidatus Sulcia muelleri*” (3, 9, 17, 18).

Blattabacterium forms a distinct clade within the phylum *Bacteroidetes*, which is relatively distant from free-living members. “*Ca. Sulcia muelleri*,” the obligate nutritional endosymbiont of cicadas, spittlebugs, and leafhoppers, is their closest relative (Fig. 2). Orthologous pairs of intact *Blattabacterium* protein coding sequences exhibit elevated rates of nonsynonymous substitutions per nonsynonymous sites (dN) compared to free-living relatives (Table 1). This genome-wide trend has

TABLE 1. Comparison of nonsynonymous divergences among pairs of obligate insect endosymbionts and related free-living bacteria

Genome pair ^a	Phylum	Host	Genome wide		Conserved core		Estimated time since divergence (MY) ^c
			<i>n</i>	Mean dN ± SD	<i>n</i>	Mean dN ± SD	
<i>Blattabacterium</i> (BPLAN-BGE)	<i>Bacteroidetes</i>	Blattaria (cockroaches)	570	0.130 ± 0.073	125	0.113 ± 0.062	150–300
“ <i>Ca. Sulcia</i> ” (DSEM-GWSS)	<i>Bacteroidetes</i>	Auchenorrhyncha (cicadas, leafhoppers, spittlebugs, etc.)	202	0.141 ± 0.080	126 ^b	0.131 ± 0.075	200
<i>Flavobacterium johnsoniae</i> - <i>Flavobacterium psychrophilum</i>	<i>Bacteroidetes</i>	Free living	1,268	0.148 ± 0.075	119 ^{b,c}	0.097 ± 0.096	NA
<i>Bacteroides thetaiotaomicron</i> - <i>Bacteroides fragilis</i>	<i>Bacteroidetes</i>	Free living	1,962	0.115 ± 0.073	123 ^{b,d}	0.047 ± 0.043	NA
“ <i>Ca. Blochmannia</i> ” (BPEN-BFL)	<i>Proteobacteria</i>	Camponotini (ants)	577	0.225 ± 0.106	126 ^b	0.186 ± 0.088	16–20
<i>Buchnera</i> (APS-SG)	<i>Proteobacteria</i>	Aphidoidea (aphids)	458	0.149 ± 0.085	125	0.112 ± 0.066	50–70
<i>Buchnera</i> (APS-BP)	<i>Proteobacteria</i>	Aphidoidea (aphids)	458	0.315 ± 0.173	125	0.238 ± 0.130	150–200
<i>E. coli</i> - <i>Salmonella enterica</i>	<i>Proteobacteria</i>	Free living	2,940	0.067 ± 0.058	125	0.020 ± 0.023	100–150

^a Accession numbers for genomes used in this analysis can be found in the supplemental material.

^b TrpD is not fused with TrpG.

^c ThrABC, LeuABCD, and LysA are absent in *F. psychrophilum*, and TrpC is a split gene in *F. johnsoniae* and *F. psychrophilum*.

^d DapD, SucA, and AceF are absent in *B. thetaiotaomicron* and *B. fragilis*.

^e Dates are based on insect fossils except for that for *E. coli*-*S. enterica*. MY, million years; NA, not available.

affected all loci, although genes encoding proteins with poorly defined functions have slightly higher dN estimates than do others (see Table S1 in the supplemental material). Elevated dN is also observed in “*Ca. Sulcia muelleri*” and the sequenced gammaproteobacterial endosymbionts (Table 1). Obligate nutritional endosymbionts encounter dramatic reductions in effective population sizes and selective regimens, which contribute to both gene loss and the elevated accumulation of nonsynonymous substitutions (2, 3, 9, 13, 18).

These two *Blattabacterium* genomes are estimated to have diverged about 140 million years ago based on the insect fossil record (19) and the observation of codiversification of *Blattabacterium* and their hosts (1, 6, 8). Although the sequence similarity is rather low (~85% nucleotide sequence identity), suggesting an ancient divergence, the conservation of gene content and colinearity is quite high. This pattern is echoed in all of the other obligate symbionts of insects for which multiple genomes are available (3, 9, 12, 15, 17, 18); in each case, despite many millions of years of divergence, almost no chromosome rearrangements or gene acquisitions are observed. The absence of *recA* has been suggested as a mechanism contributing to genome stability in obligate endosymbionts (17); however, *recA* is present in both *Blattabacterium* species.

Given the large set of shared genes in these anciently diverged endosymbionts, much of the genome reduction likely occurred early during the association of primitive cockroaches and the ancestor of *Blattabacterium*. Evidence of ongoing gene loss is present in the form of pseudogenes. Of the six BPLAN pseudogenes (Fig. 1), five are intact in BGE and are involved in a variety of pathways, including translation (*trmP/proL*) and amino acid (*cysND*), cofactor (*hemD*), and lipid (*lolD*) biosynthesis. Five additional intact open reading frames unique to BPLAN were also observed.

Gene loss shapes endosymbiont metabolisms. Both genomes encode biosynthetic pathways capable of producing nearly all of the amino acids, except asparagine and glutamine, from waste nitrogen (e.g., urea and ammonia). Interestingly, BGE possesses most of the genes involved in sulfur assimilation (*cysNDHIJ*) and they are either completely absent (*cysI*) or riddled with deletions (*cysND*) in BPLAN. This loss of function has occurred through gradual gene decay and has not dis-

rupted genome colinearity (Fig. 1). A similar pattern of lineage-specific loss of the *cys* genes was observed in *Buchnera aphidicola* strains in host aphids with different diets (17). *B. aphidicola* of *Acythrosiphon pisum* has intact *cysIHGDN* genes while in *B. aphidicola* of *Schizaphis graminum* these genes show multiple deletions and appear nonfunctional, potentially reflecting relaxed selection on sulfur assimilation genes due to the availability of organic sulfur in the *S. graminum* diet (17). Both *B. germanica* and *P. americana* are generalist detritivores, and it is not clear that host diet has shaped the loss/retention of these genes in *Blattabacterium* strains.

In addition to three conserved hypothetical proteins and a precorrin 2 dehydrogenase (*sirBC*), BGE encodes three tricarboxylic acid (TCA) cycle genes (*gltA*, *acnA*, and *icd*) that are completely missing from the BPLAN genome. However, BPLAN does encode 3-isopropylmalate dehydratase (*leuCD*) and 3-isopropylmalate dehydrogenase (*leuB*), which share functional domains with aconitate hydratase (*acnA*) and isocitrate dehydrogenase (*icd*), respectively. Phylogenetic reconstructions of aconitase family enzymes (e.g., LeuCD and AcnA) from *Bacteria*, *Archaea*, and *Eukarya* indicate that these enzymes are derived from a common ancestor, which underwent subsequent ancient duplications (5). Site-directed mutagenesis at amino acid positions 113, 115, and 116 in Icd of *Escherichia coli* resulted in broader substrate specificity and increased preference for isopropylmalate, the native substrate of LeuB, over isocitrate (4). Given the overlapping functional motifs and relatively few amino acid substitutions that could impact substrate specificity, it is possible that the losses of *acnA* and *icd* were largely neutral because of complementation by *leuCD* and *leuB*, respectively. Additionally, most insect obligate endosymbionts with completely sequenced genomes lack intact *acnA* or *icd* genes but do encode intact *leuCD* and *leuB* (Table 2), highlighting the potential dispensability of *acnA* and *icd* genes in insect endosymbionts and suggesting that endosymbiont enzymes may evolve broader substrate specificity (20). Because most of these symbionts provision their hosts with essential amino acids, retention of *leuBCD* genes is likely favored for their role in leucine biosynthesis. Remaining to be determined is the biological significance for *acnA* and *icd* retention by BGE.

TABLE 2. Loss of *acn* and *icd* and retention of *leuCD* and *leuB* is widespread among nutritional insect endosymbionts^a

Obligate insect endosymbiont	Mb	GenBank accession no.	Presence of gene:			
			<i>acn</i>	<i>leuCD</i>	<i>icd</i>	<i>leuB</i>
<i>Baumannia cicadellincola</i> HC	0.69	NC_007984				
<i>Blattabacterium</i> sp. BGE	0.64	NC_013454	+	+	+	+
<i>Blattabacterium</i> sp. BPLAN	0.64	NC_013418		+		+
<i>Buchnera aphidicola</i> APS	0.64	NC_002528		+		+
<i>Buchnera aphidicola</i> BP	0.62	NC_004545		+		+
<i>Buchnera aphidicola</i> CC	0.42	NC_008513		+		+
<i>Buchnera aphidicola</i> SG	0.64	NC_004061		+		+
" <i>Candidatus</i> <i>Azobacteroides pseudotriconymphae</i> "	1.11	NC_011565		+		+
" <i>Candidatus</i> <i>Blochmannia floridanus</i> "	0.71	NC_005061		+		+
" <i>Candidatus</i> <i>Blochmannia pennsylvanicus</i> "	0.79	NC_007292		+		+
" <i>Candidatus</i> <i>Carsonella ruddii</i> PV"	0.16	NC_008512		+		+
" <i>Candidatus</i> <i>Hodgkinia cicadicola</i> DSEM"	0.14	NC_012960				
" <i>Candidatus</i> <i>Sulcia muelleri</i> " GWSS	0.25	NC_010118		+		+
" <i>Candidatus</i> <i>Sulcia muelleri</i> " SMDSEM	0.28	NC_013123		+		+
<i>Wigglesworthia glossinidia</i>	0.70	NC_004344				

^a Endosymbionts with completely sequenced genomes were queried with orthologous proteins from closely related organisms using the blastp program.

The comparison of the *Blattabacterium* genomes provides an additional case of extreme stability of gene order and ongoing gene loss in reduced bacterial genomes. Further investigation of the substrate specificity of endosymbiont enzymes, such as the aconitases, could give insight into the kind of changes that allow these bacteria to function with so few genes or with particular genes missing from pathways that are mostly intact.

Z.L.S. was supported by the University of Arizona Center for Insect Science through National Institutes of Health training grant 2 K12 GM000708. National Science Foundation Award 0626716, to N.A.M., supported this work. P.H.D. received funding from National Science Foundation Awards 0313737 and 0723472 to N.A.M.

REFERENCES

- Clark, J. W., S. Hossain, C. A. Burnside, and S. Kambhampati. 2001. Coevolution between a cockroach and its bacterial endosymbiont: a biogeographical perspective. *Proc. R. Soc. Lond. B Biol. Sci.* **268**:393–398.
- Clark, M. A., N. A. Moran, and P. Baumann. 1999. Sequence evolution in bacterial endosymbionts having extreme base compositions. *Mol. Biol. Evol.* **16**:1586–1598.
- Degnan, P. H., A. B. Lazarus, and J. J. Wernegreen. 2005. Genome sequence of *Blochmannia pennsylvanicus* indicates parallel evolutionary trends among bacterial mutualists of insects. *Genome Res.* **15**:1023–1033.
- Doyle, S. A., S. Y. Fung, and D. E. Koshland, Jr. 2000. Redesigning the substrate specificity of an enzyme: isocitrate dehydrogenase. *Biochemistry* **39**:14348–14355.
- Irvin, S. D., and J. K. Bhattacharjee. 1998. A unique fungal lysine biosynthesis enzyme shares a common ancestor with tricarboxylic acid cycle and leucine biosynthetic enzymes found in diverse organisms. *J. Mol. Evol.* **46**:401–408.
- Lo, N., C. Bandi, H. Watanabe, C. Nalepa, and T. Beninati. 2003. Evidence for cocladogenesis between diverse dictyopteran lineages and their intracellular endosymbionts. *Mol. Biol. Evol.* **20**:907–913.
- Lopez-Sanchez, M. J., A. Neef, J. Pereto, R. Patino-Navarrete, M. Pignatelli, A. Latorre, and A. Moya. 2009. Evolutionary convergence and nitrogen metabolism in *Blattabacterium* strain Bge, primary endosymbiont of the cockroach *Blattella germanica*. *PLoS Genet.* **5**:e1000721.
- Maekawa, K., Y. C. Park, and N. Lo. 2005. Phylogeny of endosymbiont bacteria harbored by the woodroach *Cryptocercus* spp. (Cryptocercidae: Blattaria): molecular clock evidence for a late Cretaceous–early Tertiary split of Asian and American lineages. *Mol. Phylogenet. Evol.* **36**:728–733.
- McCutcheon, J. P., B. R. McDonald, and N. A. Moran. 2009. Convergent evolution of metabolic roles in bacterial co-symbionts of insects. *Proc. Natl. Acad. Sci. U. S. A.* **106**:15394–15399.
- McCutcheon, J. P., B. R. McDonald, and N. A. Moran. 2009. Origin of an alternative genetic code in the extremely small and GC-rich genome of a bacterial symbiont. *PLoS Genet.* **5**:e1000565.
- McCutcheon, J. P., and N. A. Moran. 2007. Parallel genomic evolution and metabolic interdependence in an ancient symbiosis. *Proc. Natl. Acad. Sci. U. S. A.* **104**:19392–19397.
- Moran, N. A. 2003. Tracing the evolution of gene loss in obligate bacterial symbionts. *Curr. Opin. Microbiol.* **6**:512–518.
- Moran, N. A. 1996. Accelerated evolution and Muller's ratchet in endosymbiotic bacteria. *Proc. Natl. Acad. Sci. U. S. A.* **93**:2873–2878.
- Nakabachi, A., A. Yamashita, H. Toh, H. Ishikawa, H. E. Dunbar, N. A. Moran, and M. Hattori. 2006. The 160-kilobase genome of the bacterial endosymbiont *Carsonella*. *Science* **314**:267.
- Perez-Brocal, V., R. Gil, S. Ramos, A. Lamelas, M. Postigo, J. M. Michelena, F. J. Silva, A. Moya, and A. Latorre. 2006. A small microbial genome: the end of a long symbiotic relationship? *Science* **314**:312–313.
- Sabree, Z. L., S. Kambhampati, and N. A. Moran. 2009. Nitrogen recycling and nutritional provisioning by *Blattabacterium*, the cockroach endosymbiont. *Proc. Natl. Acad. Sci. U. S. A.* **106**:19521–19526.
- Tamas, I., L. Klasson, B. Canback, A. K. Naslund, A. S. Eriksson, J. J. Wernegreen, J. P. Sandstrom, N. A. Moran, and S. G. Andersson. 2002. 50 million years of genomic stasis in endosymbiotic bacteria. *Science* **296**:2376–2379.
- van Ham, R. C., J. Kamerbeek, C. Palacios, C. Rausell, F. Abascal, U. Bastolla, J. M. Fernandez, L. Jimenez, M. Postigo, F. J. Silva, J. Tamames, E. Viguera, A. Latorre, A. Valencia, F. Moran, and A. Moya. 2003. Reductive genome evolution in *Buchnera aphidicola*. *Proc. Natl. Acad. Sci. U. S. A.* **100**:581–586.
- Vrsansky, P., V. N. Vishniakova, and A. P. Rasnitsyn. 2002. Order Blattida Latreille, 1810. The cockroaches, p. 263–270. *In* A. P. Rasnitsyn and D. L. J. Quicke (ed.), *History of insects*. Kluwer Academic Publishers, Dordrecht, Netherlands.
- Zientz, E., T. Dandekar, and R. Gross. 2004. Metabolic interdependence of obligate intracellular bacteria and their insect hosts. *Microbiol. Mol. Biol. Rev.* **68**:745–770.