

Correlation between Environmental Factors and Prevalence of *Vibrio parahaemolyticus* in Oysters Harvested in the Southern Coastal Area of Sao Paulo State, Brazil[∇]

Paulo de Souza Costa Sobrinho,[†] Maria T. Destro, Bernadette D. G. M. Franco,
and Mariza Landgraf*

Department of Food and Experimental Nutrition, Faculty of Pharmaceutical Sciences,
University of Sao Paulo, Sao Paulo, Sao Paulo, Brazil

Received 15 April 2009/Accepted 16 October 2009

The presence of *Vibrio parahaemolyticus* in 123 oyster samples collected from an estuary on the southern coast of Sao Paulo state, Brazil, was investigated. Of the 123 samples, 99.2% were positive with densities ranging from <3 to 10⁵ most probable number (MPN)/g. Densities correlated significantly with water temperature ($r = 0.48$; $P < 0.001$) but not with salinity ($r = -0.09$; $P = 0.34$). The effect of harvest site on counts was not significant ($P > 0.05$). These data provide information for the assessment of exposure of *V. parahaemolyticus* in oysters at harvest.

Infections caused by *Vibrio parahaemolyticus* have been reported in several countries (1, 3–5, 14, 15, 16, 17, 18, 19, 20, 22, 24, 26). Among other pathogenic features, *V. parahaemolyticus* strains produce a thermostable hemolysin, known as thermostable direct hemolysin (TDH), as well as TRH (a TDH-related hemolysin) (25, 29). However, not all strains are pathogenic, as less than 1% of food or environmental strains produce TDH or TRH (2, 7, 9, 10, 11).

The most important vehicle for this microorganism is raw or partially cooked shellfish (8, 13, 25, 29). In this study, the densities of *V. parahaemolyticus* in oysters collected in six oyster bed sites in the estuary of Cananeia (25°S; 48°W) in the southern coastal area of Sao Paulo state, Brazil (Fig. 1) between May 2004 and June 2005 were determined using the most probable number (MPN) technique by the method of De Paola and Kaysner (12). Each sample consisted of 15 oysters, pooled in a plastic bag, and transported in a cold box to the laboratory located in the city of Sao Paulo, Brazil. The temperature during transportation did not exceed 13°C, and the travel time was around 5 h. In the laboratory, the oysters were kept under refrigeration (4 to 8°C) and analyzed within 24 h of collection. Oysters were cleaned and shucked by the method of Cook et al. (6). Identification of *V. parahaemolyticus* was based on traditional and API 20E strip biochemical tests (bioMérieux, France), using *V. parahaemolyticus* ATCC 17802 as the reference strain. The observed prevalence of this bacterium was high, as the microorganism was detected in 99.2% (122/123) of the samples and the densities varied between 0.78 and 5.04 log MPN/g.

Strategies for the control of *V. parahaemolyticus* in oysters

depend on understanding the seasonal and geographical distribution and the effects of environmental parameters on the growth of this pathogen. To verify the influence of salinity and temperature of seawater on the density of *V. parahaemolyticus*, samples of water ($n = 123$) were collected from the same depth of oyster beds using 250-ml plastic flasks. Salinity was determined using a salinometer (model RS10; Rosemount Analytical, Cedar Grove, NJ), and the temperature was determined at the time of collection using a digital thermometer (Hanna Instruments). The results are shown in Fig. 2.

Total *V. parahaemolyticus* densities did not correlate significantly with water salinity, as determined by Pearson coefficient ($r = -0.09$; $P = 0.34$). However, the mean salinity varied significantly according to the sampling site and season ($P < 0.05$) (Table 1). The highest mean salinity (24.2 ppt) was detected at site 5 and was 1.4 times higher than at site 2 (17.3 ppt), the lowest mean salinity detected in this study.

The weak correlation between water salinity and *V. parahaemolyticus* densities in oysters suggests that salinity *per se* is a secondary factor for growth of this bacterium, as are turbidity and chlorophyll content in water (27, 30). These results agree with those obtained by Deepanjali et al. (9) and Martinez-Urtiga et al. (21), who did not find correlation between these two parameters. However, they are in contrast with the results reported by DePaola et al. (11), who observed correlation ($P < 0.05$) between salinity and total density of *V. parahaemolyticus*.

The results of this study corroborate existing evidence (10, 11, 27, 30) indicating that the temperature of seawater has a significant correlation ($r = 0.48$; $P < 0.001$) on the densities of *V. parahaemolyticus* in oysters, but they are at odds with results reported by Deepanjali et al. (9), who observed no statistically significant correlation with seawater temperature. The temperature variations observed in the present study (15°C) were lower than those observed by DePaola et al. (22°C) (11) but higher than those reported by Deepanjali et al. (10°C) (9).

The relationship between *V. parahaemolyticus* density and water temperature and salinity were analyzed by multiple linear regression. Results showed that salinity was not significant

* Corresponding author. Mailing address: Departamento de Alimentos e Nutrição Experimental, Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, Av. Prof. Lineu Prestes, 580, B.14, 05508-900 Sao Paulo, SP, Brazil. Phone: 55(11)30912199. Fax: 55(11)38154410. E-mail: landgraf@usp.br.

[†] Present address: Department of Nutrition, Federal University of the Valleys of Jequitinhonha and Mucuri, Diamantina, MG, Brazil.

[∇] Published ahead of print on 18 December 2009.




FIG. 1. Locations of oyster bed sites in the Cananéia estuary on the southern coast of São Paulo state, Brazil. (Courtesy of E. E. de Miranda and A. C. Coutinho [Embrapa Monitoramento por Satélite] [http://www.cdbrazil.cnpem.embrapa.br].)

either for linear effects or for squared effects ($P > 0.05$). For temperature, while the parameter of linear effect was significant ($P < 0.05$), the squared effect was not ($P > 0.05$). Considering the goodness of fit of the model, the following linear regression described the density in oysters the best (Fig. 3): \log_{10} MPN *V. parahaemolyticus*/g = $-0.944 + (0.175 \times \text{temperature})$. The lack of model fitness test was not significant and was considered adequate to express the relationship between *V. parahaemolyticus* density and seawater temperature, in spite of the low R^2 (0.23).

The effect of temperature was further summarized by rank correlation and the use of a smoothing technique (moving average) in which densities corresponding to temperatures within a range of 1°C were pooled to estimate an arithmetic mean of densities in successive intervals. The moving average was calculated using a length of three values. Although seawater temperature and *V. parahaemolyticus* densities were correlated in the present study, the mean densities reached a plateau at temperatures above 24°C and below 20°C (Fig. 4) where the density was not significantly influenced by temperature, consistent with observations reported also by DePaola et al. (11). Our findings could explain the lack of correlation




FIG. 2. Total densities of *Vibrio parahaemolyticus* in oysters from the southern coast of São Paulo state, Brazil. Each bar or point represents the arithmetic mean of six sites, and each error bar represents the standard deviation.

TABLE 1. Seasonal distribution of the total density of *Vibrio parahaemolyticus* in oysters, water temperature, and salinity in the southern coastal area of São Paulo state, Brazil

Variable	Season	No. of samples	Mean ^a	SD	Range
<i>Vibrio parahaemolyticus</i> density (\log_{10} MPN/g)	Winter	35	2.44 A	1.06	<0.48–4.38
	Spring	29	3.26 B	1.17	1.04–5.04
	Summer	24	3.47 B	0.75	1.54–5.04
	Fall	35	3.48 B	1.01	1.04–5.04
Temp (°C)	Winter	35	20.1 A	1.9	14.4–24.0
	Spring	29	23.6 B	1.8	20.0–26.0
	Summer	24	26.7 C	1.4	24.1–29.2
	Fall	35	23.9 B	2.2	20.6–28.3
Salinity (ppt)	Winter	35	22.3 A	4.5	12.2–29.8
	Spring	29	20.2 AB	4.4	11.2–29.4
	Summer	24	18.2 B	4.3	5.3–25.2
	Fall	35	21.8 A	3.9	8.7–28.2

^a Values with different letters are significantly different ($P < 0.05$).




FIG. 3. Goodness of fit regression model of *V. parahaemolyticus* density in oysters and water temperature.

among those parameters detected in tropical oysters by Deepanjali et al. (9) when the temperature varied from 25 to 35°C.

The influence of the season of the year and site of collection on the mean densities was assessed by analysis of variance and Tukey's test, when necessary. As shown in Table 1, the *V. parahaemolyticus* densities were similar in the samples collected during spring, summer, and autumn but differed significantly ($P < 0.05$) in those collected during winter. Densities among samples collected during summer varied less compared to other seasons. Densities above 10^5 MPN/g were detected in six (4.9%) oyster samples (three samples during spring, two samples during summer, and one sample during autumn) collected when the temperature was higher than 24°C and the salinity was higher than 15 ppt. The effect of harvest site on densities was not significant ($P > 0.05$).

Previous studies performed with oysters collected in the same region in Brazil have shown a low incidence of pathogenic *V. parahaemolyticus* (23, 28). Similar results were observed in the present study, as only one oyster sample (0.8%) and only one isolate among 2,243 isolates tested (0.044%) were




FIG. 4. Relationship between the mean density of *V. parahaemolyticus* in oysters and seawater temperature in Cananea estuary, Sao Paulo state, Brazil.

Kanagawa and *tdh* positive. Besides the Kanagawa reaction (24), all strains have been tested for *tlh*, *tdh*, and *trh* genes using PCR (12). The pathogen-positive sample presented with a low density of *V. parahaemolyticus* (3 MPN/g) and was collected during winter, when the temperature was 21°C. Due to the low incidence of pathogenic strains in the samples, correlation between pathogenicity and water temperature or salinity could not be determined.

This study indicates that the presence of *V. parahaemolyticus* in oysters cultivated in the southern coast of Sao Paulo state, Brazil, is high, but pathogenic strains are seldom detected. These results on the ecology and characteristics of *V. parahaemolyticus* are valuable for future risk assessments related to this pathogen in oysters at harvest.

This study was supported by the Conselho Nacional de Pesquisa Científico e Tecnológico (CNPq) and Fundação de Amparo a Pesquisa do Estado de Sao Paulo (FAPESP). We are grateful to FAPESP for the scholarship to P.D.S.C.S.

We thank Brendan Niemira for comments.

REFERENCES

1. Anonymous. 1999. *Vibrio parahaemolyticus*, Japan 1996–1998. Wkly. Epidemiol. Rep. 74:361–363.
2. Anonymous. 2005. Quantitative risk assessment on the public health impact of pathogenic *Vibrio parahaemolyticus* in raw oysters. U.S. Food and Drug Administration, Washington, DC.
3. Centers for Disease Control and Prevention. 1999. Outbreak of *Vibrio parahaemolyticus* infection associated with eating raw oysters and clams harvested from Long Island Sound—Connecticut, New Jersey, and New York, 1998. MMWR Morb. Mortal. Wkly. Rep. 48:48–51.
4. Centers for Disease Control and Prevention. 1998. Outbreak of *Vibrio parahaemolyticus* infections associated with eating raw oysters—Pacific Northwest, 1997. MMWR Morb. Mortal. Wkly. Rep. 47:457–462.
5. Centers for Disease Control and Prevention. 2006. *Vibrio parahaemolyticus* infections associated with consumption of raw shellfish—three states, 2006. MMWR Morb. Mortal. Wkly. Rep. 55:854–856.
6. Cook, D. W., W. Burkhardt, A. DePaola, S. A. McCarthy, and K. R. Calci. 2001. Molluscan shellfish: oysters, mussels, and clams, p. 507–514. In F. P. Downes and K. Ito (ed.), Compendium of methods for the microbiological examination of foods, 4th ed. American Public Health Association, Washington, DC.
7. Cook, D. W., J. D. Bowers, and A. DePaola. 2002. Density of total and pathogenic (*tdh*⁺) *Vibrio parahaemolyticus* in Atlantic and Gulf Coast molluscan shellfish at harvest. J. Food Prot. 65:1873–1880.
8. Daniels, N. A., L. MacKinnon, R. Bishop, S. Altekrose, B. Ray, R. M. Hammond, S. Thompson, S. Wilson, N. H. Bean, P. M. Griffin, and L. Slutsker. 2000. *Vibrio parahaemolyticus* infections in the United States, 1973–1998. J. Infect. Dis. 181:1661–1666.
9. Deepanjali, A., H. S. Kumar, I. Karunasagar, and I. Karunasagar. 2005. Seasonal variation in abundance of total and pathogenic *Vibrio parahaemolyticus* bacteria in oysters along the southwest coast of India. Appl. Environ. Microbiol. 71:3575–3580.
10. DePaola, A., L. H. Hopkins, J. T. Peeler, B. Wentz, and R. M. McPhearson. 1990. Incidence of *Vibrio parahaemolyticus* in U.S. coastal waters and oysters. Appl. Environ. Microbiol. 56:2299–2302.
11. DePaola, A., J. L. Nordstrom, J. C. Bowers, J. G. Wells, and D. W. Cook. 2003. Seasonal abundance of total and pathogenic *Vibrio parahaemolyticus* in Alabama oysters. Appl. Environ. Microbiol. 69:1521–1526.
12. DePaola, A., and C. A. Kaysner. 2004. *Vibrio*, chapter 9. In Food and Drug Administration (ed.), Bacteriological analytical manual, 8th ed., revision A, 1998. U.S. Food and Drug Administration, Washington, DC. <http://www.fda.gov/Food/ScienceResearch/LaboratoryMethods/BacteriologicalAnalyticalManualBAM/default.htm>.
13. Drake, S. L., A. DePaola, and L. A. Jaykus. 2007. An overview of *Vibrio vulnificus* and *Vibrio parahaemolyticus*. Compr. Rev. Food Sci. Food Safety 6:120–144.
14. Fuenzalida, L., C. Hernandez, J. Toro, M. L. Rioseco, J. Romero, and R. T. Espejo. 2006. *Vibrio parahaemolyticus* in shellfish and clinical samples during two large epidemics of diarrhoea in southern Chile. Environ. Microbiol. 8:675–683.
15. Gil, A. I., H. Miranda, C. F. Lanata, A. Prada, E. R. Hall, C. M. Barreno, S. Busrin, N. A. Bhuiyan, D. A. Sack, and G. B. Nair. 2007. O3:K6 serotype of *Vibrio parahaemolyticus* identical to the global pandemic clone associated with diarrhea in Peru. Int. J. Infect. Dis. 11:324–328.
16. Gonzalez-Escalona, N., V. Cachicas, C. Acevedo, M. L. Rioseco, J. A. Ver-

- gara, F. Cabello, J. Romero, and R. T. Espejo. 2005. *Vibrio parahaemolyticus* diarrhea, Chile, 1998 and 2004. *Emerg. Infect. Dis.* **11**:129–131.
17. Hara-Kudo, Y., K. Sugiyama, M. Nishibuchi, A. Chowdhury, J. Yatsuyanagi, Y. Ohtomo, A. Saito, H. Nagano, T. Nishina, H. Nakagawa, H. Konuma, M. Miyahara, and S. Kumagai. 2003. Prevalence of pandemic thermostable direct hemolysin-producing *Vibrio parahaemolyticus* O3:K6 in seafood and the coastal environment in Japan. *Appl. Environ. Microbiol.* **69**:3883–3891.
18. Harth, E., L. Matsuda, C. Hernández, M. L. Rioseco, J. Romero, N. González-Escalon, J. Martínez-Urtaza, and R. T. Espejo. 2009. Epidemiology of *Vibrio parahaemolyticus* outbreaks, southern Chile. *Emerg. Infect. Dis.* **15**:163–168.
19. Leal, N. C., S. C. da Silva, V. O. Cavalcanti, Â. C. T. Figueiroa, V. V. F. Nunes, I. S. Miralles, and E. Hofer. 2008. *Vibrio parahaemolyticus* serovar O3:K6 gastroenteritis in northeast Brazil. *J. Appl. Microbiol.* **105**:691–697.
20. Martínez-Urtaza, J., B. Huapava, R. G. Gavilan, V. Blanco-Abad, J. Ansedo-Bermejo, C. Cadarso-Suarez, A. Figueiras, and J. Trinanes. 2008. Emergence of Asiatic vibrio diseases in South America in phase with El Niño. *Epidemiology* **19**:829–837.
21. Martínez-Urtaza, J., A. Lozano-Leon, J. Varela-Pet, J. Trinanes, Y. Pazos, and O. Garcia-Martin. 2008. Environmental determinants of the occurrence and distribution of *Vibrio parahaemolyticus* in the Rias of Galicia, Spain. *Appl. Environ. Microbiol.* **74**:265–274.
22. Martínez-Urtaza, J., L. Simental, D. Velasco, A. DePaola, M. Ishibashi, Y. Nakaguchi, M. Nishibuchi, D. Carrera-Flores, C. Rey-Alvarez, and A. Pousa. 2005. Pandemic *Vibrio parahaemolyticus* O3:K6, Europe. *Emerg. Infect. Dis.* **11**:1319–1320.
23. Matté, G. R., M. H. Matté, I. G. Rivera, and M. T. Martins. 1994. Distribution of potentially pathogenic vibrios in oysters from a tropical region. *J. Food Prot.* **57**:870–873.
24. Olea, A. M., C. Gonzalez, M. Chiu, C. Vallebuona, M. Labraña, and F. Martiniello. 2005. Brote de gastroenteritis por *Vibrio parahaemolyticus* en Chile. *Rev. Chilena Salud Pública* **9**:51–53.
25. Oliver, J. D., and J. B. Kaper. 2001. *Vibrio* species, p. 263–300. In M. P. Doyle, L. R. Beuchat, and T. J. Montville (ed.), *Food microbiology: fundamentals and frontiers*, 2nd ed. ASM Press, Washington, DC.
26. Pan, T. M., T. K. Wang, C. L. Lee, S. W. Chien, and C. B. Horng. 1997. Food-borne disease outbreaks due to bacteria in Taiwan, 1986 to 1995. *J. Clin. Microbiol.* **35**:1260–1262.
27. Phillips, A. M. B., A. DePaola, J. Bowers, S. Ladner, and D. J. Grimes. 2007. An evaluation of the use of remotely sensed parameters for prediction of incidence and risk associated with *Vibrio parahaemolyticus* in Gulf Coast oysters (*Crassostrea virginica*). *J. Food Prot.* **70**:879–884.
28. Ristori, C. A., S. T. Iaria, D. S. Gelli, and I. N. G. Rivera. 2007. Pathogenic bacteria associated with oysters (*Crassostrea brasiliensis*) and estuarine water along the south coast of Brazil. *Int. J. Environ. Health Res.* **17**:259–269.
29. Su, Y.-C., and C. Liu. 2007. *Vibrio parahaemolyticus*: a concern of seafood safety. *Food Microbiol.* **24**:549–558.
30. Zimmerman, A. M., A. DePaola, J. C. Bowers, J. A. Krantz, J. L. Nordstrom, C. N. Johnson, and D. J. Grimes. 2007. Variability of total and pathogenic *Vibrio parahaemolyticus* densities in northern Gulf of Mexico water and oysters. *Appl. Environ. Microbiol.* **73**:7589–7596.