

Molecular Basis of Sulfonamide and Trimethoprim Resistance in Fish-Pathogenic *Aeromonas* Isolates[∇]

Kristina Kadlec,^{1*} Ellen von Czapiewski,^{1,2#} Heike Kaspar,² Jürgen Wallmann,² Geovana Brenner Michael,¹ Ulrike Steinacker,² and Stefan Schwarz¹

Institute of Farm Animal Genetics, Friedrich-Loeffler-Institut (FLI), Neustadt-Mariensee, Germany,¹ and Federal Office of Consumer Protection and Food Safety (BVL), Berlin, Germany²

Received 11 March 2011/Accepted 14 June 2011

Sulfonamide-trimethoprim-resistant *Aeromonas salmonicida* and motile *Aeromonas* spp. from diseased fish of the GERM-Vet study carried the *sulI* gene together with mostly cassette-borne trimethoprim resistance genes, including the novel gene *dfrA28*. The seven *dfrA* and *dfrB* genes identified were located mostly in class 1 integrons which commonly harbored other gene cassettes.

Aeromonads play an important role as pathogens not only in fish but also in humans and other animals (5). In humans and animals, members of the genus *Aeromonas* may be involved in a variety of intestinal and extraintestinal diseases (5, 14). In fish, the nonmotile *Aeromonas salmonicida* is considered an obligate pathogen and causes a typical furunculosis among salmonids characterized by deep abscesses and hemorrhages on the skin and mouth. Among nonsalmonids, *A. salmonicida* causes atypical furunculosis, also known as “goldfish ulcer” (23). Motile aeromonads, including *A. hydrophila*, *A. veronii*, *A. sobria*, *A. bestiarum*, and *A. caviae*, are considered facultative pathogens and mainly affect immunocompromised fish, causing superficial to deep skin lesions which may progress to ulcers, necrosis, or hemorrhagic septicemia (9). Infections caused by motile aeromonads are probably the most common bacterial disease of freshwater fish (12). Antimicrobial resistance genes, including cassette-borne resistance genes in class 1 integrons, have been described as occurring in *A. salmonicida* and in motile aeromonads (4, 8, 16–18).

Fish-pathogenic bacteria were included for the first time in the German national resistance monitoring program GERM-Vet in 2005 to gain insight into their antimicrobial susceptibilities. A total of 186 fish-pathogenic aeromonads, 173 motile aeromonads and 13 *A. salmonicida* isolates, collected from all over Germany between January 2005 and October 2006 from different disease conditions of commercially reared fish and ornamental fish, were tested for their antimicrobial susceptibilities by broth microdilution according to CLSI document M49-A (2). Since only a combination of trimethoprim-sulfonamide is approved for antimicrobial therapy of fish in Germany, a particular focus was set on resistance to trimethoprim as well as trimethoprim-sulfamethoxazole (SXT) (1:19) and the molecular basis of SXT resistance. Sulfonamides alone were not included in the GERM-Vet test panel. Resistance to trimethoprim is often associated with gene cassettes located in

class 1 or class 2 integrons, and the sulfonamide resistance gene *sulI* is part of the 3'-conserved segment of class 1 integrons. Thus, all SXT-resistant isolates were screened by PCR for class 1 and class 2 integrons and associated gene cassettes (6). Same-size amplicons were compared by restriction analysis, and at least one representative of each amplicon type was cloned and sequenced completely. Plasmid pPCR-Script Amp SK(+) (Stratagene) or pCR-Blunt (Invitrogen) served as the cloning vector, and the *Escherichia coli* strain XL-10-Gold Kan or *E. coli* strain One Shot TOP10 served as the recipient. Sequence analysis started with the M13 reverse and forward primers and was completed by primer walking. In addition, the resistance phenotype conferred by the cassette-located genes was confirmed in the *E. coli* clones by susceptibility testing according to CLSI document M31-A3 (3). The species of the SXT-resistant isolates were determined by PCR amplification of an internal part of the *gyrB* gene with subsequent sequence analysis (24).

High SXT MIC values of >32/608 µg/ml were seen for 33 *Aeromonas* species isolates and two *A. salmonicida* isolates, all of which also exhibited trimethoprim MICs of >128 µg/ml. All 35 isolates carried the sulfonamide resistance gene *sulI*, as confirmed by PCR. Of these, 29 isolates (27 *Aeromonas* species and the 2 *A. salmonicida* isolates) harbored a single class 1 integron while 4 *Aeromonas* sp. isolates carried two class 1 integrons. No class 2 integrons were detected. Among the class 1 integrons, gene cassettes with the seven different trimethoprim resistance genes, *dfrA1*, *dfrA12*, *dfrA14*, *dfrA28*, *dfrB1*, *dfrB3*, or *dfrB4*, were identified by sequence analysis. PCR analysis confirmed the presence of the trimethoprim resistance gene *dfrA1* outside a class 1 integron in four SXT-resistant *Aeromonas* sp. isolates. Two of these isolates, however, carried class 1 integrons with *catB3-aadA2* gene cassettes. Twelve different gene cassette arrangements were detected (Fig. 1 and Table 1). The identified genes located in these gene cassettes matched the resistance phenotypes of the *Aeromonas* isolates and of the *E. coli* clones. The *dfrA* or *dfrB* gene cassettes were usually accompanied by gene cassettes conferring other resistance properties or by *orfV* or *orfF* cassettes whose functions are unknown (Table 1). The most frequently detected cassette combination was *dfrA12-*

* Corresponding author. Mailing address: Institute of Farm Animal Genetics, Friedrich-Loeffler-Institut (FLI), Höltystr. 10, 31535 Neustadt-Mariensee, Germany. Phone: 49-5034-871-242. Fax: 49-5034-871-246. E-mail: kristina.kadlec@fli.bund.de.

These authors contributed equally to the study.

∇ Published ahead of print on 15 July 2011.




FIG. 1. Schematic presentation of the 12 gene cassette arrangements detected among the *A. salmonicida* and motile *Aeromonas* species isolates. The reading frames are shown as arrows, with the arrowhead indicating the direction of transcription. Trimethoprim resistance genes are shown in black, and aminoglycoside resistance genes are in gray. A distance scale in kb is given below each map. The 5' conserved and the 3' conserved segments of the class 1 integrons are depicted as "5'CS" and "3'CS" on a gray background.

TABLE 1. Presence of different gene cassettes within class 1 integrons among the 33 SXT-resistant motile aeromonads and the two *A. hydrophila* isolates

Gene cassette combination	Associated resistance phenotype	No. of isolates	Origin(s) (n) ^a	Aeromonad(s) in which combination was detected (n) ^b	EMBL accession no.
<i>dfrA28-orfV</i>	Trimethoprim	1	Koi carp	<i>A. veronii</i> biovar <i>sobria</i>	FM877476
<i>dfrA12-orfF</i>	Trimethoprim	1	Koi carp	<i>A. veronii</i> biovar <i>sobria</i>	FM877477
<i>orf-aadA5</i>	Streptomycin-spectinomycin	1	Koi carp	<i>A. veronii</i> biovar <i>sobria</i>	FM877481
<i>dfrB3-aadA1</i>	Trimethoprim, streptomycin-spectinomycin	1	Koi carp	<i>A. sobria</i>	FM877478
<i>dfrA1-aadA1</i>	Trimethoprim, streptomycin-spectinomycin	4	Koi carp (2), goldfish, salmon	<i>A. sobria</i> , <i>A. veronii</i> biovar <i>sobria</i> , <i>A. veronii</i> biovar <i>veronii</i> , <i>A. salmonicida</i>	FM877479
<i>dfrA14-recombined aadA6</i>	Trimethoprim, streptomycin-spectinomycin	1	Ram cichlid	<i>A. sobria</i>	FM877480
<i>catB3-aadA2</i>	Chloramphenicol, streptomycin-spectinomycin	3	Koi carp (3)	<i>A. hydrophila</i> (2), <i>A. bestiarum</i>	FM877482
<i>dfrA12-orfF-aadA2</i>	Trimethoprim, streptomycin-spectinomycin	19	Koi carp (13), goldfish, rainbow trout (2), nase, ND (2)	<i>A. hydrophila</i> (11), <i>A. bestiarum</i> (2), <i>A. caviae</i> , <i>A. veronii</i> biovar <i>sobria</i> (4), <i>A. veronii</i> biovar <i>veronii</i>	FM877483
<i>dfrB4-catB3-aadA1</i>	Trimethoprim, chloramphenicol, streptomycin-spectinomycin	1	Koi carp	<i>A. veronii</i> biovar <i>veronii</i>	FM877484
<i>dfrB1-aadA1-catB2</i>	Trimethoprim, streptomycin-spectinomycin, chloramphenicol	2	Rainbow trout, ND	<i>A. sobria</i> , <i>A. salmonicida</i>	FM877485
<i>dfrA12-orfF</i> with integrated IS630- <i>aadA2</i>	Trimethoprim, streptomycin-spectinomycin	1	Koi carp	<i>A. veronii</i> biovar <i>veronii</i>	FM877486
<i>arr-3-aacA4-bla_{OXA-10}-aadA1</i>	Rifampin, gentamicin, penicillins, streptomycin-spectinomycin	1	Koi carp	<i>A. hydrophila</i>	FM877487

^a ND, no data available; *n*, no. of isolates; koi carp, *Cyprinus carpio*; goldfish, *Carassius auratus*; salmon, *Salmo salar*; ram cichlid, *Mikrogeophagus ramirezi*; clown loach, *Chromobotia makracanthus*; rainbow trout, *Oncorhynchus mykiss*; nase, *Chondrostoma nasus*.

^b The species was determined by sequencing an internal part of the *gvrB* gene (24). *n*, no. of isolates.

orfF-aadA2, present in 19 isolates, followed by *dfrA1-aadA1*, present in 4 isolates.

The *dfrA28* gene was a novel *dfrA* gene identified for the first time during this study. The *dfrA28* gene cassette had a total size of 562 bp and a 59-base element of 82 bp. The *dfrA28* gene had a GTG start codon and comprised 474 bp. It coded for a dihydrofolate reductase of 157 amino acids. Identities at the nucleotide and amino acid sequence levels to the most-related *dfrA27* gene and the DfrA27 protein from *E. coli* (GenBank accession no. EU675686) were 96% and 97%, respectively. Downstream of the *dfrA28* cassette, a structure was detected which showed, at least in the 5'-terminal region, similarities to a gene cassette. The reading frame *orfV*, 249 bp in size, did not exhibit any similarities to known reading frames and ended in the 3' conserved segment of the class 1 integron. A 59-base element was not detectable. The combination *dfrA12-orfF-aadA2* with the insertion sequence IS630 integrated into *orfF* and the cassette combination *arr-3-aacA4-bla_{OXA-10}-aadA1* also have not been described before.

However, most of the other cassette combinations have been observed in class 1 integrons of bacteria from fish or other animals. Identical or closely related *dfrA12-orfF* and *catB3-aadA2* combinations have been found in *A. hydrophila* from foodborne outbreak-suspect samples and environmental sources in Taiwan (1). The *aadA5* cassette in the *orf-aadA5* combination was indistinguishable from an *aadA5* cassette found in a class 1 integron of *E. coli* from a catfish (11). The *dfrA1-aadA1* combination is widespread among Gram-negative bacteria, and very similar gene cassettes have been detected in

Salmonella enterica serovar *Infantis* from swine (13), *E. coli* from swine (6), *Salmonella enterica* serovar *Typhimurium* from horses (21), and *E. coli* from a wastewater treatment plant (10). The combination of *dfrA14* and recombined *aadA6* cassettes has previously been detected in *E. coli* of porcine origin (6). The combination *dfrA12-orfF-aadA2* has been seen in *Salmonella* *Typhimurium* from imported seafood (7), *Salmonella enterica* serovar *Schwarzengrund* from chicken (22), and *A. hydrophila* (15) or *E. coli* (6) from swine. The combination *dfrB1-aadA1-catB2* was seen as part of larger *bla_{VIM-1}*-carrying integron structures in *Klebsiella pneumoniae* (19) and *E. coli* (20).

This study provided for the first time data on the SXT susceptibility status of fish-pathogenic aeromonads from Germany and the trimethoprim and sulfonamide resistance genes present. The association of SXT resistance with class 1 integrons, which in part carried gene cassettes for other resistance properties, bears the risk of coselection and persistence of other resistance genes under the selective pressure imposed by the use of trimethoprim-sulfonamide combinations. The observation that similar or even identical gene cassettes have been detected in bacteria from fish, humans, food-producing animals, and/or companion animals underlines the presence of a common resistance gene pool.

ACKNOWLEDGMENTS

We thank Anne-Kathrin Hinz und Katharina Stolz for excellent technical assistance.

This study was financially supported by the Bundesverband für Tiergesundheit e.V. (BfT).

REFERENCES

1. Chang, Y. C., D. Y. Shih, J. Y. Wang, and S. S. Yang. 2007. Molecular characterization of class 1 integrons and antimicrobial resistance in *Aeromonas* strains from foodborne outbreak-suspect samples and environmental sources in Taiwan. *Diagn. Microbiol. Infect. Dis.* **59**:191–197.
2. CLSI. 2006. Methods for broth dilution susceptibility testing of bacteria isolated from aquatic animals; approved guideline. CLSI document M49-A. Clinical and Laboratory Standards Institute, Wayne, PA.
3. CLSI. 2008. Performance standards for antimicrobial disk and dilution susceptibility test for bacteria isolated from animals; approved standard, 3rd edition. CLSI document M31-A3. Clinical and Laboratory Standards Institute, Wayne, PA.
4. Jacobs, L., and H. Y. Chenia. 2007. Characterization of integrons and tetracycline resistance determinants in *Aeromonas* spp. isolated from South African aquaculture systems. *Int. J. Food Microbiol.* **114**:295–306.
5. Janda, J. M., and S. L. Abbott. 2010. The genus *Aeromonas*: taxonomy, pathogenicity, and infection. *Clin. Microbiol. Rev.* **23**:35–73.
6. Kadlec, K., and S. Schwarz. 2008. Analysis and distribution of class 1 and class 2 integrons and associated gene cassettes among *Escherichia coli* isolates from swine, horses, cats and dogs collected in the BfT-GermVet monitoring study. *J. Antimicrob. Chemother.* **62**:469–473.
7. Khan, A. A., et al. 2009. Identification and characterization of class 1 integron resistance gene cassettes among *Salmonella* strains isolated from imported seafood. *Appl. Environ. Microbiol.* **75**:1192–1196.
8. L'Abée-Lund, T. M., and H. Sørum. 2001. Class 1 integrons mediate antibiotic resistance in the fish pathogen *Aeromonas salmonicida* worldwide. *Microb. Drug Resist.* **7**:263–272.
9. McGarey, D. J., et al. 1991. The role of motile aeromonads in the fish disease, ulcerative disease syndrome (UDS). *Experientia* **47**:441–444.
10. Moura, A., I. Henriques, R. Ribeiro, and A. Correia. 2007. Prevalence and characterization of integrons from bacteria isolated from a slaughterhouse wastewater treatment plant. *J. Antimicrob. Chemother.* **60**:1243–1250.
11. Nawaz, M., et al. 2009. Molecular characterization of tetracycline-resistant genes and integrons from avirulent strains of *Escherichia coli* isolated from catfish. *Foodborne Pathog. Dis.* **6**:553–559.
12. Noga, E. J. 2000. Fish disease: diagnosis and treatment. Iowa State University Press, Ames, IA.
13. O'Mahony, R., et al. 2005. Antimicrobial resistance in isolates of *Salmonella* spp. from pigs and the characterization of an *S. Infantis* gene cassette. *Foodborne Pathog. Dis.* **2**:274–281.
14. Parker, J. L., and J. G. Shaw. 2011. *Aeromonas* spp. clinical microbiology and disease. *J. Infect.* **62**:109–118.
15. Poole, T. L., T. R. Callaway, K. M. Bischoff, C. E. Warnes, and D. J. Nisbet. 2006. Macrolide inactivation gene cluster *mphA-mrx-mphR* adjacent to a class 1 integron in *Aeromonas hydrophila* isolated from a diarrhoeic pig in Oklahoma. *J. Antimicrob. Chemother.* **57**:31–38.
16. Rosser, S. J., and H. K. Young. 1999. Identification and characterization of class 1 integrons in bacteria from an aquatic environment. *J. Antimicrob. Chemother.* **44**:11–18.
17. Schmidt, A. S., M. S. Bruun, I. Dalsgaard, and J. L. Larsen. 2001. Incidence, distribution, and spread of tetracycline resistance determinants and integron-associated antibiotic resistance genes among motile aeromonads from a fish farming environment. *Appl. Environ. Microbiol.* **67**:5675–5682.
18. Schmidt, A. S., M. S. Bruun, J. L. Larsen, and I. Dalsgaard. 2001. Characterization of class 1 integrons associated with R-plasmids in clinical *Aeromonas salmonicida* isolates from various geographical areas. *J. Antimicrob. Chemother.* **47**:735–743.
19. Tato, M., T. M. Coque, F. Baquero, and R. Cantón. 2010. Dispersal of carbapenemase *bla_{VIM-1}* gene associated with different Tn402 variants, mercury transposons, and conjugative plasmids in Enterobacteriaceae and *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **54**:320–327.
20. Tórtola, M. T., et al. 2005. First detection of a carbapenem-hydrolyzing metalloenzyme in two enterobacteriaceae isolates in Spain. *Antimicrob. Agents Chemother.* **49**:3492–3494.
21. Vo, A. T., E. van Duijkeren, A. C. Fluit, and W. Gaastra. 2007. A novel *Salmonella* genomic island 1 and rare integron types in *Salmonella* Typhimurium isolates from horses in The Netherlands. *J. Antimicrob. Chemother.* **59**:594–599.
22. Vo, A. T., E. van Duijkeren, W. Gaastra, and A. C. Fluit. 2010. Antimicrobial resistance, class 1 integrons, and genomic island 1 in *Salmonella* isolates from Vietnam. *PLoS One* **5**:e9440.
23. Wiklund, T., and I. Dalsgaard. 1998. Occurrence and significance of atypical *Aeromonas salmonicida* in non-salmonid and salmonid fish species: a review. *Dis. Aquat. Organ.* **32**:49–69.
24. Yáñez, M. A., V. Catalán, D. Apráiz, M. J. Figueras, and A. J. Martínez-Murcia. 2003. Phylogenetic analysis of members of the genus *Aeromonas* based on *gyrB* gene sequences. *Int. J. Syst. Evol. Microbiol.* **53**:875–883.