

Naringenin Regulates Expression of Genes Involved in Cell Wall Synthesis in *Herbaspirillum seropedicae*[∇]

M. Z. Tadra-Sfeir,¹ E. M. Souza,¹ H. Faoro,¹ M. Müller-Santos,¹ V. A. Baura,¹ T. R. Tuleski,¹
L. U. Rigo,¹ M. G. Yates,¹ R. Wasseem,² F. O. Pedrosa,¹ and R. A. Monteiro^{1*}

*Department of Biochemistry and Molecular Biology, Universidade Federal do Paraná, Curitiba, PR, Brazil,¹ and
Department of Genetics, Universidade Federal do Paraná, Curitiba, PR, Brazil²*

Received 1 September 2010/Accepted 14 January 2011

Five thousand mutants of *Herbaspirillum seropedicae* SmR1 carrying random insertions of transposon pTnMod-OGmKmlacZ were screened for differential expression of LacZ in the presence of naringenin. Among the 16 mutants whose expression was regulated by naringenin were genes predicted to be involved in the synthesis of exopolysaccharides, lipopolysaccharides, and auxin. These loci are probably involved in establishing interactions with host plants.

The betaproteobacterium *Herbaspirillum seropedicae* is an endophytic diazotroph that forms nitrogen-fixing associations with maize (*Zea mays*), rice (*Oryza sativa*), sorghum (*Sorghum bicolor*), and sugar cane (*Saccharum officinarum*), as well as such diverse plants as bananas (*Musa* spp.) and pineapple (*Ananas comosus*) (1, 5, 27). As such, *H. seropedicae* is a potential nitrogen biofertilizer that also produces phytohormones that may stimulate the growth of plants (4). Studies demonstrated that the inoculation of rice with *H. seropedicae* promoted a yield increase equivalent to treatment with 40 kg N/ha (2). Coinoculation of micropropagated sugar cane with *Gluconacetobacter diazotrophicus* and *Herbaspirillum* sp. resulted in an increase of the total plant biomass (25).

Exactly how *H. seropedicae* colonizes Gramineae and other plants is not known, and there is even less information available on the role of plant metabolites in the regulation of bacterial invasion and colonization of the inner tissues. That such associations involve molecular communication between the host plant and bacteria, resulting in modified patterns of gene expression, is clear from studies of other rhizospheric bacteria (15). In legume-*Rhizobium* interactions, flavonoids released by plant roots induce sets of genes involved in nodulation. As a result, lipochitooligosaccharides are excreted and symbiotic forms of exopolysaccharides (EPSs), lipopolysaccharides (LPSs), and glucans synthesized, all of which modulate the nodulation process (8, 15, 26). Previous studies showed that naringenin (50 μ mol/liter) stimulated the root colonization of *Arabidopsis thaliana* by *H. seropedicae* (16) and the intercellular colonization of wheat roots by *Azorhizobium caulinodans* (33).

To identify bacterial genes whose expression is under the control of the flavonoid naringenin, the chromosome of *H. seropedicae* strain SmR1 (23) was randomly mutagenized using a lacZ-Km-Gm cassette carried by the plasmid pTnMod-OGmKmlacZ, and strains resistant to both kanamycin and

gentamicin were selected (30). Twelve thousand mutant strains were obtained, and 5,000 screened for differential expression of the promoterless lacZ reporter gene in the presence of naringenin (50 μ mol/liter) using NFbHP-malate-agar medium (19) containing X-Gal (5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside) (30 μ g/ml). One hundred ninety-six mutants bearing mutated genes potentially controlled by naringenin were preselected, and their β -galactosidase activities determined (7, 24). Of these, 16 isolates carried mutated genes that reacted differently to the presence of naringenin: 4 of the mutants carried genes that were upregulated, and 12 carried genes that were downregulated by naringenin (Table 1). The mutated genes of these strains were identified by sequencing as described previously (30).

The fraction of the genome covered by the set of insertion mutants was calculated according to the method of Jacobs et al. (17). The total length of the *H. seropedicae* genome is 5,513,887 bp, containing 4,737 open reading frames (ORFs) with an average length of 1,029 bp. Since a total of 5,000 mutants were tested and assuming that half of these (2,500) would have the lacZ gene inserted in the mutated gene transcription orientation, approximately 38% of the genome was covered by the mutant collection screened (17).

Among the mutants in genes downregulated (2- to 3-fold) by naringenin, four were affected in genes probably involved in the synthesis of cell wall components. Strain MHS01 was mutated in *epsG*, which is involved in exopolysaccharide biosynthesis; MHS02 in a gene for a probable O-antigen acetylase; MHS05 in *ampG*, which codes for a muropeptide permease of the major facilitator superfamily; and MHS15 in a gene coding for a probable glucosyl transferase bearing low similarity to GumH, a glucosyl transferase from *Xylella fastidiosa* that is involved in exopolysaccharide biosynthesis. Interestingly, strain MHS03 was mutated in a gene coding for a conserved protein with no function assigned but located in an operon together with two genes coding for putative outer membrane proteins. One of these is a probable outer membrane porin of the OmpA family, suggesting that the mutated gene may also play a role in outer membrane structure. LPSs consist of an O-antigen, a lipid A, and core oligosaccharides

* Corresponding author. Mailing address: Department of Biochemistry and Molecular Biology, Universidade Federal do Paraná, C. Postal 19046, CEP81531980, Curitiba, PR, Brazil. Phone: 55 41 3361 1659. Fax: 55 41 3361 1578. E-mail: roseadele@ufpr.br.

[∇] Published ahead of print on 21 January 2011.

TABLE 1. Identification of genes from *H. seropedicae* that are regulated by the flavonoid naringenin

Mutant strain ^a	Accession no. of the mutated locus	Deduced gene product ^b	Relative naringenin regulation (avg expression ± SD) shown by:	
			β-Gal activity ^c	RT-PCR ^d
MHS01	YP_003775401.1	EPS biosynthesis protein (EpsG)	Downregulated (0.31 ± 0.05)	Downregulated (0.07 ± 0.045)
MHS02	YP_003775614	Probable O-antigen acetylase	Downregulated (0.34 ± 0.09)	Downregulated (0.23 ± 0.11)
MHS03	YP_003773926.1	Conserved hypothetical protein	Downregulated (0.50 ± 0.01)	Downregulated (0.52 ± 0.082)
MHS04	YP_003774406	Probable acyltransferase protein	Downregulated (0.44 ± 0.07)	Downregulated (0.43 ± 0.047)
MHS05	YP_003773531.1	Probable muropeptide permease of the major facilitator superfamily protein (AmpG)	Downregulated (0.33 ± 0.05)	Downregulated (0.12 ± 0.043)
MHS06	YP_003778011.1	Probable acyl CoA:acetate/3-ketoacid CoA transferase, beta subunit protein	Upregulated (1.65 ± 0.21)	Upregulated (1.23 ± 0.012)
MHS07	YP_003774121.1	Probable tRNA (5-methylaminomethyl-2-thiouridylate)-methyltransferase protein (TrmU)	Downregulated (0.66 ± 0.11)	Downregulated (0.24 ± 0.063)
MHS08	YP_003776865.1	Hypothetical protein	Upregulated (1.66 ± 0.07)	Upregulated (1.13 ± 0.09)
MHS09	YP_003775366.1	Probable 5-aminolevulinic acid synthase protein	Upregulated (2.04 ± 0.03)	Downregulated (0.93 ± 0.046)
MHS10	YP_003776866.1	Hypothetical protein	Downregulated (0.18 ± 0.001)	Downregulated (0.71 ± 0.01)
MHS11	YP_003778130.1	Conserved hypothetical protein	Upregulated (1.22 ± 0.015)	Downregulated (0.81 ± 0.056)
MHS12	YP_003775661.1	Probable NAD-dependent aldehyde dehydrogenase protein	Downregulated (0.27 ± 0.03)	Downregulated (0.258 ± 0.075)
MHS13	YP_003777134.1	Hypothetical protein	Downregulated (0.54 ± 0.12)	Downregulated (0.21 ± 0.038)
MHS14	YP_003777655.1	Probable indole pyruvate ferredoxin oxidoreductase, alpha and beta subunits protein	Downregulated (0.042 ± 0.02)	Downregulated (0.15 ± 0.015)
MHS15	YP_003777590.1	Glucosyltransferase hypothetical protein	Downregulated (0.54 ± 0.12)	Downregulated (0.23 ± 0.04)
MHS16	YP_003775234.1	Conserved hypothetical protein	Downregulated (0.12 ± 0.04)	Downregulated (0.17 ± 0.053)

^a Sixteen insertion mutants of *H. seropedicae* SmR1 carry genes whose expression is modulated by naringenin.

^b BLASTn and SMART comparisons of the genes with publicly available databases (accessed on 28 July 2010) are shown.

^c Expression levels were measured as β-galactosidase (β-Gal)-specific activities of the parental strain and *lacZ* fusion mutants, and relative expression is reported as the ratio of β-galactosidase activity in the presence of 50 μmol/liter naringenin to β-galactosidase activity in the absence of naringenin.

^d Relative expression was determined by the 2^{-ΔΔCT} (threshold cycle) method (22); the relative expression of the *epsB* gene located in the same operon as *epsG* was 0.082 ± 0.01. The results represent three replicate experiments. The RT-PCR experiments were performed using Power SYBR green master mix (Applied Biosystems, Carlsbad, CA), and reactions were run on a Step One Plus real-time PCR system (Applied Biosystems).

and are present as a monolayer on the outer membrane of Gram-negative bacterial cells. Strain MHS02 carries a mutation that may affect the acetylation of the O-antigen part of the LPS of *H. seropedicae*. Similar changes to the O-antigen of rhizobia result in altered symbiotic phenotypes (9, 21).

Changes in bacterial outer surface structures, such as LPS and EPS, often result in increased sensitivity to cationic peptide antibiotics, detergents, and various stresses and affect symbiotic development (9, 10, 18, 20, 21, 31). Accordingly, we challenged the collection of mutants with increasing concentrations of sodium dodecyl sulfate (SDS). In contrast to the parental strain, isolates with mutations in genes related to the synthesis of EPS (MHS01) and muropeptide permease (MHS05) were unable to grow in the presence of 0.5% (wt/vol) SDS (data not shown). Since mutants affected in LPS synthesis had an altered sensitivity to SDS, we analyzed the LPS profiles of isolates MHS01 and MHS05. LPSs were extracted using the phenol procedure (13) and separated on denaturing polyacrylamide gels (20). Only mutation of the muropeptide permease gene caused significant changes to the LPS profile (Fig. 1A). Although the band pattern is similar to that of the parental strain, the amount of LPS recovered per cell of strain MSH05 is clearly reduced, suggesting that LPS may not be qualitatively different but is present in a lower concentration in the *ampG* strain's outer membrane. The AmpG protein is located in the




FIG. 1. LPS profiles and colonization of the roots of *Zea mays* by the parental *H. seropedicae* strain SmR1 and *ampG* mutant MHS05. (A) Sodium dodecyl sulfate-PAGE electrophoresis patterns of LPS isolated from the parental strain (lane 1) and the mutant derivative (lane 2) grown in NfbHPN-malate medium. (B) Colonization of *Z. mays* roots. Results are shown as the average numbers of bacteria · g⁻¹ fresh root ± standard deviations. The values above the columns represent the difference in colonization between parental and mutant strains. Asterisks indicate statistically (Student *t* test) significant differences between the parental strain and the *ampG* mutant in colonization of maize roots. *, *P* < 0.005; **, *P* < 0.01.

membrane of Gram-negative bacteria and is necessary for cell wall peptidoglycan recycling (11). The peptidoglycan is both a highly complex and essential macromolecule of bacterial cell walls (6). The results suggest that the lack of AmpG in strain MSH05 leads to an anomaly in the peptidoglycan layer affecting the synthesis or targeting of LPS in *H. seropedicae*.

To check whether any of the mutations influenced the colonization of maize (*Zea mays*), the protocol of Balsanelli et al. (3) was used to test bacterial cell adhesion to root surface and internal colonization. No difference was observed between the parental strain (SmR1) and mutants in growth and root surface attachment (data not shown). On the other hand, the endophytic population of MSH05 was reduced by 10-fold compared to that of the parental strain both 1 and 3 days after inoculation, suggesting that the modification in LPS in this strain is important for bacterial establishment in plant tissues (Fig. 1B). Previously, Balsanelli et al. (3) showed that strains unable to synthesize LPS were severely impaired in maize colonization. These authors also showed that the *mlb* gene, whose product is involved in LPS biosynthesis, was upregulated by naringenin (4-fold), in agreement with our results showing that naringenin can modulate genes involved in the biosynthesis and modification of LPS. All other strains had the same pattern of colonization as the parental strain (data not shown).

Genes coding for a putative methyltransferase (MHS07), a probable acyltransferase (MHS04), an aldehyde dehydrogenase (MHS12), and indole pyruvate ferredoxin oxidoreductase (MHS14) were also downregulated by naringenin. Since the mutated indole pyruvate ferredoxin oxidoreductase in strain MHS14 may be involved in indole acetic acid (IAA) biosynthesis, the results suggest that naringenin may modulate indole acetic acid synthesis in *H. seropedicae*. The relationship of plant flavonoid and auxin production was also observed in *Azospirillum brasilense*, where mutation of *ipdC* of the IAA biosynthetic pathway caused a reduction of exudation of flavonoids by maize roots (14), and flavonoids induce IAA production in *Rhizobium* NGR234 (32). In the latter organism, the genes *y4wE* (class 2 amino transferase) or *yw4F* (monooxygenase) are regulated by flavonoids through NodD.

Four strains were mutated in genes upregulated by naringenin. These include those coding for a putative acyl coenzyme A (CoA):acetate/3-ketoacid CoA transferase (MHS06), a hypothetical protein (MHS08), an aminotransferase similar to 5-aminolevulinic acid synthase (MHS09), and a MarR transcription regulator that may be involved in multidrug resistance (MHS11). The deduced product of the gene mutated in strain MHS08 is similar to a protein of *Vibrio cholera* of the type II secretory pathway. This system is used by many Gram-negative bacteria, mainly to secrete extracellular enzymes that are associated with the degradation of host tissue (28, 29). The gene coding for the aminotransferase (MHS09) is located in an operon together with a gene coding for a permease of the MFS family, which is possibly related to the tetracenomycin C resistance protein. Cho et al. (12) identified a similar multidrug resistance permease gene of *Escherichia coli* that is upregulated by an infusion containing polyphenols. In strain MHS16, the gene mutated is located in an operon together with another acetyltransferase, an organization that is conserved in *Burkholderia vietnamiensis*. Thus, altogether, at least three different

genes coding for acyltransferases seem to have their expression affected by naringenin.

Four other mutations were in hypothetical genes of *H. seropedicae* whose products had no significant matches in all databases (MHS03, MHS10, MHS13, and MHS16). The genes mutated in strains MHS03 and MHS10 probably code for membrane proteins.

Real-time PCR analysis was used to validate the data obtained with *lacZ* fusions. To do this, total RNA was isolated from cultures grown in the presence and absence of naringenin (50 μ mol/liter) using the Trizol procedure (Sigma Corp., St. Louis, MO). Two micrograms of RNA was used to synthesize cDNA with a high-capacity cDNA reverse transcription kit (Applied Biosystems, Carlsbad, CA). The 16S rRNA gene was used as the internal control. Primers for the gene *epsB* located in the same operon as *epsG* were also used, to confirm the coregulation of genes related to EPS synthesis.

The RT-PCR analyses confirmed regulation of the mutated genes by naringenin except for strains MHS08, MHS09, and MHS11 (Table 1). In these three strains, the β -galactosidase activity indicated upregulation by naringenin, whereas RT-PCR indicated no regulation (MHS08 and MHS09) or even slight downregulation (MHS11), raising the possibility of post-transcriptional regulation of these genes. Finally, the *epsB* and *epsG* genes responded similarly to naringenin, suggesting that the *eps* operon is regulated by the flavonoid.

Our results show that *H. seropedicae* responds to the plant flavonoid naringenin by modifying the pattern of gene expression. The products of four genes with altered expression seem to be involved in the synthesis of the outer membrane of the cell wall. Changes in the cell surface are documented in other bacteria that interact with plants (e.g., rhizobia) (9, 20, 31) and probably play a role in the interaction between *H. seropedicae* and its host plants. Furthermore, the data indicate that naringenin may also regulate genes involved in phytohormone production in *H. seropedicae*.

This work was supported by the Brazilian agencies CAPES and INCT-FBN/CNPq.

We thank Roseli Prado and Julieta Pie for technical assistance.

REFERENCES

- Baldani, J. I., V. L. D. Baldani, L. Seldin, and J. Döbereiner. 1986. Characterization of *Herbaspirillum seropedicae* gen. nov., sp. nov., a root-associated nitrogen-fixing bacterium. *Int. J. Syst. Bacteriol.* **33**:167–172.
- Baldani, J. I., and V. L. D. Baldani. 2005. History on the biological nitrogen fixation research in graminaceous plants: special emphasis on the Brazilian experience. *An. Acad. Bras. Cienc.* **77**:549–579.
- Balsanelli, E., et al. 2010. *Herbaspirillum seropedicae* *rfbB* and *rfbC* genes are required for maize colonization. *Environ. Microbiol.* **12**:2233–2244.
- Bastián, F., et al. 1998. Production of indole-3-acetic acid and gibberellins A₁ and A₃ by *Acetobacter diazotrophicus* and *Herbaspirillum seropedicae* in chemically-defined culture media. *Plant Growth Regul.* **24**:7–11.
- Boddey, R. M., et al. 1995. Biological nitrogen fixation associated with sugar cane and rice: contributions and prospects for improvement. *Plant Soil* **74**:195–209.
- Boneca, I. G. 2005. The role of peptidoglycan in pathogenesis. *Curr. Opin. Microbiol.* **8**:46–53.
- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**:248–254.
- Broughton, W. J., S. Jabbouri, and X. Perret. 2000. Keys to symbiotic harmony. *J. Bacteriol.* **182**:5641–5652.
- Broughton, W. J., et al. 2006. Flavonoid-inducible modifications to rhamnan O antigens are necessary for *Rhizobium* sp. strain NGR234-legume symbiosis. *J. Bacteriol.* **188**:3654–3663.
- Campbell, G. R. O., B. L. Reuhs, and G. C. Walker. 2002. Chronic intra-

- cellular infection of alfalfa nodules by *Sinorhizobium meliloti* requires correct lipopolysaccharide core. Proc. Natl. Acad. Sci. U. S. A. **99**:3928–3943.
11. **Chahboune, A., M. Decaffmeyer, R. Brasseur, and B. Joris.** 2005. Membrane topology of the *Escherichia coli* AmpG permease required for recycling of cell wall anhydromuropeptides and AmpC beta-lactamase induction. Antimicrob. Agents Chemother. **49**:1145–1149.
 12. **Cho, Y. S., N. L. Schiller, H. Y. Kahng, and K. H. Oh.** 2007. Cellular responses and proteomic analysis of *Escherichia coli* exposed to green tea polyphenols. Curr. Microbiol. **55**:501–506.
 13. **Currie, C. G., and I. R. Poxton.** 1999. The lipopolysaccharide core type of *Escherichia coli* O157:H7 and other non-O157 verotoxin-producing *E. coli*. FEMS Immunol. Med. Microbiol. **24**:57–62.
 14. **Dardanelli, M. S., et al.** 2008. Effect of *Azospirillum brasilense* coinoculated with *Rhizobium* on *Phaseolus vulgaris* flavonoids and Nod factor production under salt stress. Soil Biol. Biochem. **40**:2713–2721.
 15. **Deakin, W. J., and W. J. Broughton.** 2009. Symbiotic use of pathogenic strategies: rhizobial protein secretion systems. Nat. Rev. Microbiol. **7**:312–320.
 16. **Gough, C., et al.** 1997. Specific flavonoids promote intercellular root colonization of *Arabidopsis thaliana* by *Azorhizobium caulinodans* ORS571. Mol. Plant Microbe Interact. **10**:560–570.
 17. **Jacobs, M. A., et al.** 2003. Comprehensive transposon mutant library of *Pseudomonas aeruginosa*. Proc. Natl. Acad. Sci. U. S. A. **100**:14339–14344.
 18. **Jofré, E., A. Lagares, and G. Mori.** 2004. Disruption of dTDP-rhamnose biosynthesis modifies lipopolysaccharide core, exopolysaccharide production and root colonization in *Azospirillum brasilense*. FEMS Microbiol. Lett. **231**:267–275.
 19. **Klassen, G., F. O. Pedrosa, E. M. Souza, S. Funayama, and L. U. Rigo.** 1997. Effect of nitrogen compounds on nitrogenase activity in *Herbaspirillum seropedicae* SMR1. Can. J. Microbiol. **43**:887–891.
 20. **Le Quéré, A. J., et al.** 2006. Structural characterization of a K-antigen capsular polysaccharide essential for normal symbiotic infection in *Rhizobium* sp. NGR234. Deletion of the *rkpMNO* locus prevents synthesis of 5,7-diacetamido-3,5,7,9-tetradeoxy-non-2-ulonic acid. J. Biol. Chem. **281**:28981–28992.
 21. **Lerouge, I., and J. Vanderleyden.** 2002. O-antigen structural variation: mechanisms and possible roles in animal/plant-microbe interactions. FEMS Microbiol. Rev. **26**:17–47.
 22. **Livak, K. J., and T. D. Schmittgen.** 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-ΔΔC_T} method. Methods **25**:402–408.
 23. **Machado, I. M., M. G. Yates, H. B. Machado, E. M. Souza, and F. O. Pedrosa.** 1996. Cloning and sequencing of the nitrogenase structural genes *nifHDK* of *Herbaspirillum seropedicae*. Braz. J. Med. Biol. Res. **29**:1599–1602.
 24. **Miller, J. H.** 1972. Experiments in molecular genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
 25. **Pedraza, R. O.** 2008. Recent advances in nitrogen-fixing acetic acid bacteria. Int. J. Food Microbiol. **125**:25–35.
 26. **Perret, X., C. Staehelin, and W. J. Broughton.** 2000. Molecular basis of symbiotic promiscuity. Microbiol. Mol. Biol. Rev. **64**:180–201.
 27. **Pimentel, J. P., et al.** 1991. Dinitrogen fixation and infection of grass leaves by *Pseudomonas rubrisubalbicans* and *Herbaspirillum seropedicae*. Plant Soil **137**:61–65.
 28. **Pugsley, A. P.** 1993. The complete general secretory pathway in gram-negative bacteria. Microbiol. Rev. **57**:50–108.
 29. **Sandkvist, M.** 2001. Biology of type II secretion. Mol. Microbiol. **40**:271–283.
 30. **Schwab, S., et al.** 2007. Identification of NH₄⁺-regulated genes of *Herbaspirillum seropedicae* by random insertional mutagenesis. Arch. Microbiol. **187**:379–386.
 31. **Staehelin, C., et al.** 2006. Exo-oligosaccharides of *Rhizobium* sp. NGR234 are required for symbiosis with various legumes. J. Bacteriol. **188**:6168–6178.
 32. **Theunis, M., H. Kobayashi, W. J. Broughton, and E. Prinsen.** 2004. Flavonoids, NodD1, NodD2 and *nod*-box NB15 modulate expression of the *y4wEFG* locus that is required for indole-3-acetic acid synthesis in *Rhizobium* sp. NGR234. Mol. Plant Microbe Interact. **17**:1153–1161.
 33. **Webster, G., et al.** 1998. The flavonoid naringenin stimulates the intercellular colonization of wheat roots by *Azorhizobium caulinodans*. Plant Cell Environ. **21**:373–383.