

Genetic Analysis of the *Cronobacter sakazakii* O4 to O7 O-Antigen Gene Clusters and Development of a PCR Assay for Identification of All *C. sakazakii* O Serotypes

Yamin Sun,^{a,b,c} Min Wang,^{a,b,c} Quan Wang,^{a,b,c} Boyang Cao,^{a,b,c} Xin He,^{a,b,c} Kun Li,^{a,b,c} Lu Feng,^{a,b,c} and Lei Wang^{a,b,c,d}

The Key Laboratory of Molecular Microbiology and Technology of the Ministry of Education, Nankai University, Tianjin, China^a; TEDA School of Biological Sciences and Biotechnology, Nankai University, TEDA, Tianjin, China^b; Tianjin Key Laboratory of Microbial Functional Genomics, TEDA, Tianjin, China^c; and Tianjin Biochip Corporation, TEDA, Tianjin, China^d

The Gram-negative bacterium *Cronobacter sakazakii* is an emerging food-borne pathogen that causes severe invasive infections in neonates. Variation in the O-antigen lipopolysaccharide in the outer membrane provides the basis for Gram-negative bacteria serotyping. The O-antigen serotyping scheme for *C. sakazakii*, which includes seven serotypes (O1 to O7), has been recently established, and the O-antigen gene clusters and specific primers for three *C. sakazakii* serotypes (O1, O2, and O3) have been characterized. In this study, the *C. sakazakii* O4, O5, O6, and O7 O-antigen gene clusters were sequenced, and gene functions were predicted on the basis of homology. *C. sakazakii* O4 shared a similar O-antigen gene cluster with *Escherichia coli* O103. The general features and anomalies of all seven *C. sakazakii* O-antigen gene clusters were evaluated and the relationship between O-antigen structures and their gene clusters were investigated. Serotype-specific genes for O4 to O7 were identified, and a molecular serotyping method for all *C. sakazakii* O serotypes, a multiplex PCR assay, was developed by screening against 136 strains of *C. sakazakii* and closely related species. The sensitivity of PCR-based serotyping method was determined to be 0.01 ng of genomic DNA and 10³ CFU of each strain/ml. This study completes the elucidation of *C. sakazakii* O-antigen genetics and provides a molecular method suitable for the identification of *C. sakazakii* O1 to O7 strains.

Cronobacter (formerly *Enterobacter sakazakii*) was reclassified as a genus, and now seven named species—*Cronobacter sakazakii*, *C. malonaticus*, *C. muytjensii*, *C. dublinensis*, *C. turicensis*, *C. condimenti*, and *C. universalis*—have been found (20, 21, 23a). *Cronobacter* is an emerging opportunistic pathogen causing severe invasive infections in neonates (11, 15, 35). Clinical strains of *Cronobacter* spp. have been linked to several outbreaks of neonatal meningitis and necrotizing enterocolitis, leading to a high mortality rate (estimated between 33 and 80%) in vulnerable infants (15, 28). *C. sakazakii* appears to be the most important species in this genus for its dominant isolation frequency and important clinical significance (3, 23, 27, 33).

The O antigen forms part of the lipopolysaccharide (LPS) in the outer membrane of Gram-negative bacteria and is one of the most variable constituents on the cell surface (40). Its variation provides the basis for the serotyping schemes of Gram-negative bacteria. Since it was shown that specific O serotypes were associated with pathogens causing enteritis epidemics in newborn infants between 1945 and 1950, serotyping has been the most widely used method for identifying strains for epidemiological purposes. The O-antigen serotyping scheme for *C. sakazakii*, which includes seven serotypes (O1 to O7), was recently established for the first time using traditional immunological technologies (46). Several O serotypes were also designed for other *Cronobacter* species, such as *C. muytjensii*, *C. malonaticus*, and *C. turicensis* (22).

Serotyping is still widely used to characterize isolates for monitoring outbreaks and general bacterial surveillance (17), although other useful molecular typing approaches, such as pulsed-field gel electrophoresis, multilocus sequence typing, and repetitive sequence-based PCR, have been developed (3, 6, 16, 27). Traditional serotyping technology using antisera is specific and still routinely used; however, this approach is labor-intensive, and antisera for

all serogroups are difficult to obtain. Therefore, rapid molecular methods for *C. sakazakii* serotyping are required.

Genes for O-antigen synthesis are normally located on the chromosome as an O-antigen gene cluster, and genetic variation in the gene cluster is the major basis for the diversity of O-antigen forms (40). Similar to *Escherichia coli*, the *C. sakazakii* O-antigen gene cluster is located between *galF* and *gnd* (34). Furthermore, a conserved 39-bp JUMPstart sequence required for the regulation of downstream genes, is located in the intergenic region between *galF* and the O-antigen gene cluster (18, 32). Three main classes of genes, including nucleotide sugar synthesis genes, sugar transferase genes, and O-unit processing genes, have been identified in O-antigen gene clusters (40). Particular O-antigen genes, such as O-unit processing genes, are highly specific to individual O serogroups and are used in molecular assay for rapid identification and detection of *E. coli*, *Shigella*, and *Salmonella* relevant strains (7, 8, 14). To date, O-antigen gene clusters and specific primers for three *C. sakazakii* serotypes (O1, O2, and O3) have been characterized (22, 34). The O-antigen gene clusters of O5 to O7 were referred to in our previously publications but were not analyzed in detail (2, 44).

In the present study, the O-antigen gene clusters of the remaining *C. sakazakii* O4, O5, O6, and O7 were sequenced and analyzed,

Received 10 December 2011 Accepted 15 March 2012

Published ahead of print 23 March 2012

Address correspondence to Lei Wang, wanglei@nankai.edu.cn.

Y.S. and M.W. contributed equally to this article.

Supplemental material for this article may be found at <http://aem.asm.org/>.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AEM.07825-11

TABLE 1 Sources of strains used in this study

Bacterium	No. of strains and source ^a	No. of strains
<i>Cronobacter sakazakii</i>	3 ^A , 2 ^B , 39 ^C , 3 ^D , 4 ^E , 2 ^F , 2 ^G , 3 ^H , 4 ^I , 4 ^J , 5 ^K , 48 ^L	119
O1	1 ^A , 5 ^C , 2 ^D , 2 ^E , 1 ^F , 1 ^G , 1 ^H , 2 ^I , 2 ^K , 32 ^L	49
O2	2 ^A , 2 ^B , 16 ^C , 1 ^E , 4 ^J , 2 ^K , 7 ^L	34
O3	7 ^C , 1 ^K	8
O4	3 ^C , 1 ^F , 4 ^L	8
O5	3 ^C , 1 ^I	4
O6	1 ^C , 1 ^I , 6 ^L	8
O7	4 ^C , 1 ^D , 1 ^E , 1 ^G , 1 ^H	8
Other <i>Cronobacter</i> spp.	1 ^A , 2 ^E , 2 ^I , 1 ^L , 1 ^M	2
<i>C. malonaticus</i> O1 ^b	2 ^E	2
<i>C. muytjensii</i>	1 ^A	1
<i>C. dublinensis</i>	1 ^I , 1 ^L	2
<i>C. turicensis</i>	1 ^I , 1 ^M	2
<i>Enterobacter cloacae</i>	1 ^A , 1 ^N	2
<i>Enterobacter aerogenes</i>	2 ^N	2
<i>Escherichia coli</i>	1 ^A , 1 ^O	2
O111	1 ^A	1
O157	1 ^O	1
<i>Salmonella enterica</i>	1 ^A , 1 ^O	2
O44	1 ^A	1
D1	1 ^O	1
<i>Shigella flexneri</i>	1 ^A , 1 ^L	2
2a	1 ^A	1
5a	1 ^L	1
Total		136

^a Sources are indicated by superscript letters as follows: A, American Type Culture Collection, USA; B, Czech Collection of Microorganisms (CCM), Czech Republic; C, environmental isolates from food by Tianjin Entry-Exit Inspection and Quarantine Bureau, China; D, environmental isolates from food by Liaoning Entry-Exit Inspection and Quarantine Bureau, China; E, Chinese Academy of Inspection and Quarantine, Beijing, China; F, environmental isolates from food by Jilin Entry-Exit Inspection and Quarantine Bureau, China; G, environmental isolates from food by Neimenggu Entry-Exit Inspection and Quarantine Bureau, China; H, environmental isolates from food by Shenyang Entry-Exit Inspection and Quarantine Bureau, China; I, environmental isolates from food by Xinjiang Entry-Exit Inspection and Quarantine Bureau, China; J, environmental isolates from food by Guangdong Entry-Exit Inspection and Quarantine Bureau, China; K, environmental isolates from food by Hunan Entry-Exit Inspection and Quarantine Bureau, China; L, environmental isolates from food by Beijing Entry-Exit Inspection and Quarantine Bureau, China; M, Environmental isolates from food by Hubei Entry-Exit Inspection and Quarantine Bureau, China; N, China General Microbiological Culture Collection Center (CGMCC), China; and O, National Center for Medical Culture Collection (CMCC), China.

^b Based on previously published data (22).

providing the possibility of a systematic characterization of the genetics of this pathogen. Specific genes from the four serotypes (O4 to O7) were also identified, thus allowing development of molecular serotyping assay for all seven *C. sakazakii* serotypes in combination with the previously identified O1 to O3 specific genes.

MATERIALS AND METHODS

Bacterial strains. Bacterial strains used in the present study are listed in Table 1, and all of these strains have been identified in our previous study (46). *C. sakazakii* type strains O4, O5, O6, and O7 (laboratory stock numbers G2594, G2706, G2704, and G2592) were isolated from powdered

infant formula collected from India, Ireland, China, and France by the Tianjin Entry-Exit Inspection and Quarantine Bureaus of China in 2006 and 2007 (46).

Genomic DNA extraction and O-antigen gene cluster amplification.

Bacterial genomic DNA was extracted from each strain by using bacterial genomic DNA purification kits (Tiangen Biotech Co., Ltd., Beijing, China) according to the instructions provided by the manufacturer. Long-range PCR of the O-antigen gene cluster was performed with an Expand Long Template PCR system (TaKaRa Biotechnology, Shiga, Japan) using the following primers for amplification of the JUMPStart sequence and *gnd* gene, respectively: wl-10324 (5'-GCACTGGTAGCTATTGAGCCAGGGGCGGTAGCAT-3') and wl-2211 (5'-ACTGCCATACCGACGACGCCGATCTGTTGCTTGG-3'). The PCR amplification was performed in 50- μ l volumes containing 1 \times LA buffer (plus MgCl₂), 0.4 mM concentrations each dATP, dCTP, dGTP, and dTTP, 0.4 μ M concentrations of each primer, 2.5 U of TakaRa LA *Taq* polymerase, and 2 μ l of template DNA (~500 ng). The following PCR conditions were used for amplification: 32 cycles of denaturation at 94°C for 30 s, annealing at 61°C for 45 s, and extension at 68°C for 15 min.

Sequencing and analysis of O-antigen gene clusters. The PCR products were gel purified using UNIQ-10 columns (Sangon, Shanghai, China) and were sheared at speed code 9 (20 cycles) to an estimated fragment length of 1,000 to 2,000 bp by HydroShear fragmentation (DigiLab, CA). The resulting DNA fragments were repaired and then cloned into the pUC18 vector to produce a shotgun bank. Sequencing was carried out using an ABI 3730 automated DNA sequencer (Applied Biosystems, Foster City, CA). Sequencing data were assembled using the Staden Package software (45) and annotated with the Artemis program (41). BLAST analysis was used to search available databases such as GenBank (5). The Clusters of Orthologous Groups (COG) (48) and Pfam (9) protein motif databases were used to search for conserved protein domains. The program TMHMM 2.0 (25) was used to identify potential transmembrane segments. Sequence alignments and comparisons were performed using the CLUSTAL W program (49).

PCR primer specificity analysis. Primers were designed based on the *wzx* and *wzy* gene sequences using Primer Premier 5.0 software program (Premier Biosoft International, Palo Alto, CA) (Table 2). The sequences of the primers were compared to the sequences in GenBank and the unpublished O-antigen gene sequences in our own laboratory to ensure their specificity. Genomic DNAs extracted from the strains listed in Table 1 were used as templates for amplification. The multiplex PCR was performed by mixing all primers in a final volume of 50 μ l containing the following components: 1 \times PCR buffer; 2.5 mM MgCl₂; 400 μ M concentrations (each) of dATP, dCTP, dGTP, and dTTP; 0.06 to 0.10 μ M primer sets listed in Table 2; and 2.5 U of *Taq* DNA polymerase and DNA template (50 to 100 ng). The following PCR conditions were used for amplification: an initial denaturation step at 95°C for 5 min, followed by 30 cycles of 94°C for 30 s, 53°C for 30 s, and 72°C for 1 min, with a final extension at 72°C for 5 min. Samples of the PCR products (5 μ l) were subjected to agarose gel electrophoresis for examination.

PCR assay sensitivity analysis. The PCR assay sensitivity was analyzed as follows. Serial dilutions of *C. sakazakii* O1 to O7 strain (laboratory stock numbers G2539, G2356, G2726, G2594, G2706, G2704, and G2592) genomic DNA (10, 1, 0.1, 0.01, 0.001, 0.0001, and 0.00001 ng) were used as templates to test the sensitivity of the multiplex PCR assay. The sensitivity for isolates in pure culture was also tested. The *C. sakazakii* O1 to O7 strains were grown in Luria-Bertani (LB) medium to log phase and serially diluted 10-fold in sterile water. Aliquots from the same dilution were removed for planting onto LB agar and for extraction of their genomic DNA to be used as the template for the developed PCR assay. The CFU were counted after overnight growth at 37°C. Amplified products were subjected to agarose gel electrophoresis.

Nucleotide sequence accession numbers. The DNA sequences of the *C. sakazakii* O4, O5, O6, and O7 O-antigen gene clusters have been de-

TABLE 2 Primers used in this study^a

<i>C. sakazakii</i> serotype	Target gene	Primer	Sequence (5'-3')	Final concn (μM)	Amplicon size (bp)
O1	<i>wzy</i>	wl-35646	CCCGCTTGTATGGATGTT	0.10	364
		wl-35647	CTTTGGGAGCGTTAGGTT	0.10	
O2	<i>wzy</i>	wl-37256	ATTGTTTGGCATGGTGAG	0.06	152
		wl-37257	AAAACAATCCAGCAGCAA	0.06	
O3	<i>wzy</i>	wl-37258	CTCTGTTACTCTCCATAGTGTC	0.10	704
		wl-37259	GATTAGACCACCATAGCCA	0.10	
O4	<i>wzy</i>	wl-39105	ACTATGGTTTGGCTATACTCCT	0.06	890
		wl-39106	ATTCATATCCTGCGTGCC	0.06	
O5	<i>wzy</i>	wl-39873	GATGATTTTGTAAAGCGGTCT	0.10	235
		wl-39874	ACCTACTGGCATAGAGGATAA	0.10	
O6	<i>wzy</i>	wl-40041	ATGGTGAAGGGAACGACT	0.06	424
		wl-40042	ATCCCCGTGCTATGAGAC	0.06	
O7	<i>wzx</i>	wl-40039	CATTTCAGATTATACCTTTC	0.06	615
		wl-40040	ACACTGGCGATTCTACCC	0.06	

^a Annealing temperature, 53°C.

posited in GenBank under accession numbers JQ674747, JQ674748, JQ674749, and JQ674750, respectively.

RESULTS

Sequence analysis of *C. sakazakii* O4, O5, O6, and O7 O-antigen gene clusters. The region between JUMPStart and *gnd* was sequenced. Sequences of 11,887, 12,299, 7,388, and 8,932 bp were obtained from *C. sakazakii* O4, O5, O6, and O7 type strains, respectively. A total of 12, 13, 7, and 8 open reading frames (ORFs) (excluding the *gnd* gene), including *orf4-1* to *orf4-12* for O4, *orf5-1* to *orf5-13* for O5, *orf6-1* to *orf6-7* for O6, and *orf7-1* to *orf7-8* for O7, were predicted, respectively (Fig. 1A), and functions such as sugar synthesis, sugar transfer, O-antigen processing, and other functions were predicted based on homology with proteins in available databases (Tables 3 to 6). All ORFs had the same transcriptional direction from JUMPStart to *gnd*, with GC contents ranging from 29.61 to 49.86%, which was lower than that of the *C. sakazakii* genome (57%) (26).

Nucleotide sugar biosynthesis genes. For the O4 O-antigen gene cluster, five proteins encoded by *orf4-1* to *orf4-5* share high-level identity levels (55 to 84%) with known RmlB, RmlA, FdtA, FdtC, and FdtB from *E. coli* O103. They have been reported to be involved in the synthesis of dTDP-3,6-dideoxy-3-(3-hydroxybutanoylamino)-D-galactose (dTDP-D-Fuc3NHb) in *E. coli* O103 (10). A protein encoded by *orf4-12* shares 67% identity with the *Haemophilus parainfluenzae* UDP-glucose 4-epimerase (Gne), which catalyzes the conversion of 2-acetamido-2-deoxy-D-glucose (GlcNAc) to 2-acetamido-2-deoxy-D-galactose (GalNAc) (4). Therefore, *orf4-1*, *orf4-2*, *orf4-3*, *orf4-4*, *orf4-5*, and *orf4-12* were named *rmlB*, *rmlA*, *fdtA*, *fdtC*, *fdtB*, and *gne*, respectively.

The O-antigen structures of O5, O6 and O7 have been defined (2, 44). Five sugars, including one 3-acetamido-3-deoxy-D-quinovose (D-Qui3NAc), two L-rhamnoses (L-Rha), one 3-deoxy-D-manno-2-octulosonic acid (Kdo), and one D-GlcNAc, were found in O5. Five sugars—D-GlcNAc, D-GalNAc, D-galactose (D-Gal), D-glucose (D-Glc), and Kdo—were found in O6, and five sugars, including two N-acetyl-L-fucosamines (L-FucNAc), two D-GlcNAc, and one D-galacturonic acid (D-GalA), were found in O7.

Genes involved in the biosynthesis of common sugar nucleotide precursors (GlcNAc, Glc, and Gal) were not located between the JUMPStart and *gnd* gene (43). Four proteins encoded by *orf5-1*, *orf5-2*, *orf5-3*, and *orf5-4* share high-level identity (69 to

97%) with known RmlB, RmlD, RmlA, and RmlC proteins, which are responsible for L-Rha biosynthesis (12, 13). Three proteins encoded by *orf5-5*, *orf5-6*, and *orf5-7* share identity (63 to 76%) with QdtA, QdtC, and QdtB, which are involved in the synthesis of D-Qui3NAc (37). Three proteins encoded by *orf7-4*, *orf7-5*, and *orf7-6* share 78 to 93% identity with the enzymes FnlA, FnlB, and FnlC, which catalyze L-FucNAc biosynthesis (24). Genes responsible for the synthesis of Kdo in O5, Kdo and GalNAc in O6, and GalA in O7 are not present in their O-antigen gene clusters.

Sugar transferase genes. Four, three, three, and three sugar transferase genes were identified in strains O4, O5, O6 and O7, respectively.

Proteins encoded by *orf4-7*, *orf4-8*, *orf4-10*, and *orf4-11* belong to glycosyltransferase families 1 (PF00534) and 2 (PF00535) and share between 56 to 71% identity with *E. coli* glycosyltransferases. Therefore, they were proposed to encode glycosyltransferases and named *wepD*, *wepE*, *wepF*, and *wepG*, respectively.

Two proteins encoded by *orf5-9* and *orf5-10* belong to glycosyltransferase family 2 (PF00535) and 1 (PF00534), respectively, and share identity (39 to 64%) with glycosyltransferases of other genus. Protein encoded by *orf5-12* was found to share 32% identity with the *Streptococcus thermophilus* glycosyltransferase Eps5K. Therefore, *orf5-9*, *orf5-10*, and *orf5-12* were proposed to encode glycosyltransferases and named *wepH*, *wepI*, and *wepK*, respectively.

Proteins encoded by *orf6-1* and *orf6-6* belong to glycosyltransferase family 2 (PF00535) and also share 38 and 50% identity with glycosyltransferases of other genus. A protein encoded by *orf6-5* shares 26% identity with an *Erwinia* glycosyltransferase. *orf6-1*, *orf6-5*, and *orf6-6* were then proposed to be glycosyltransferase genes and named *wepL*, *wepN*, and *wepO*, respectively.

Three proteins encoded by *orf7-2*, *orf7-3*, and *orf7-7* belong to glycosyltransferase family 1 (PF00534). Two proteins encoded by *orf7-2* and *orf7-3* share 35 to 28% identity with glycosyltransferases of other genus and are named *wepQ* and *wepR*, respectively. A protein encoded by *orf7-7* shares high-level identity with WbuB of *E. coli*, which is an L-fucosamine transferase. Therefore, the protein encoded by *orf7-7* is proposed to be responsible for L-FucNAc transfer and named WbuB.

O-antigen-processing genes. Both *wzx* and *wzy* genes were found in O4, O5, and O6 serotypes, whereas only *wzx* genes were




FIG 1 (A) Predicted genes of O-antigen gene clusters for *C. sakazakii* O1 to O7 serotypes. (B) Comparison of the O-antigen gene clusters of *C. sakazakii* O4 and *E. coli* O103. (C) O-antigen chemical structures of *C. sakazakii* O1, O2, O5, O6, and O7.

identified in O7. The putative Wzx (O-antigen flippase) from the O4 to O7 serotypes have 11 to 13 well-proportioned transmembrane, which is typical of the topological characteristics of Wzx (31). They also shared 24 to 79% identity with Wzx of other strains. The putative Wzy (O-antigen polymerase) from O4 to O6 serotypes were found to have 10 to 11 predicted transmembrane segments with large periplasmic loops of 46 to 53 amino acid residues, which is typical of the topological character of Wzy (8), and shared 24 to 51% identity values with the Wzy of other strains. Therefore, these genes were proposed to encode the O-antigen flippase and polymerase, respectively.

Additional genes identified. The function of *orf5-11* was not able to be predicted based on searches from the available databases. However, on the basis of the observation that the O antigen for O5 contains five sugars, a glycosyltransferase is absent in the gene cluster (1, 40). *orf5-11* was thus proposed as a glycosyltransferase gene and named *wepJ*. A protein encoded by *orf6-7* shares 32% identity with an *Erwinia* acetyltransferase responsible for the O-acetyl group in the O6 antigen and was named WepP. The protein encoded by *orf6-2* shares 34% identity with capsule polysaccharide biosynthesis proteins of other species, although its

function in O-antigen synthesis is not known. *orf6-2* was named *wepM*. The protein encoded by *orf7-8* shares 69% identity with the WbuC of *Shigella*, which is a proposed remnant gene product (30). Therefore, *orf7-8* was named *wbuC*.

Selection of target genes. The *wzx* and *wzy* DNA sequences of seven *C. sakazakii* serotypes were aligned using CLUSTAL W software. The *wzx* and *wzy* DNA sequences are highly divergent and serotype specific, with identity levels ranging from 30.1 to 54.6% and 28.9 to 45.0%, respectively (data not shown). Therefore, the *wzx* and *wzy* genes represent as suitable targets for the development of a *C. sakazakii* molecular serotyping scheme.

Primer specificity. A total of 136 strains, including 49, 34, 8, 8, 4, 8, and 8 *C. sakazakii* O1 to O7 strains, respectively, 2 *C. malonaticus* strains, 1 *C. muytjensii* strain, 2 *C. dublinensis* strains, 2 *C. turicensis* strains, and 10 strains of closely related species, were used to test the primer specificity (Table 1). The specificity of each primer pairs was cross-tested with other O serotype strains and the closely related strains. The results showed only the DNA from of corresponding *C. sakazakii* O serotype strains generated the expected PCR products and no PCR products were amplified from other isolates (see Fig. S1a to g in the supplemental material).

TABLE 3 Characteristics of ORFs in the *C. sakazakii* O4 O-antigen gene cluster^a

ORF	Gene	Gene position (length [aa])	G+C content (%)	Pfam conserved domain(s) (Pfam no./e value)	Similar protein(s), strain(s) (GenBank accession no.)	% Identical (aa overlap)	Putative function of protein
4-1	<i>rmlB</i>	124–1215 (363)	45.51	NAD-dependent epimerase/dehydratase family (PF01370/1.7e-78)	dTDP-glucose 4,6-dehydratase, <i>E. coli</i> O103:H2 (YP_003222430)	84 (356)	Glucose dehydratase
4-2	<i>rmlA</i>	1217–2080 (287)	42.24	Nucleotidyltransferase (PF00483/6.8e-70)	D-Glucose-1-phosphate thymidyltransferase, <i>E. coli</i> O103:H2 strain 12009 (YP_003222429)	84 (286)	Glucose-1-phosphate thymidyltransferase
4-3	<i>fdtA</i>	2082–2483 (133)	35.57	WxcM-like, C-terminal (PF05523/1.3e-54)	dTDP-6-deoxy-3,4-keto-hexulose isomerase, <i>E. coli</i> O103:H2 strain 12009 (YP_003222428)	64 (131)	Hexose isomerase
4-4	<i>fdhC</i>	2476–3009 (177)	32.95		Putative butyryltransferase, <i>E. coli</i> O103:H2 (EDV84013)	55 (177)	Butyryltransferase
4-5	<i>fdtB</i>	3014–4120 (368)	39.74	DegT/DnrJ/EryC1/StrS aminotransferase family (PF01041/4e-118)	Putative aminotransferase, <i>E. coli</i> (AAS73167)	69 (372)	Aminotransferase
4-6	<i>wzx</i>	4108–5394 (428)	35.50	Polysaccharide biosynthesis protein (PF01943/7.3e-13)	O-antigen flippase Wzx, <i>E. coli</i> O103:H2 (YP_003222425)	79 (419)	Flippase
4-7	<i>wepD</i>	5378–6370 (330)	34.44	Glycosyltransferase family 2 (PF00535/2.1e-32)	Putative glycosyltransferase, <i>E. coli</i> (EDV84144)	56 (323)	Glycosyltransferase
4-8	<i>wepE</i>	6367–7257 (297)	34.34	Glycosyltransferase family 2 (PF00535/7.6e-29)	Putative glycosyltransferase, <i>E. coli</i> O103:H2 (YP_003222423)	71 (296)	Glycosyltransferase
4-9	<i>wzy</i>	7259–8413 (384)	33.5		Putative O-antigen polymerase, <i>E. coli</i> (ABK27352)	51 (378)	Polymerase
4-10	<i>wepF</i>	8410–9507 (365)	39.7	Glycosyltransferase group 1 (PF00534/3.5e-37)	Putative glycosyltransferase, <i>E. coli</i> (AAS73172)	67 (361)	Glycosyltransferase
4-11	<i>wepG</i>	9511–10632 (374)	39.48	Glycosyltransferase group 1 (PF00534/4.5e-31)	Putative galactosyltransferase, <i>E. coli</i> O103:H2 (ABK27323)	60 (373)	Galactosyltransferase
4-12	<i>gne</i>	10684–11706 (340)	42.03	NAD-dependent epimerase/dehydratase family (PF01370/1.7e-58)	UDP-glucose 4-epimerase, <i>H. parainfluenzae</i> ATCC 33392 (ZP_08147973)	67 (337)	UDP-glucose 4-epimerase

^a aa, amino acid(s).

Development of a multiplex PCR assay for identification of all *C. sakazakii* serotypes. Primer concentrations of 0.05, 0.06, 0.08, 0.10, and 0.20 μ M were analyzed to identify the optimal PCR conditions, respectively, and the best combinations (0.10 μ M for *C. sakazakii* O1, O3, and O5 serotypes and 0.06 μ M for *C. sakazakii* O2, O4, O6, and O7 serotypes) are used and listed in Table 2. Using the optimized specific primer concentrations, all of the seven primer pairs were mixed and used to screen DNA isolated from 136 strains. The results demonstrated all of the *C. sakazakii* strains produced the expected PCR products and non-*sakazakii* isolates gave no PCR products. Representative PCR results using *C. sakazakii* O1 to O7 genomic template DNA are shown in Fig. 2a. Finally, coded DNA samples from 20 strains of *Cronobacter* spp. were randomly selected and used as templates to further evaluate the specificity and sensitivity of the multiplex PCR assay using a double-blind test. The PCR results identified 2, 2, 1, 3, 1, 2, and 2 strains as *C. sakazakii* O1 to O7, respectively, and 7 strains as other *Cronobacter* spp. These results were consistent with conventional antisera serotyping methods (46).

Sensitivity of serotyping using genomic DNA and pure culture. The sensitivity of the multiplex PCR assay for the serotyping of O1 to O7 was evaluated using a serial 10-fold dilution ranging from 0.00001 to 10 ng of genomic DNA from *C. sakazakii* O serotypes O1 to O7 as templates. Positive signals were generated using 0.01 ng of

DNA for all seven serotypes. Representative PCR results using *C. sakazakii* O6 diluted genomic template DNA are shown in Fig. 2b. Similarly, serially diluted (10^1 to 10^8 CFU/ml) pure cultures of *C. sakazakii* O serotypes O1 to O7 were used as templates, and positive signals were obtained using 10^3 CFU/ml for all seven serotypes. Representative PCR results using *C. sakazakii* O6 genomic template DNA exacted from diluted pure cultures are shown in Fig. 2c.

DISCUSSION

O-antigen gene clusters of *C. sakazakii* O4 to O7 were sequenced and analyzed in the present study. The O1 to O3 gene clusters have been identified previously. Therefore, in combination, these data complete identification of all O-antigen gene clusters among the seven serotypes, thus providing the potential for systematic analysis of all *C. sakazakii* O-antigen gene clusters.

C. sakazakii O-antigen gene clusters of all seven serotypes are located on the chromosome between the housekeeping genes *galF* and *gnd*, as are the O-antigen gene clusters of *E. coli*, *Shigella* sp., *Salmonella* sp., and *Citrobacter freundii* (52). Based on the elucidated O-antigen structures of *C. sakazakii* O1, O2, O5 to O7, and O1 to O7 antigen gene clusters, it was determined that these O antigens contain rare sugars, including D-GalNAc, D-ClcA, D-GalA, D-Qui3NAc, 3-acylamido-3,6-dideoxy-D-glucose (d-Qui3NAcyl), D-Fuc3NHb, L-Rha, Kdo, and L-FucNAc (Fig. 1C).

TABLE 4 Characteristics of ORFs in the *C. sakazakii* O5 O-antigen gene cluster^a

ORF	Gene	Gene position (length [aa])	G+C content (%)	Pfam conserved domain(s) (Pfam no./e-value)	Similar protein(s), strain(s) (GenBank accession no.)	% Identity (aa overlap)	Putative function of protein
5-1	<i>rmlB</i>	183–1265 (360)	49.86	NAD-dependent epimerase/dehydratase family (PF01370/3.6e-81)	dTDP-D-glucose-4,6-dehydratase, <i>C. sakazakii</i> (ABX51885.1)	97 (360)	Glucose dehydratase
5-2	<i>rmlD</i>	1268–2164 (298)	49.72	RmlD substrate-binding domain (PF04321/2.2e-111)	dTDP-6-deoxy-L-mannose dehydrogenase, <i>C. sakazakii</i> (ABX51886.1)	96 (295)	Mannose dehydrogenase
5-3	<i>rmlA</i>	2214–3092 (292)	47.55	Nucleotidyltransferase (PF00483/6.5e-73)	Glucose-1-phosphate thymidyltransferase, <i>C. sakazakii</i> (ABX51887.1)	94 (292)	Glucose-1-phosphate thymidyltransferase
5-4	<i>rmlC</i>	3096–3644 (182)	40.98	dTDP-4-dehydrorhamnose 3,5-epimerase (PF00908/2.2e-78)	Putative dTDP-4-dehydrorhamnose 3,5-epimerase, <i>Shigella</i> sp. strain D9 (ZP_05431654.1)	69 (185)	dTDP-4-dehydrorhamnose 3,5-epimerase
5-5	<i>qdtA</i>	3647–4066 (139)	38.57	dTDP-4-dehydrorhamnose 3,5-epimerase (PF00908/2.2e-78)	QdtA <i>E. coli</i> (ADI77029)	76 (129)	dTDP-4-oxo-6-deoxy-D-glucose-3,4-oxoisomerase
5-6	<i>qdtC</i>	4041–4511 (156)	38.85	Transferase hexapeptide (PF00132/0.02)	QdtC <i>E. coli</i> (ADI77035)	68 (147)	Transacetylase
5-7	<i>qdtB</i>	4498–5607 (369)	41.17	DegT/DnrJ/EryC1/StrS aminotransferase family (PF01041/1.4e-111)	QdtB <i>E. coli</i> (ADI77030)	63 (366)	Transaminase
5-8	<i>wzx</i>	5604–6854 (416)	36.45	Polysaccharide biosynthesis protein (PF01943/2e-10)	O-antigen flippase <i>C. sakazakii</i> (ABX51878)	47 (413)	Flippase
5-9	<i>wepH</i>	6905–7807 (300)	31.67	Glycosyltransferase family 2 (PF00535/5.5e-24)	Putative glycosyltransferase <i>P. rettgeri</i> DSM 1131 (EFE51496)	39 (281)	Glycosyltransferase
5-10	<i>wepI</i>	7956–9011 (351)	33.61	Glycosyltransferase group 1 (PF00534/8.9e-06)	α -D-GlcNAc α -1,2-L-rhamnosyltransferase <i>V. mimicus</i> VM223 (EEY46453)	64 (349)	Glycosyltransferase
5-11	<i>wepJ</i>	9067–10158 (363)	32.60		Hypothetical protein CLOBOL_01216, <i>C. boltea</i> ATCC BAA-613 (EDP18345)	26 (255)	Glycosyltransferases
5-12	<i>wepK</i>	10158–11072 (304)	29.61		Hypothetical protein glycosyltransferase Eps5K, <i>S. thermophilus</i>	32 (297)	Glycosyltransferases
5-13	<i>wzy</i>	11056–12153 (365)	31.23		Conserved hypothetical protein, <i>Bacteroides</i> sp. strain 1_1_6 (ZP_04850089.1)	24 (241)	Polymerase

^a aa, amino acid(s).

Genes responsible for GalNAc, Kdo, and GalA are not present in O-antigen gene clusters. Kdo is the obligatory component of the LPS core, and two genes, *ugd* and *gla*, responsible for the synthesis of GalA are involved in colonic acid synthesis. Therefore, it is reasonable that the genes responsible for the synthesis Kdo and GalA are located outside of the O-antigen gene cluster. In fact, these genes could be found in the genome sequences of *C. sakazakii* (NC_009778) and *C. turicensis* (NC_013282) (the locus tags are ESA-01502, ESA-02426, ESA-3591, ESA3590, ESA-02586, ESA-01534, and ESA-01535 in *C. sakazakii* and CTU-24190, CTU-15240, CTU-03800, CTU-03810, CTU-13590, CTU-23880, and CTU-23870 in *C. turicensis*). The gene for the synthesis of GalNAc is *gne*, which could be located inside or outside of the O-antigen gene cluster. In *C. sakazakii* O4, *gne* was found, while in *C. sakazakii* O1 and O6, *gne* is absent in their O-antigen gene clusters. By amplifying *gne* using the primer pair targeting the gene (5-ACAGATTGGTGTATGTTTCG and 5-ATCAAAGCAATA TCCACC), we found the *gne* could be located elsewhere in the chromosomes of *C. sakazakii* O1 and O6 (data not shown). Similar cases have been also been identified in *E. coli* (12, 13).

It has been reported in *E. coli* that the gene (*wecA*) responsible for the transfer of GalNAc-1-phosphate or GlcNAc-1-phosphate to the carrier lipid undecaprenol phosphate (UndP) for initiation of O unit synthesis is located outside the O-antigen gene cluster (1, 40). Based on the known O-antigen structures and gene clusters, five or six sugars are known to be present in *C. sakazakii* O antigens, and one fewer glycosyltransferase than sugars in O units has been noted. Therefore, a similar arrangement is proposed for *C. sakazakii*. For O5, no acetyltransferase gene (necessary for the O-acetyl modification of the O antigen) was found in its O-antigen gene cluster, indicating that this gene may be located on the prophage in the chromosome, similar to *Shigella* (19, 29, 38).

Three different pathways (the Wzx/Wzy, the ABC transporter, and the synthase pathways) have been described for the assembly and processing of O antigens (39). Most O antigens, particularly heteropolymeric O antigens, appear to be synthesized by the Wzx/Wzy pathway. The seven *C. sakazakii* O serotypes express a heteropolymeric O antigen and are synthesized by the Wzy/Wzx pathway (22, 34). The *wzy* gene is not present in its O-antigen gene cluster of *C. sakazakii* O7. However, the LPS of *C. sakazakii* O7 was

TABLE 5 Characteristics of ORFs in the *C. sakazakii* O6 O-antigen gene cluster^a

ORF	Gene	Gene position (length [aa])	G+C content (%)	Pfam conserved domain(s) (Pfam no./e-value)	Similar protein(s), strain(s) (GenBank accession no.)	% Identity (aa overlap)	Putative function of protein
6-1	<i>wepL</i>	47–829 (260)	31.30	Glycosyltransferase family 2 (PF00535/4.2e-25)	Glycosyltransferase, <i>F. bacterium</i> BAL38 (ZP_01733772.1)	38 (118)	Glycosyltransferase
6-2	<i>wepM</i>	813–2072 (413)	30.13	Capsule polysaccharide biosynthesis protein (PF05159/1.7e-22)	Capsule polysaccharide biosynthesis protein family protein, <i>Marinomonas</i> sp. (ZP_010075937.1)	34 (334)	Capsule polysaccharide biosynthesis protein family protein
6-3	<i>wzx</i>	2065–3270 (401)	32.18	Polysaccharide biosynthesis protein (PF01943/2.6e-26)	Polysaccharide biosynthesis protein, <i>B. helcogenes</i> P36-108 (YP_004162007)	27 (344)	Flippase
6-4	<i>wzy</i>	3260–4486 (408)	30.65		Wzy, <i>P. mirabilis</i> (ADL32282)	26 (195)	Polymerase
6-5	<i>wepN</i>	4488–5327 (279)	32.75	Core-2/I-branching enzyme (PF02485/2.1e-17)	Glycosyltransferase family 14, <i>M. paludis</i> DSM (ZP_07747376)	26 (294)	Glycosyltransferase
6-6	<i>wepO</i>	5324–6160 (278)	34.52	Glycosyltransferase family 2 (PF00535/2e-23)	Glycosyltransferase WceM, <i>Erwinia</i> sp. strain Ejp617 (ADP12985.1)	50 (267)	Glycosyltransferase
6-7	<i>wepP</i>	6185–7225 (346)	32.76	Acytransferase family (PF01757/1.2e-23)	Exopolysaccharide biosynthesis protein, <i>Erwinia</i> sp. strain Ejp617 (ADP10993)	32 (352)	Acetyltransferase

^a aa, amino acid(s).

extracted and separated by SDS–15% PAGE (data not shown). The results showed that *C. sakazakii* O7 produced normal LPS, which indicated that this strain has a functional *wzy* gene (36). Similar arrangements occur in *Salmonella* serogroups A, B, and D, which have a functional *wzy* gene outside the O-antigen gene cluster (50).

In the present study, O-antigen gene clusters of *C. sakazakii* O4 and *E. coli* O103 were observed to share an overall DNA identity of 60 to 74% (Fig. 1B), which is lower than the identities between housekeeping genes of bacteria (79 to 92%) (data not shown). The O antigen is exposed on the cell surface and undergo higher selec-

tion pressure for better adaptation to the situation; therefore, the genes in the O-antigen gene cluster were found to diverge at a higher rate than the housekeeping genes (42). If the overall levels of similarity in O-antigen gene clusters are higher than that for housekeeping genes, the O antigen may arise as a result of acquisition of the gene cluster by interspecies lateral transfer since species divergence (42). However, in the present study, the O-antigen gene clusters had a lower level of identity than did the housekeeping genes. Two possible explanations were proposed for this case. One is that the O-antigen gene cluster was present in the common ancestral species, and the O-antigen genes diverged at a high rate

TABLE 6 Characteristics of ORFs in the *C. sakazakii* O7 O-antigen gene cluster^a

ORF	Gene	Gene position (length [aa])	G+C content (%)	Pfam conserved domain(s) (Pfam no./e-value)	Similar protein(s), strain(s) (GenBank accession no.)	% Identical (aa overlap)	Putative function of protein
7-1	<i>wzx</i>	69–1358 (430)	33.02	Polysaccharide biosynthesis protein (PF01943/8.6e-23)	Polysaccharide biosynthesis protein Wzx, <i>V. vulnificus</i> (ABD38622)	31 (404)	Flippase
7-2	<i>wepQ</i>	1286–2741 (485)	29.42	Glycosyltransferase group 1 (PF00534/5.1e-12)	Glycosyltransferase group 1, <i>A. baumannii</i> ABNIH2 (EGT89242.1)	35 (264)	Glycosyltransferase
7-3	<i>wepR</i>	2743–3918 (391)	31.71	Glycosyltransferase group 1 (PF00534/4.3e-13)	Glycosyltransferase group 1, <i>Marinomonas</i> sp. (YP_001339674.1)	28 (392)	Glycosyltransferase
7-4	<i>fmlA</i>	3911–4945 (344)	41.06	Polysacc_synt_2 (PF02719.8/2.7e-109)	L-Fucosamine synthetase, <i>E. coli</i> (ABM53659)	93 (344)	Fucosamine synthetase
7-5	<i>fmlB</i>	4947–6050 (367)	41.30	NAD-dependent epimerase/dehydratase family (PF01370.14/1.2e-13)	FnlB, epimerase, <i>S. dysenteriae</i> (ACA24843)	78 (367)	Epimerase
7-6	<i>fmlC</i>	6050–7180 (376)	40.93	UDP-N-acetylglucosamine 2-epimerase (PF02350.12/5.5e-95)	C-2 epimerase, <i>E. coli</i> MS 145-7 (ZP_07692194.1)	92 (376)	Epimerase
7-7	<i>wbuB</i>	7180–8388 (402)	42.10	Glycos_transf_1 (PF00534.13/1.6e-09)	L-Fucosamine transferase, <i>S. dysenteriae</i> (ACA24845)	76 (398)	Glycosyltransferase
7-8	<i>wbuC</i>	8379–8783 (135)	42.71	Domain of unknown function (DUF1971/0.0009)	WbuC, <i>S. dysenteriae</i> (ACA24846)	69 (131)	Glycosyltransferase

^a aa, amino acid(s).




FIG 2 Specificity and sensitivity analysis of PCR assay for *C. sakazakii* O1 to O7. (a) Agarose gel electrophoresis of the PCR products obtained from respective *C. sakazakii* O1 to O7 strains by using the multiplex PCR assay. Lanes 1 and 10, DL2000 DNA markers; lane 2, O1; lane 3, O2; lane 4, O3; lane 5, O4; lane 6, O5; lane 7, O6; lane 8, O7; lane 9, negative control. (b) Agarose gel electrophoresis of the PCR products obtained from serially diluted chromosomal DNA of *C. sakazakii* O6 strain G2704. Lanes 1 and 10, DL2000 DNA markers; lane 2, negative control; lane 3, 0.00001 ng; lane 4, 0.0001 ng; lane 5, 0.001 ng; lane 6, 0.01 ng; lane 7, 0.1 ng; lane 8, 1 ng; lane 9, 10 ng. (c) Agarose gel electrophoresis of the PCR products obtained from serially diluted culture of *C. sakazakii* O6 strain G2704. Lanes 1 and 9, DL2000 DNA markers; lane 2, negative control; lane 3, 10^1 CFU/ml; lane 4, 10^2 CFU/ml; lane 5, 10^3 CFU/ml; lane 6, 10^4 CFU/ml; lane 7, 10^5 CFU/ml; lane 8, 10^6 CFU/ml.

than the housekeeping genes. The other is that the O-antigen gene cluster was acquired by one species from the other for a long time and was undergoing higher selection pressure than were the housekeeping genes. Most O antigens are distinct among different genus. Therefore, it is proposed that this O-antigen form is highly adaptive in *C. sakazakii* and *E. coli*, which occupy different niches, and may be important for the pathogenicity of these strains.

The O-antigen gene clusters of *C. sakazakii* O6 and *C. malonaticus* O2 were found to share high-level identity (99.3%), which may imply recent lateral transfer of the respective O-antigen gene cluster between these species. Similar cases have also been reported in *E. coli* strains (53). The O-antigen gene clusters

shared by *C. sakazakii* O3 and *C. muytjensii* O1, *C. turicensis* O1, and *C. malonaticus* O1 have also been recently reported (18). However, these pairs of O-antigen gene clusters exhibit a lower identity of 91%, in the range of or lower than the corresponding housekeeping gene identities (89 to 91% and 93 to 94%, respectively) (data not shown). These data indicate that the O-antigen gene clusters could not be evolved from recent lateral transfer. It is likely that these two O-antigen forms have the same case with that for *C. sakazakii* O4 and *E. coli* O103 discussed above.

Development of molecular serotyping methods is important for epidemiological surveillance of *C. sakazakii*, which is becoming an emerging food-borne pathogen infecting humans, especially infants. Some O-antigen genes, such as *wzx* and *wzy*, are highly specific to individual O serotypes, thus providing the potential for use in molecular assay for rapid identification and serotyping of relevant strains (51), which has been proved in the identification of *C. sakazakii* serotypes O1, O2, and O3 (22, 34). Many PCR-based typing methods targeting these O-antigen-specific genes have proved to be reliable, rapid, and sensitive for identifying isolates of *E. coli*, *Shigella*, and *Salmonella* from clinical, food, and environmental samples (7, 8, 14). In the present study, *C. sakazakii* O4 to O7 specific genes were identified, and O1 to O7 specific primers based on the targeted genes were designed, synthesized, and screened. The specific gene-based PCR serotyping method used here has been shown to be accurate and sensitive (22) and therefore provides a potential method that is suitable for serotyping *C. sakazakii* O1 to O7 strains.

ACKNOWLEDGMENTS

This study was supported by the National 973 Program of China grants 2012CB721001, 2012CB721101, and 2011CB504900; National Natural Science Foundation of China (NSFC) key program grant 31030002; NSFC General Program grants 31000044 and 81171524; National 863 Program of China grant 2011BAK10B02; Fundamental Research Funds for the Central Universities (65010721, 65020121, and 65020061); Research Fund for the Doctoral Program of Higher Education of China (grant 20090031120023); and National Key Programs for Infectious Diseases of China (2009ZX10004-108).

REFERENCES

- Alexander DC, Valvano MA. 1994. Role of the *rfe* gene in the biosynthesis of the *Escherichia coli* O7-specific lipopolysaccharide and other O-specific polysaccharides containing N-acetylglucosamine. *J. Bacteriol.* 176:7079–7084.
- Arbatsky NP, et al. 2011. Elucidation of the structure and characterization of the gene cluster of the O-antigen of *Cronobacter sakazakii* G2592, the reference strain of *C. sakazakii* O7 serotype. *Carbohydr. Res.* 346:1169–1172.
- Baldwin A, et al. 2009. Multilocus sequence typing of *Cronobacter sakazakii* and *Cronobacter malonaticus* reveals stable clonal structures with clinical significance which do not correlate with biotypes. *BMC Microbiol.* 9:223.
- Bengoechea JA, et al. 2002. Functional characterization of Gne (UDP-N-acetylglucosamine-4-epimerase), Wzz (chain length determinant), and Wzy (O-antigen polymerase) of *Yersinia enterocolitica* serotype O:8. *J. Bacteriol.* 184:4277–4287.
- Benson DA, et al. 2000. GenBank. *Nucleic Acids Res.* 28:15–18.
- Caubilla-Barron J, et al. 2007. Genotypic and phenotypic analysis of *Enterobacter sakazakii* strains from an outbreak resulting in fatalities in a neonatal intensive care unit in France. *J. Clin. Microbiol.* 45:3979–3985.
- DeRoy C, et al. 2004. Detection of *Escherichia coli* serogroups O26 and O113 by PCR amplification of the *wzx* and *wzy* genes. *Appl. Environ. Microbiol.* 70:1830–1832.
- Feng L, et al. 2004. Identification of *Escherichia coli* O114 O-antigen gene cluster and development of an O114 serogroup-specific PCR assay. *J. Clin. Microbiol.* 42:3799–3804.
- Finn RD, et al. 2006. Pfam: clans, web tools and services. *Nucleic Acids Res.* 34:D247–D251.

10. Fratamico PM, DebRoy C, Strobaugh TP, Jr, Chen CY. 2005. DNA sequence of the *Escherichia coli* O103 O antigen gene cluster and detection of enterohemorrhagic *E. coli* O103 by PCR amplification of the *wzx* and *wzy* genes. *Can. J. Microbiol.* 51:515–522.
11. Giovannini M, et al. 2008. *Enterobacter sakazakii*: an emerging problem in paediatric nutrition. *J. Int. Med. Res.* 36:394–399.
12. Giraud MF, et al. 1999. Purification, crystallization, and preliminary structural studies of dTDP-6-deoxy-D-xylo-4-hexulose 3,5-epimerase (RmlC), the third enzyme of the dTDP-L-rhamnose synthesis pathway, from *Salmonella enterica* serovar Typhimurium. *Acta Crystallogr. D Biol. Crystallogr.* 55:706–708.
13. Giraud MF, et al. 1999. Overexpression, purification, crystallization and preliminary structural study of dTDP-6-deoxy-L-lyxo-4-hexulose reductase (RmlD), the fourth enzyme of the dTDP-L-rhamnose synthesis pathway, from *Salmonella enterica* serovar Typhimurium. *Acta Crystallogr. D Biol. Crystallogr.* 55:2043–2046.
14. Han W, et al. 2007. DNA microarray-based identification of serogroups and virulence gene patterns of *Escherichia coli* isolates associated with porcine postweaning diarrhea and edema disease. *Appl. Environ. Microbiol.* 73:4082–4088.
15. Healy B, et al. 2010. *Cronobacter (Enterobacter sakazakii)*: an opportunistic foodborne pathogen. *Foodborne Pathog. Dis.* 7:339–350.
16. Healy B, et al. 2008. Evaluation of an automated repetitive sequence-based PCR system for subtyping *Enterobacter sakazakii*. *J. Food Prot.* 71:1372–1378.
17. Herikstad H, Motarjemi Y, Tauxe RV. 2002. *Salmonella* surveillance: a global survey of public health serotyping. *Epidemiol. Infect.* 129:1–8.
18. Hobbs M, Reeves PR. 1994. The JUMPstart sequence: a 39-bp element common to several polysaccharide gene clusters. *Mol. Microbiol.* 12:855–856.
19. Huan PT, Whittle BL, Bastin DA, Lindberg AA, Verma NK. 1997. *Shigella flexneri* type-specific antigen V: cloning, sequencing and characterization of the glucosyl transferase gene of temperate bacteriophage SfV. *Gene* 195:207–216.
20. Iversen C, et al. 2007. The taxonomy of *Enterobacter sakazakii*: proposal of a new genus *Cronobacter* gen. nov. and descriptions of *Cronobacter sakazakii* comb. nov., *Cronobacter sakazakii* subsp. *sakazakii*, comb. nov., *Cronobacter sakazakii* subsp. *malonaticus* subsp. nov., *Cronobacter turicensis* sp. nov., *Cronobacter muytjensii* sp. nov., *Cronobacter dublinensis* sp. nov., and *Cronobacter* genomospecies 1. *BMC Evol. Biol.* 7:64.
21. Iversen C, et al. 2008. *Cronobacter* gen. nov., a new genus to accommodate the biogroups of *Enterobacter sakazakii*, and proposal of *Cronobacter sakazakii* gen. nov., comb. nov., *Cronobacter malonaticus* sp. nov., *Cronobacter turicensis* sp. nov., *Cronobacter muytjensii* sp. nov., *Cronobacter dublinensis* sp. nov., *Cronobacter* genomospecies 1, and of three subspecies, *Cronobacter dublinensis* subsp. *dublinensis* subsp. nov., *Cronobacter dublinensis* subsp. *lausannensis* subsp. nov., and *Cronobacter dublinensis* subsp. *lactaridii* subsp. nov. *Int. J. Syst. Evol. Microbiol.* 58:1442–1447.
22. Jarvis KG, et al. 2011. Molecular characterization of *Cronobacter* lipopolysaccharide O-antigen gene clusters and development of serotype-specific PCR assays. *Appl. Environ. Microbiol.* 77:4017–4026.
23. Joseph S, Forsythe SJ. 2011. Predominance of *Cronobacter sakazakii* sequence type 4 in neonatal infections. *Emerg. Infect. Dis.* 17:1713–1715.
- 23a. Joseph S, et al. 2011. *Cronobacter condimenti* sp. nov., isolated from spiced meat and *Cronobacter universalis* sp. nov., a novel species designation for *Cronobacter* sp. genomospecies 1, recovered from a leg infection, water, and food ingredients. *Int. J. Syst. Evol. Microbiol.* 9:225.
24. Kneidinger B, et al. 2003. Three highly conserved proteins catalyze the conversion of UDP-N-acetyl-D-glucosamine to precursors for the biosynthesis of O antigen in *Pseudomonas aeruginosa* O11 and capsule in *Staphylococcus aureus* type 5: implications for the UDP-N-acetyl-L-fucosamine biosynthetic pathway. *J. Biol. Chem.* 278:3615–3627.
25. Krogh A, Larsson B, von Heijne G, Sonnhammer EL. 2001. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J. Mol. Biol.* 305:567–580.
26. Kucerova E, et al. 2010. Genome sequence of *Cronobacter sakazakii* BAA-894 and comparative genomic hybridization analysis with other *Cronobacter* species. *PLoS One* 5:e9556.
27. Kuhnert P, Korczak BM, Stephan R, Joosten H, Iversen C. 2009. Phylogeny and prediction of genetic similarity of *Cronobacter* and related taxa by multi-locus sequence analysis (MLSA). *Int. J. Food Microbiol.* 136:152–158.
28. Lai KK. 2001. *Enterobacter sakazakii* infections among neonates, infants, children, and adults. Case reports and a review of the literature. *Medicine (Baltimore)* 80:113–122.
29. Lindberg AA, Wollin R, Gernski P, Wohlhieter JA. 1978. Interaction between bacteriophage Sf6 and *Shigella flexneri*. *J. Virol.* 27:38–44.
30. Liu B, et al. 2008. Structure and genetics of *Shigella* O antigens. *FEMS Microbiol. Rev.* 32:627–653.
31. Liu D, Cole RA, Reeves PR. 1996. An O-antigen processing function for Wzx (RfbX): a promising candidate for O-unit flippase. *J. Bacteriol.* 178:2102–2107.
32. Marolda CL, Valvano MA. 1998. Promoter region of the *Escherichia coli* O7-specific lipopolysaccharide gene cluster: structural and functional characterization of an upstream untranslated mRNA sequence. *J. Bacteriol.* 180:3070–3079.
33. Miled-Bennour R, et al. 2010. Genotypic and phenotypic characterization of a collection of *Cronobacter (Enterobacter sakazakii)* isolates. *Int. J. Food Microbiol.* 139:116–125.
34. Mullane N, et al. 2008. Molecular analysis of the *Enterobacter sakazakii* O-antigen gene locus. *Appl. Environ. Microbiol.* 74:3783–3794.
35. Mullane NR, et al. 2007. *Enterobacter sakazakii* an emerging bacterial pathogen with implications for infant health. *Minerva Pediatr.* 59:137–148.
36. Nakhmchik A, Wilde C, Rowe-Magnus DA. 2007. Identification of a Wzy polymerase required for group IV capsular polysaccharide and lipopolysaccharide biosynthesis in *Vibrio vulnificus*. *Infect. Immun.* 75:5550–5558.
37. Pfohl A, et al. 2008. Biosynthesis of dTDP-3-acetamido-3,6-dideoxy- α -D-glucose. *Biochem. J.* 410:187–194.
38. Pupo GM, Lan R, Reeves PR. 2000. Multiple independent origins of *Shigella* clones of *Escherichia coli* and convergent evolution of many of their characteristics. *Proc. Natl. Acad. Sci. U. S. A.* 97:10567–10572.
39. Raetz CR, Whitfield C. 2002. Lipopolysaccharide endotoxins. *Annu. Rev. Biochem.* 71:635–700.
40. Reeves PP, Wang L. 2002. Genomic organization of LPS-specific loci. *Curr. Top. Microbiol. Immunol.* 264:109–135.
41. Rutherford K, et al. 2000. Artemis: sequence visualization and annotation. *Bioinformatics* 16:944–945.
42. Samuel G, Hogbin JP, Wang L, Reeves PR. 2004. Relationships of the *Escherichia coli* O157, O111, and O55 O-antigen gene clusters with those of *Salmonella enterica* and *Citrobacter freundii*, which express identical O antigens. *J. Bacteriol.* 186:6536–6543.
43. Samuel G, Reeves P. 2003. Biosynthesis of O-antigens: genes and pathways involved in nucleotide sugar precursor synthesis and O-antigen assembly. *Carbohydr. Res.* 338:2503–2519.
44. Shashkov AS, Arbatsky NP, Knirel YA. 2011. Structures and genetics of Kdo-containing O-antigens of *Cronobacter sakazakii* G2706 and G2704, the reference strains of serotypes O5 and O6. *Carbohydr. Res.* 346:1924–1929.
45. Staden R. 1996. The Staden sequence analysis package. *Mol. Biotechnol.* 5:233–241.
46. Sun Y, et al. 2011. Development of an O-antigen serotyping scheme for *Cronobacter sakazakii*. *Appl. Environ. Microbiol.* 77:2209–2214.
47. Reference deleted.
48. Tatusov RL, Koonin EV, Lipman DJ. 1997. A genomic perspective on protein families. *Science* 278:631–637.
49. Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22:4673–4680.
50. Wang L, Andrianopoulos K, Liu D, Popoff MY, Reeves PR. 2002. Extensive variation in the O-antigen gene cluster within one *Salmonella enterica* serogroup reveals an unexpected complex history. *J. Bacteriol.* 184:1669–1677.
51. Wang L, Reeves PR. 1998. Organization of *Escherichia coli* O157 O antigen gene cluster and identification of its specific genes. *Infect. Immun.* 66:3545–3551.
52. Wang L, Wang Q, Reeves PR. 2010. The variation of O antigens in gram-negative bacteria. *Subcell. Biochem.* 53:123–152.
53. Wang Q, et al. 2009. Genetic and structural analyses of *Escherichia coli* O107 and O117 O-antigens. *FEMS Immunol. Med. Microbiol.* 55:47–54.