

# Acid-Adapted Strains of *Escherichia coli* K-12 Obtained by Experimental Evolution

Mark M. Harden, Amanda He, Kaitlin Creamer, Michelle W. Clark, Issam Hamdallah, Keith A. Martinez II, Robert L. Kresslein, Sean P. Bush, Joan L. Slonczewski

Department of Biology, Kenyon College, Gambier, Ohio, USA

Enteric bacteria encounter a wide range of pHs throughout the human intestinal tract. We conducted experimental evolution of *Escherichia coli* K-12 to isolate clones with increased fitness during growth under acidic conditions (pH 4.5 to 4.8). Twenty-four independent populations of *E. coli* K-12 W3110 were evolved in LBK medium (10 g/liter tryptone, 5 g/liter yeast extract, 7.45 g/liter KCl) buffered with homopiperazine-*N,N'*-bis-2-(ethanesulfonic acid) and malate at pH 4.8. At generation 730, the pH was decreased to 4.6 with HCl. By 2,000 generations, all populations had achieved higher endpoint growth than the ancestor at pH 4.6 but not at pH 7.0. All evolving populations showed a progressive loss of activity of lysine decarboxylase (CadA), a major acid stress enzyme. This finding suggests a surprising association between acid adaptation and moderation of an acid stress response. At generation 2,000, eight clones were isolated from four populations, and their genomes were sequenced. Each clone showed between three and eight missense mutations, including one in a subunit of the RNA polymerase holoenzyme (*rpoB*, *rpoC*, or *rpoD*). Missense mutations were found in *adiY*, the activator of the acid-inducible arginine decarboxylase (*adiA*), and in *gcvP* (glycine decarboxylase), a possible acid stress component. For tests of fitness relative to that of the ancestor, *lacZ::kan* was transduced into each strain. All acid-evolved clones showed a high fitness advantage at pH 4.6. With the cytoplasmic pH depressed by benzoate (at external pH 6.5), acid-evolved clones showed decreased fitness; thus, there was no adaptation to cytoplasmic pH depression. At pH 9.0, acid-evolved clones showed no fitness advantage. Thus, our acid-evolved clones showed a fitness increase specific to low external pH.

Enteric bacteria experience a wide range of pHs throughout the human intestinal tract: from pH 1.5 to 3.5 in the stomach, then a rise to pH 4 to 7 in the duodenum and pH 7 to 9 in the jejunum, and then a decline to pH 5 to 6 in the cecum (1, 2). *Escherichia coli* has evolved to grow over a relatively wide range of intestinal pHs from pH 4.5 to 9.0 (3–5) and to survive without growth for several hours at pH 2 (6–8; for reviews of acid stress and extreme-acid survival, see references 9 to 11). The ability to survive the low pH of the stomach—and enter rapid growth as the pH increases in the lower digestive tract—may contribute to the low infectious doses characteristic of enteric pathogens, such as *E. coli* O157:H7 (12).

The adaptations of *E. coli* to growth in mild acid (pH 5 to 6) have been studied previously (9–11). Growth in mild acid may be enhanced by components of metabolism and transport, such as the lysine and arginine decarboxylases (13, 14), and by cyclopropane fatty acid synthase (8, 15, 16). A transcriptomic study of growth at pH 5 versus pH 7 shows that acid upregulates proton pumps of the electron transport system and outer membrane proteins, the chemotaxis regulon for proton-driven motility, periplasmic acid chaperones *hdeA* and *hdeB*, the *mar* multidrug resistance regulon, oxidative stress genes, and numerous genes with unidentified products (17).

Nevertheless, we still know little about how *E. coli* grows below pH 5, at the lower limit of its pH range, just above the range of nongrowth survival (pH 1.5 to 4.0). The range of pHs occurring during the transition from nongrowth to growth conditions is experienced by bacteria as they pass through the stomach and enter the intestine. Understanding growth in and adaptation to this lower range of growth pH is particularly important, because the gut microbiome may include hundreds of competing species (18, 19) and the ability of bacteria to outgrow their neighbors at pH 4 to 5 could influence which strains colonize the gut. The same

pH range also occurs in many fruit juices vulnerable to bacterial contamination (20).

To investigate genes that impact growth below pH 5, we used the approach of experimental evolution (21, 22). In an evolution experiment, multiple replicates of a single microbial lineage are repeatedly diluted and cultured under specific regulated conditions (23, 24). If the specified environment deviates from the optimal conditions for peak fitness, then there exists strong selective pressure for mutant alleles that enhance fitness in the new environment. *E. coli* has been the subject of a number of evolution studies involving adaptation to high temperatures (25), high ethanol concentrations (26), growth in complex media (27), and resistance to antibiotics (28).

Experimental evolution offers several advantages for the study of bacterial stress responses (29, 30). First, the approach allows the researcher to tightly regulate the microbe's environment for thou-

Received 22 October 2014 Accepted 28 December 2014

Accepted manuscript posted online 2 January 2015

Citation Harden MM, He A, Creamer K, Clark MW, Hamdallah I, Martinez KA, II, Kresslein RL, Bush SP, Slonczewski JL. 2015. Acid-adapted strains of *Escherichia coli* K-12 obtained by experimental evolution. *Appl Environ Microbiol* 81:1932–1941. doi:10.1128/AEM.03494-14.

Editor: J. L. Schottel

Address correspondence to Joan L. Slonczewski, slonczewski@kenyon.edu. M.M.H., A.H., and K.C. contributed equivalent leadership to the project.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/AEM.03494-14>.

Copyright © 2015, American Society for Microbiology. All Rights Reserved. doi:10.1128/AEM.03494-14

TABLE 1 Strains isolated after 2,000 generations<sup>a</sup>

Strain	Population	Isolate no.
JLSE0079	H9	1
JLSE0080	H9	2
JLSE0083	B11	1
JLSE0084	B11	2
JLSE0091	F11	1
JLSE0092	F11	2
JLSE0137	F9	2
JLSE0138	F9	3

<sup>a</sup> The sequence data are available in the NCBI SRA under accession number SRP041420.

sands of generations, which allows multiple selective sweeps of potentially unpredicted mutations to spread through the population. The evolving populations can be frozen regularly and stored indefinitely, allowing the researcher to set up direct comparisons between the ancestral strain and the adapted strains. Whole-genome resequencing can then permit identification of mutations in genes relevant to acid stress. Such an analysis might reveal surprising mechanisms of the stress response that differ from those identified by traditional gene-knockout investigation. Finally, experimental evolution allows us to determine whether acid adaptation comes at the price of a fitness trade-off in other environments.

Previous experimental evolution studies of acid stress are limited. *E. coli* experimental evolution in moderate acid (pH 5.3) yields an acid-dependent fitness increase, but genome sequence analysis has not been used to identify the underlying genetic changes (31). In another study, populations that evolved for 1,000 generations with 20 rounds of exposure to pH 2.5 showed 20 new mutations, including mutations in *evgS* alleles that relieve log-phase repression of extreme-acid resistance (32). Here we report on the isolation and characterization of eight *E. coli* strains from 4 of 24 populations cultured at pH 4.8 (for 730 generations) and then at pH 4.6 (2,000 generations in total). We found putative novel contributors to the acid stress response, including a strong selection for altered RNA polymerase.

## MATERIALS AND METHODS

**Bacterial strains and media.** *Escherichia coli* K-12 W3110 was the ancestral strain of all acid-adapted populations (33). Additional strains derived from *E. coli* K-12 W3110 were isolated during the course of the experiment (Table 1). For competition assays assessing the relative fitness of ancestral and adapted strains, a *lacZ::kan* insertion allele was introduced into each strain by P1 phage transduction from Keio collection strain GJ1886 (Coli Genetic Stock Center, Yale University) (34). The *lacZ::kan* insertion strains were maintained with 50 µg/ml kanamycin.

All bacterial cultures were grown in media based on LBK medium (10 g/liter tryptone, 5 g/liter yeast extract, 7.45 g/liter KCl) (17). The culture media were buffered with either 100 mM homopiperazine-*N,N'*-bis-2-(ethanesulfonic acid) (HOMOPIPES; pK<sub>a</sub> = 4.55, 8.12), 100 mM 2-(*N*-morpholino)ethanesulfonic acid (MES; pK<sub>a</sub> = 5.96), or 100 mM 3-morpholinopropane-1-sulfonic acid (MOPS; pK<sub>a</sub> = 7.20), depending upon the desired pH range of 4.4 to 4.8, 5.5 to 6.5, or 7.0, respectively (17). The pK<sub>a</sub> values presented above are those found at 25°C. For media buffered to pH 4.5 to 4.8, the LBK medium was supplemented with 10 g/liter *D*-l-malic acid (pK<sub>a</sub> = 3.40, 5.11) (LBK<sub>mal</sub>). The pH of the medium was adjusted as necessary with either 5 M HCl or 5 M KOH. For all experiments, the pH of the medium was tested immediately before and after bacterial growth.

**Experimental evolution.** A protocol based on successful evolution protocols described in the literature (22, 25, 27) was devised for experimental evolution of *E. coli* growth at external pH values below 5. Twenty-four populations were established from the same parental freezer stock of strain W3110. Evolving populations (200 µl each) were maintained in 96-well plates in a microplate reader. In each plate, 24 evolving populations occupied three columns of eight wells, with the columns being separated by an empty column to minimize the possibility of cross contamination. The initial inoculation of each well was with a 1:100 dilution of an overnight culture of the ancestral strain W3110. Populations were evolved in LBK<sub>mal</sub> with 100 mM HOMOPIPES at pH 4.8 (with daily dilutions of 1:4,000) for approximately 730 generations. The number of generations was estimated using the assumption that a dilution of 1:4,000 allows 12 generations of binary fission per day. After generation 730, the pH of the dilution medium was adjusted to 4.6 and the populations evolved for a further 1,270 generations (daily dilution, 1:100). A dilution of 1:100 allows 100-fold growth daily, producing 6.64 generations. The well plate was maintained at 37°C in a SpectraMax Plus384 MicroPlate reader (Molecular Devices). For the initial 22 h of the incubation, the well plate was shaken for 3 s every 15 min, after which the optical density at 450 nm (OD<sub>450</sub>) of each population was recorded. After 22 h, the plate was maintained at 37°C without shaking for 1 to 4 h until the next daily dilution.

For the first 730 generations, at the end of each daily cycle the endpoint cultures were supplemented with 100 µl glycerol (50%, vol/vol) and stored at -80°C. In the event that the cultures needed to be restarted from the freezer, aliquots of the most recent frozen cultures were diluted 1:50 into sterile growth medium on the first day and were then diluted as normal on the following days. After 730 generations (61 daily dilutions) of continuous growth in moderate acid, the pH of the LBK<sub>mal</sub> growth medium was lowered to 4.6. The daily dilution was decreased to 1:100, to decrease the population bottleneck (23). The glycerol solution for storing cultures was buffered with 100 mM MES, and the pH was adjusted to 6.5, to decrease the stress of the storage condition. All other aspects of the protocol remained unchanged.

After a cumulative 2,000 generations of evolution (61 dilutions at 1:4,000, 190 dilutions at 1:100), four populations were selected at random and streaked out on LBK agar. From each population, two clones were selected for genome sequencing (Table 1).

**Endpoint growth measurement at pH 4.6.** As one measure of evolving fitness, the endpoint growth level (culture density at stationary phase) was measured for cultures derived from each of the 24 experimental populations. The reference populations were chosen after approximately 25, 730, and 2,000 generations. Reference populations were thawed, diluted 1:50 in LBK<sub>mal</sub> at pH 4.8 (100 mM HOMOPIPES), and placed into rows of wells (volume, 200 µl) in the same microplate. The cultures were then incubated in a well plate at 37°C in a SpectraMax Plus384 MicroPlate reader (Molecular Devices) for 22 h. Every 15 min for the duration of the growth period, the well plate was shaken for 3 s, after which the OD<sub>450</sub> of each population was recorded. After 22 h, each population was diluted 1:100 into pH 4.6 medium for growth. A total of three serial dilutions at pH 4.6 (1:100) and 22-h incubations were performed. After each growth period, the endpoint values of the OD<sub>450</sub> were recorded and averaged.

**Lysine decarboxylase assays.** Isolated colonies were tested for lysine decarboxylase and other catabolic enzyme activities using an Enteropluri-Test indicator battery (Liofilchem). For large-scale screening, lysine decarboxylase was assayed by spotting an isolated colony into 200 µl Moeller decarboxylase broth, modified from the procedure of the company BD (BBL, 2011). The medium contained 5 g/liter *L*-lysine, 1 g/liter glucose, 3 g/liter yeast extract, and 15 mg/liter bromocresol purple. Samples were incubated at 37°C in the wells of a microplate; the plate was sealed for anaerobiosis. After 24 h, the plate was inspected visually. A purple color (alkalization) indicated positive lysine decarboxylase activity, whereas a yellow color indicated low or absent activity. Visual inspection was confirmed by analysis of the absorbance readings recorded for

each well at 340 nm and 590 nm using the SpectraMax Plus384 microplate reader (Molecular Devices).

**Whole-genome sequencing.** The genomes of the ancestral and evolved strains were sequenced. For each strain, the total DNA was extracted using a DNeasy DNA extraction kit (Qiagen). The DNA purity was confirmed by measuring the ratios of the absorbance at 260 nm/280 nm and 260 nm/230 nm. The genomes were sequenced with an Illumina MiSeq sequencer at the Michigan State University Research Technology Support Facility. Sequence reads were mapped to the *E. coli* K-12 W3110 reference genome (GenBank accession number [NC\\_007779.1](https://ncbi.nlm.nih.gov/nuccore/NC_007779.1)) (35). The quality scores (Q) for the sequences ranged from 25 to 33, with the average Q value being 30.8; this value corresponds to an error rate of 1/1,000. Sequences with 30-fold coverage or greater were accepted for mutation calls. Mutations were predicted with the breseq computational pipeline (<http://barricklab.org/breseq/>) (22).

**Competition assays.** Evolved clones were assayed for changes in fitness relative to that of the ancestral strain (36). Competitors were differentiated in coculture on the basis of *lac* marker states. Each evolved strain (*lac*-positive [*lac*<sup>+</sup>] phenotype) was cocultured with the *lac*-negative construct of the ancestral wild type (W3110 *lacZ::kan*) at pH 4.6. To ensure that the *lac*-negative phenotype did not confer a fitness advantage or disadvantage in the relevant environment, a competition assay between the *lac*-negative ancestor (W3110 *lacZ::kan*) and the original ancestor (W3110) was carried out simultaneously with each experiment. The two strains were cultured separately overnight at 37°C for 24 h with aeration in LBK<sub>mal</sub> at pH 4.8 (100 mM HOMOPIPES). The cultures were rediluted 1:100 into pH 4.8 media and incubated for an additional 16 to 18 h with aeration.

To commence competition, 125 μl of an overnight culture from a *lac*-negative strain and 125 μl of an overnight culture from a *lac*<sup>+</sup> strain were combined and diluted (100-fold) in 25 ml of LBK<sub>mal</sub> at pH 4.6 (100 mM HOMOPIPES). Samples were immediately serially diluted in M63 minimal medium [0.4 g/liter KH<sub>2</sub>PO<sub>4</sub>, 0.4 g/liter K<sub>2</sub>HPO<sub>4</sub>, 2 g/liter (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 7.45 g/liter KCl] and plated on LBK agar supplemented with ChromoMax IPTG (isopropyl-β-D-thiogalactopyranoside)-X-Gal (5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside) solution (4 μl/ml; Fisher). Cultures were incubated at 37°C with aeration for 24 h, and then the samples were again diluted and plated. The colony counts of white (*lac*-negative) and blue (*lac*<sup>+</sup>) colonies were used to calculate the total number of doublings for each competitor (31). The fitness of the evolved clone was compared to that of the ancestor by finding the difference between the two clones' number of doublings. The relative fitness (*W*) was calculated as  $[\log_2 100(N_{t2}) - \log_2(N_{02})]/[\log_2 100(N_{t1}) - \log_2(N_{01})]$ , where *N*<sub>01</sub> is the colony count for the ancestor in the mixed culture before dilution, *N*<sub>t1</sub> is the colony count for the ancestor after 100-fold dilution and 24 h of growth, *N*<sub>02</sub> is the colony count for the evolved clone in the mixed culture before dilution, and *N*<sub>t2</sub> is the colony count for the evolved clone after 100-fold dilution and 24 h of growth (36). Our colony counts allowed calculation of *W* over values up to 3; values higher than 3 were cut off at 3, because the low colony counts of the outcompeted strain caused high levels of variation.

Competition assays were carried out with 4- to 6-fold replication. For each competition, we included an equal number of trials in which the ancestor contained *lacZ::kan* and an equal number of trials in which the acid-adapted strain contained *lacZ::kan*. Similar fitness competitions were performed using the growth media LBK medium with 100 mM PIPES [piperazine-*N,N'*-bis(2-ethanesulfonic acid)], pH 6.5, and 15 mM benzoic acid and LBK medium with 150 mM TAPS (3-[[1,3-dihydroxy-2-(hydroxymethyl)-2-propanyl]amino]-1-propanesulfonic acid), pH 9.0.

**Log-phase growth of isolates at pH 4.5.** Strains were cultured initially in LBK<sub>mal</sub> buffered to pH 4.8 (100 mM HOMOPIPES) at 37°C for 24 h with aeration. For initial overnight cultures, pH 4.8 was selected to allow physiological acclimation and obtain consistent culture densities for the adapted and ancestral clones. Overnight cultures were rediluted 1:100 in fresh overnight medium and grown for 16 to 18 h. Triplicate cultures of 15

ml LBK<sub>mal</sub> at pH 4.5 (100 mM HOMOPIPES) were inoculated via a 1:200 dilution of the overnight culture, and the initial OD<sub>600</sub> of each culture was recorded. The cultures were incubated with aeration in a water bath at 37°C. The OD<sub>600</sub> was measured after 1 h and then subsequently at approximately 20-min intervals until the cultures began to enter stationary phase. The growth rate of each culture was calculated with respect to the time period corresponding to log-phase growth (37).

**Cotransductant strains and competition assays at pH 4.6.** A *thiG761::kan* insertion (80 to 90% linked to *rpoC*) was introduced into an evolved clone containing an *rpoC* mutation (clone B11-1) by P1 phage transduction using strain JW5549-1 (Keio collection). Cotransduction with the linked *kan* marker replaced the mutated region of the gene with the ancestral sequence. The replacement was confirmed by PCR amplification and sequencing of the gene. The forward primer sequence used was 5'-CCTCGGAGATGATTTGCGTG-3', and the reverse primer sequence used was 5'-GAGCTGTTCAAACCGTTCATC-3' (844 bp between primers). The cotransductant strain was competed against the parental clone in LBK<sub>mal</sub> at pH 4.6 following the same protocol described above.


**Nucleotide sequence accession number.** Sequence data have been deposited in the NCBI Sequence Read Archive (SRA) under accession number [SRP041420](https://ncbi.nlm.nih.gov/sra/SRP041420).

## RESULTS

**Adaptation of 24 experimental *E. coli* populations at pH 4.6.** We established culture conditions for maintaining experimental *E. coli* lineages below pH 5. In LBK medium, *E. coli* catabolizes diverse metabolites to optimize growth at low pH (9). The added components, HOMOPIPES and D/L-malate (LBK<sub>mal</sub>), buffered the acidity below pH 5. Both D- and L-malate may be catabolized by reactions that consume acid (38, 39) and thus potentially supply the experimental lineages with a novel means of enhancing acid tolerance. Using this medium, LBK<sub>mal</sub>, we established that our laboratory strain of *E. coli* K-12 W3110 could grow consistently in a microplate with the medium adjusted to pH 4.8.

Twenty-four bacterial populations were evolved under a daily serial dilution regime in 96-well microtiter plates in LBK medium supplemented with HOMOPIPES and D/L-malate, buffered initially at pH 4.8, as described in Materials and Methods. After 730 generations of growth, trial experiments of three successive 1:100 dilutions revealed that cultures of all experimental populations could be maintained at pH 4.6. We therefore decreased the pH of the culture conditions to pH 4.6 to provide even greater selective pressure on the experimental populations. We also decreased the daily dilution factor to 1:100, in order to decrease the daily population bottleneck and thus sample a greater number of possible mutations.

Over the course of the experiment, we observed significant increases in the endpoint culture density of all evolving populations, consistent with adaptation to low pH (Fig. 1). We measured the endpoint culture density of experimental populations sampled from populations frozen at several different generation numbers (generations 25, 730, and 2,000). The 25-generation plate (the second 4,000-fold dilution plate after the ancestor) was chosen instead of the ancestral strain, in order to test cultures frozen in plates under equivalent conditions. By generation 730, the acid-evolved populations showed an increased ability to grow at pH 4.6, although their endpoint densities declined by the third daily dilution. Comparison of the results by the *t* test to determine statistically significant differences (*P* < 0.001) verified the increase in endpoint growth between generations 25 and 730 and between generations 730 and 2,000 (evolved at pH 4.6). By generation 2,000, all populations reached endpoint densities higher than



**FIG 1** Growth of evolved populations at pH 4.6. The endpoint growth of the 24 populations from generations 25, 730, and 2,000 was compared. Frozen populations were thawed and diluted 1:50 in LBK<sub>mal</sub>, pH 4.8, for initial recovery and then incubated in a microplate reader for 22 h. Culture was conducted for three subsequent days by daily dilution 100-fold at pH 4.6. On each day, cultures were incubated at 37°C in a microplate reader with shaking every 15 min. The final OD<sub>450</sub> was measured after 22 h. Error bars represent SEMs ( $n = 24$ ).

those at generations 25 and 730. Furthermore, the 2,000-generation populations grew consistently over three serial dilutions at pH 4.6; a *t* test showed no difference among the endpoint reads for the three successive dilutions ( $P = 0.54$ ). Similar experiments conducted at pH 7.0 and at pH 9.0 showed no fitness advantage or disadvantage (data not shown). Thus, the growth advantage of the evolved populations was specific to pH 4.6, and the evolved populations incurred no fitness disadvantage at high pH.

**Downregulation of lysine decarboxylase activity.** *E. coli* bacteria respond to acid stress through various shifts in metabolism, such as upregulation of the lysine and arginine decarboxylases (13, 40). We therefore screened isolates from our evolving populations for several metabolic enzyme activities, using the Entero-Pluri-Test. Several of our 2,000-generation clones tested negative for the activity of the degradative lysine decarboxylase (CadA). This was surprising, given that in the ancestor, the gene encoding CadA is one of the genes that is the most highly induced by acid (13, 41).

We investigated the loss of lysine decarboxylase activity by screening three generations of our 24 evolving populations for enzyme activity (Table 2). For each generation number (760, 1,400, 2,000), three isolates from each of the 24 populations were screened for lysine decarboxylase activity after 24 h growth in Moeller broth with lysine. In this assay, the ancestor W3110 con-

sistently tested positive (purple), and isolates with a *cadA* deletion consistently tested negative (yellow). Among the sampled populations, we observed a progressive decline in the prevalence of decarboxylase-positive clones. At generation 760, only one isolate (one of three isolates from population B7) tested negative. At generation 1,400, 26 isolates from 15 populations tested negative for lysine decarboxylase function, and by generation 2,000, all but one isolate (an isolate from population A11) tested negative. Further screening of the populations revealed a minority of decarboxylase-positive isolates remaining in several populations, such as population F9 (data not shown).

Upon further incubation at 37°C, all sampled clones subsequently tested positive after 2 or 3 days. This observation suggests that the enzyme was present but downregulated. Indeed, genome sequencing subsequently confirmed that the *cadA* gene and its known regulators showed no mutations in the downregulated strains (see “Genomic analysis of acid-evolved isolates” below). The basis of *cadA* downregulation remains to be elucidated, but the occurrence of *cadA* downregulation in all 24 populations suggests a beneficial fitness effect under our acid stress conditions.

**Genomic analysis of acid-evolved isolates.** To identify the genetic basis of the improved fitness of the evolved clones at low pH, we sequenced the genomes of eight clones from generation 2,000 (SRA accession number SRP041420). Two clones were isolated from each of four experimental populations, populations H9, B11, F11, and F9. Among the sequenced clones, two (clones F9-2 and F9-3 from population F9) displayed lysine decarboxylase activity comparable to that of the ancestor; the other six evolved clones tested negative.

For comparison with our acid-evolved clones, we resequenced the ancestral strain W3110 from our original reference freezer stock (D8-W), as well as a clone from a second freezer stock in our collection (D13-W) (see Table S1 in the supplemental material). Our two W3110 clones showed nearly identical sequences. However, several differences appeared between the genome of our lab strain and the sequenced *E. coli* K-12 W3110 reference genome (ASM1024v1). For example, our lab strain showed deletions of the *insD-insC* and *insH-alsK* regions. Also, a number of single nucleotide polymorphisms (SNPs) appeared in the *icd* coding sequence. The source of the unusual *icd* variability is unknown.

Mutations were identified in the sequenced genomes by aligning each isolate’s raw reads with the *E. coli* K-12 W3110 reference genome (35) using the breseq computational pipeline (<http://barricklab.org/breseq>) (24). Mutation calls were accepted for sites showing 30-fold or greater coverage. Each acid-adapted strain had between two and six nonsynonymous mutations. Among the evolved genomes, we identified 24 mutations in coding regions (Table 3), which included 20 missense mutations, 2 single-base insertions (*hepA*, *mppA*), 1 9-bp insertion (*yejM*), and

**TABLE 2** Lysine decarboxylase activity of clones from acid-evolved populations<sup>a</sup>

No. of generations	No. of isolates in the following population testing negative for lysine decarboxylase activity:																							
	A7	B7	C7	D7	E7	F7	G7	H7	A9	B9	C9	D9	E9	F9	G9	H9	A11	B11	C11	D11	E11	F11	G11	H11
760		1																						
1,400		1	2		3			2	1	1	2	2	3	1	2		3	1	1		1			
2,000	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	2	3	3	3	3	3	3	3

<sup>a</sup> For each generation, three isolates from each population were assayed. The number of isolates of three isolates tested that lost lysine decarboxylase activity (indicated by a yellow color in Moeller broth) after 24 h of culture in a microplate is indicated.

TABLE 3 Mutations identified in coding regions in the 2,000th generation clones<sup>a</sup>

Gene	Function	Mutation	Identification of a mutation in clone:							
			H9-1	H9-2	B11-1	B11-2	F11-1	F11-2	F9-2	F9-3
<i>adiY</i>	Activator of <i>adiA</i>	Missense (R119P)			X	X				
<i>flu</i>	AG43 transport (biofilm)	Missense (A162E)	X	X						
<i>ftsX</i>	Cell division (IM transporter)	Missense (A101V)			X	X				
<i>fucI</i>	L-Fucose isomerase	Missense (R363S)	X	X						
<i>gcvP</i>	Glycine decarboxylase	Missense (T854A)		X						
<i>hepA</i>	RNAP recycling factor	Insertion (1139, +G)			X	X				
<i>icd</i>	Isocitrate dehydrogenase	Missense (D398E)	X	X	X	X	X	X		
<i>menH</i>	Menaquinone biosynthesis	Missense (D79Y)			X	X				
<i>metE</i>	Homocysteine transmethylase	Missense (R337S)				X				
<i>mhpB</i>	Phenylpropionate degradation	Missense (D226Y)							X	
<i>mppA</i>	Murein tripeptide transporter	Insertion (970, +A)	X	X						
<i>msbA</i>	Lipid A transport IM to OM	Missense (G150C)					X	X		
<i>rlmE</i>	23S rRNA methyltransferase	Missense (P157L)						X		
<i>rpoB</i>	RNAP beta subunit	Missense (A679V)	X	X						
<i>rpoC</i>	RNAP beta prime subunit	Missense (I774S)							X	X
<i>rpoC</i>	RNAP beta prime subunit	Missense (V507L)			X	X				
<i>rpoD</i>	RNAP sigma 70	Missense (M273I)					X	X		
<i>spr</i>	Murein endopeptidase	Missense (Q73K)						X		
<i>wzzE</i>	ECA polysaccharide chain length	Missense (P314Q)							X	X
<i>yeiH</i>	IM protein	Missense (A245S)		X						
<i>yejM</i>	IM hydrolase	Insertion (position 531) of 9 bp (TTTATCGCC)	X	X					X	X
<i>yfdI</i>	IM protein	Missense (V7L)			X					
<i>yfjK</i>	Ionizing radiation resistance	Missense (T400I)								X
<i>yjiK</i>	IM protein	Silent (TCG → TCA)	X							

<sup>a</sup> The genomes of the experimental clones and the ancestral strain were sequenced and aligned with the sequence of the *E. coli* K-12 W3110 reference genome (35) using the breseq computational pipeline (<http://barricklab.org/breseq>). The mutations listed represent differences between the genome of the ancestor and the adapted clones, excluding intergenic mutations. An X signifies that the mutation is present in that clone. IM, inner membrane; OM, outer membrane; ECA, enterobacterial common antigen.

1 silent mutation (*yjiK*). The high ratio of nonsynonymous mutations to synonymous mutations across the evolved isolates suggests that these mutations were likely to be adaptive under the experimental conditions (42). An additional 12 mutations in intergenic regions were also identified (Table 4). Pairs of clones isolated from the same population typically displayed both shared and unshared mutations; the shared mutations presumably arose earlier in the ancestor common to that population. Only one mutation (*yejM* insertion) that was clearly absent from the ancestor occurred in two acid-evolved populations. Multiple mutations

were found in *icd*, but numerous SNPs in *icd* were found in the ancestral stock, suggesting a mutation frequency unrelated to the evolution experiment. The diversity of our evolved clones suggests that our remaining populations, of the original 24, contained further mutations of interest.

Interestingly, only one of the sequenced mutations occurred in a gene with a well-known role in acid stress tolerance, *adiY*, the activator of the acid-inducible arginine decarboxylase (AdiA). The *adiY* mutation appeared in the two isolates of population B11 (B11-1 and B11-2). The other mutations that we found are likely

TABLE 4 Intergenic mutations identified in the 2,000th generation acid-adapted clones<sup>a</sup>

Proximal gene(s)	Mutation (positions, nucleotide change)	Identification of a mutation in clone:							
		H9-1	H9-2	B11-1	B11-2	F11-1	F11-2	F9-2	F9-3
<i>matP</i> →/← <i>ompA</i>	Intergenic (+40/+36, C to A)	X							
<i>racR</i> ←/→ <i>ydaS</i>	Intergenic (-61/-63, C to A)						X		
<i>Y75_p4289</i> ←/→ <i>eaeH</i>	Intergenic (-545/-1035, C to A)		X						
<i>proL</i> →	Noncoding (57/77 nt, C to A)							X	X
<i>nuoA</i> ←/← <i>lrhA</i>	Intergenic (-87/+544, C to A)							X	X
<i>yfjL</i> ←/← <i>yfjM</i>	Intergenic (-258/+102) +8 bp (GCACTATG)						X		
<i>rhaA</i> →/→ <i>rhaD</i>	Intergenic (+308/-143, C to A)	X	X						
<i>rrsC</i> ←	Noncoding (1417/1542 nt, +A)					X	X		
<i>pabA</i> →/→ <i>argD</i>	Intergenic (+34/-52, A to G)			X	X				
<i>yjbI</i> →/→ <i>ubiC</i>	Intergenic (+84/-139, G to T)	X	X						
<i>dcuA</i> ←/← <i>aspA</i>	Intergenic (-202/+104, G to T)			X	X				
<i>acs</i> ←/→ <i>ntfA</i>	Intergenic (-228/-165, G to A)							X	X

<sup>a</sup> The mutations listed indicate intergenic differences between the adapted clones and the ancestral wild type. A positive distance from a proximal gene represents the downstream location of the mutation relative to the gene's translation stop site, whereas a negative distance represents the mutation's upstream location relative to the gene's translation start site. nt, nucleotide position number.

to yield previously unrecognized mechanisms of acid stress response.


Every evolved genome had one mutation in a subunit of the RNA polymerase (RNAP) holoenzyme (43). The specific RNAP component that was affected differed among the populations. Population H9 isolates shared a mutation in *rpoB* (beta subunit; A679V), whereas B11 and F9 isolates each showed mutations in *rpoC* (beta prime subunit; V507L and I774S, respectively.) The *rpoC* beta prime subunit I774S mutation maps near the beta/beta prime interface and confers resistance to the CBR703 series of RNAP inhibitors (44). The F9 isolates shared a mutation in *rpoD* (housekeeping sigma 70; M2731). In each population, the RNAP mutation was shared by both clones, a result that would be consistent with early fixation in the evolution experiment. Besides the RNAP subunit mutations, the B11 strains also had a frameshift mutation in the HepA (RapA) helicase, which is involved in recycling RNAP and in UV resistance (45, 46).

**Relative fitness of 2,000th generation isolates at pH 4.6.** Fitness competitions were performed on our eight sequenced clones, in order to measure their relative fitness gains under acidic conditions. We cocultured the evolved clones against the ancestor over a 24-h growth period in LBK<sub>mal</sub> buffered to pH 4.6. Competitor colonies were counted before and after growth by plating on X-Gal agar, on which the different *lac* marker states could be discerned. The relative fitness of each evolved isolate was measured by determining the number of generations that each strain accumulated during the 24-h period and then dividing the isolate's number of doublings by that of the ancestor. Thus, a value of *W* of greater than 1 indicates that the *lac*<sup>+</sup> isolate had a fitness advantage, whereas a value of *W* of less than 1 indicates that the *lac*-negative ancestor had a fitness advantage. For each isolate, equal numbers of competitions in which the *lac* phenotypes were reversed (*lac*<sup>+</sup> isolate versus *lac*-negative ancestor) were performed. In order to confirm the neutrality of the *lac*-negative marker, the *lac*-negative ancestor was also competed with the *lac*<sup>+</sup> ancestor. The mean relative fitness of W3110 versus W3110 *lacZ::kan* was not significantly different (one-sample *t* test) from 1.0 and was a mean of 1.045 ± 0.07 with 99% confidence.

All eight evolved clones outcompeted the ancestor and displayed relative fitness values of 1.5 or higher (Fig. 2A). The magnitude of the fitness advantage varied among the clones, with a somewhat lower fitness advantage being noted in the population F11 isolates, but all isolates outcompeted the ancestor. These fitness values are notably larger than those typically reported for bacterial evolution experiments, including those conducted at pH 5.3 (31).

**Fitness with cytoplasmic pH depressed by benzoate.** The observed improved fitness of the evolved clones at pH 4.6 might be due to increased tolerance for depressed cytoplasmic pH, which occurs at values of external pH below 5 (9). To test this hypothesis, we measured the fitness of four evolved clones (one from each of the pairs described above) relative to that of the ancestor in a medium buffered to 6.5 but containing a permeant acid (benzoic acid, benzoate) which depresses the cytoplasmic pH (Fig. 2B). All four evolved clones showed a loss of fitness under these conditions, indicating that their improved fitness at pH 4.6 is not due to increased tolerance of the lowered cytoplasmic pH.


**Fitness at pH 9.0.** Fitness improvement in one environment often comes at the cost of reduced fitness in another (23). We



**FIG 2** Fitness changes in acid-evolved clones. For each pair of strains, the relative fitness (*W*) was calculated as described in Materials and Methods. To control for any fitness effect of the transduced *lac*-negative allele, the original *lac*<sup>+</sup> ancestral wild type was cocultured with the *lac*-negative wild type (gray bars). (A) pH 4.6. Clones were serially diluted (1:200) in LBK<sub>mal</sub> buffered to pH 4.8 and incubated overnight twice prior to each experiment. Each isolate was then cocultured with a *lac*-negative variant of the ancestral wild type (or else the *lac*-negative isolate was cocultured with a *lac*<sup>+</sup> ancestor) for 24 h in LBK<sub>mal</sub> buffered to pH 4.6. Error bars represent SEMs ( $n = 6$  for evolved strains,  $n = 18$  for ancestral strain W3110). (B) Benzoate. Strains were serially diluted 1:200 and incubated overnight twice in LBK medium buffered to pH 6.5. For the competition, each isolate and ancestor were cocultured for 24 h in LBK medium with 15 mM benzoic acid buffered with 100 mM PIPES at pH 6.5 ( $n = 4$  for evolved strains,  $n = 8$  for W3110). (C) pH 9.0. Strains were serially diluted and incubated overnight twice in LBK medium buffered to pH 8.5. For the competition, each isolate and ancestor were cocultured for 24 h in LBK medium buffered with 150 mM TAPS at pH 9.0 ( $n = 4$  for evolved strains,  $n = 8$  for W3110). \*,  $P \leq 0.05$  (*t* test) in comparison with the results for W3110.

sought to determine if the improved fitness of the evolved clones at low pH carried the trade-off of reduced fitness at high pH. To test this possibility, we measured the fitness of four of the evolved clones relative to that of the ancestor in a medium buffered to pH 9 (Fig. 2C). The evolved clones showed insignificant to slightly negative fitness changes, in addition to displaying variation among replicates greater than that observed at low pH. This result shows that the improved fitness of our acid-evolved clones is specific to growth at low pH, with little to no correlated fitness change occurring at high pH.

**Early-log-phase growth of 2,000th generation isolates at pH 4.6.** The relative fitness of bacterial strains evolved in our microplates could be due to various factors, including the culture density reached by stationary phase (Fig. 1) as well as the rate of growth in log phase. To separate these factors, we measured the growth rates of our acid-evolved isolates under conditions conducive to optimal log-phase growth (rotary shaking of baffled flasks). We cultured four isolates (representing the four populations) in LBK<sub>mal</sub> buffered to pH 4.5 and compared their early-log-phase growth rate to that of their ancestor (Fig. 3). The four isolates showed a significant increase in the log-phase growth rate com-



**FIG 3** Early-log-phase growth rates of acid-evolved clones in LBK<sub>mal</sub> buffered to pH 4.5. Clones were twice serially diluted (100-fold) and cultured overnight in LBK<sub>mal</sub> buffered to pH 4.8. Cultures were then diluted 1:200 into LBK medium buffered to pH 4.5 and then grown until mid-log phase at 37°C under aerated conditions. Growth rates were calculated from OD<sub>600</sub> measurements taken every 20 min throughout early-log-phase growth. Error bars indicate SEMs ( $n = 3$ ). \*, significant differences ( $P < 0.05$ ) from the results for ancestor strain W3110, based on  $t$  test.

pared to that of the ancestor. When cultured at pH 7, however, the log-phase growth rate of each strain was comparable to that of the ancestral wild type (data not shown).

**Relative fitness of evolved strain B11-1 with an *rpoC* mutation.** The evolved clone from population B11 had a missense mutation (V507L) in the *rpoC* gene (Table 3). In order to investigate the effect of the *rpoC* mutation on fitness, we transduced the linked W3110 *thiG761::kan* allele into strain B11-1 to produce a cotransductant strain in which the mutated *rpoC* is reverted to the ancestral sequence. The sequence reversion was confirmed by PCR. The cotransductant strain was competed against its parental mutant strain in LBK<sub>mal</sub> pH 4.6. The mean fitness of B11-1 *lacZ::kan* (containing *rpoC* V507L) relative to that of B11-1 *thiG761::kan* (containing a reverted *rpoC*) was  $1.358 \pm 0.03$  (99% confidence,  $n = 3$ , one-sample  $t$  test,  $P = 0.008$ ). The decreased fitness of B11-1 *thiG761::kan* with reverted *rpoC* suggests that the *rpoC* mutation provides a competitive advantage for acid growth.

## DISCUSSION

In this study, we examined the adaptation of *E. coli* to growth below pH 5, a condition relevant to understanding both enteric bacterial passage into the intestine (1, 18) and fruit juice contamination (20). Following 2,000 generations of experimental evolution at the lower limit of *E. coli*'s tolerable pH range, we found that eight evolved clones isolated from 4 of our 24 populations displayed a substantially higher fitness than the ancestral clone under growth conditions of pH 4.6 (Fig. 2). Indeed, we observed a fitness improvement much greater than that which has been seen in previous long-term evolution experiments at pH 5.3 (31). Our experiment was performed in a complex medium, LBK<sub>mal</sub>, which broadened the range of possible metabolic responses and, hence, targets of selection to the acidic conditions (9, 27). Evolved populations showed both a significantly higher endpoint culture den-

sity (Fig. 1) and significantly higher early-log-phase growth rates (Fig. 3), suggesting that these conditions selected for improvements in both the early and late parts of the growth cycle.

**Mutations in the RNA polymerase holoenzyme of acid-adapted strains.** All acid-adapted evolved clones that we examined have a missense mutation in a component of the RNAP holoenzyme (43, 47, 48). Our fitness competition assays suggest that the altered *rpoC* allele of evolved clone B11-1 confers a competitive advantage for acidic growth. In other reports, RNAP mutations arise in some but not all laboratory evolution experiments; the mechanism by which they alter fitness is poorly understood (23, 30).

For our acid-evolved strains, we proposed three possible mechanisms by which the observed RNAP mutations might have enhanced fitness at low pH. (i) The RNAP mutations could increase the stability of the complex at low pH, when the cytoplasm is slightly acidified by proton leakage from the external medium. In this case, we would expect the acid-evolved strains to show increased fitness in the presence of benzoic acid, which acidifies the cytoplasm (49). However, our acid-evolved clones showed decreased fitness in benzoate (Fig. 2B). Thus, under our conditions, acid-stressed bacteria did not evolve resistance to permeant acids.

(ii) The RNAP mutations could destabilize the complex and thus moderate the overexpression of acid stress genes. Such overexpression at the edge of the permissive pH range could be deleterious for growth, for example, by catabolizing amino acids needed for protein synthesis. Destabilizing mutations of RNAP have been described, although the details of their mechanism remain poorly understood (50). Downregulation via RNAP destabilization could be consistent with our observed downregulation of lysine decarboxylase in all evolving populations (Table 2). The gene *cadA* is one of the most highly expressed acid stress genes in *E. coli* (13, 17, 51).

(iii) The RNAP mutations could have caused the rapid adjustment of multiple regulons in small ways that cumulatively increase fitness under the selected condition. For example, under an experimental evolution involving adaptation to minimal glycerol medium, Herring et al. found that early RNAP substitutions conferred the greatest fitness advantage of all mutations isolated (52). Moreover, many other SNPs confer a fitness advantage only when combined with RNAP. The authors hypothesized that the RNAP mutations fixed very early in the experiment due to fitness benefits stemming from their manifold effects on gene regulation. They further hypothesized that subsequent mutations might have functioned to compensate for the inevitable negative side effects of altered RNAP function (47, 53).

**Putative acid stress response genes.** In our analysis of acid-evolved genome sequences, we identified mutations in several genes known to be important for *E. coli*'s acid stress response. The most notable example of these was in *adiY* (mutated in the B11 population of isolates). The *adiY* gene encodes an AraC-like regulatory protein that can induce expression of *AdiA* (arginine decarboxylase) and *AdiC* (arginine- $\alpha$ -glutamate antiporter), the essential components of the arginine-dependent acid resistance system (6, 40). The effect of the *adiY* mutation could be mediated by either an increased or a decreased rate of *adiAC* transcription.

Two other strains, F9-2 and F9-3, had a mutation in a regulatory upstream of the *nuo* operon, which codes for NADH:ubiquinone oxidoreductase (respiratory complex I), a component of the

electron transport chain (54, 55). The respiratory complex I is a proton pump of the electron transport chain known to be upregulated at low external pH (17). If this mutation upstream of *nuoA* were to increase the rate of transcription of the *nuo* operon, it could potentially increase acid stress by increasing the bacterium's net proton efflux.

In one clone, H9-2, we identified a missense mutation in *gcvP*, which encodes glycine decarboxylase P, part of the glycine cleavage system (56). Transcription of *gcvP* is positively regulated by glycine (57) and is responsible for the reversible breakdown of glycine into carbon dioxide, ammonia, and a cofactor-bound methylene group (58). The glutamate, arginine, lysine, and ornithine decarboxylase reactions have been demonstrated to play a role in maintaining neutral cytoplasmic pH. It is possible that the catabolic glycine decarboxylase pathway similarly serves to buffer the cytoplasm via the release of ammonia during glycine decarboxylation. Indeed, Lu et al. recently demonstrated that the conversion of glutamine to glutamate, with the release of ammonia, can serve a protective role for *E. coli* during exposure at pH 2.5 (59).

The protein Ag43, encoded by *flu*, was mutated in clones isolated from population H9. Ag43 is an autotransporter protein involved in cellular aggregation and biofilm formation, and production of Ag43 in a planktonic culture is associated with bacterial autoaggregation (forming clumps) (60, 61). Expression of *flu* is subject to phase variation, in which individuals in the same clonal population vary qualitatively in their production of the protein (60, 61). The missense mutation (A162E) that we identified is located in the protein's passenger domain,  $\alpha^{43}$ , which protrudes from the outer membrane into the extracellular space (60). This suggests that the mutant strains might have an altered ability to adhere to other cells, perhaps by binding to each other more tightly within the culture. Aggregation could benefit acid-stressed cells by reducing the total area of the cell surface exposed to the acidic medium, thereby reducing the rate of proton influx from the environment across the membrane.

Another possible acid stress gene is *msbA*, which encodes an ATP-dependent inner membrane flippase responsible for lipid transport from the inner leaflet to the outer leaflet of the bacterium's inner membrane. Loss of function of MsbA causes phospholipids and lipopolysaccharides (LPSs) to collect at the cell's inner membrane, thereby compromising the cell's structural integrity (62). The two evolved clones from population F11 showed the presence of a G150C missense mutation in one of MsbA's transmembrane domains. It is possible that this mutation could alter membrane integrity, which could in turn impact acid sensitivity.

This study has established eight acid-adapted *E. coli* clones for future study, each of which shows significant, though varying, fitness improvements relative to their ancestor. Many exciting genetic and evolutionary questions about these strains remain to be addressed. In future studies, we intend to investigate the role of RNAP mutations in gene expression and relative fitness, as well as the putative contributions of the other potentially beneficial mutant alleles that we have identified.

#### ACKNOWLEDGMENTS

This work was supported by National Science Foundation grant MCB-1329815.

We are most grateful to Richard Lenski, Zachary Blount, and Irina

Artsimovitch for valuable discussions of our data. We also thank Jeffrey Barrick, Rohan Maddamsetti, Peter Lund, and Houra Merrikh for stimulating discussions. We thank Harper Kerkhoff and Veronica De Pascuale for expert technical assistance.

#### REFERENCES

- Evans D, Pye G, Bramley R, Clark A, Dyson T, Hardcastle J. 1988. Measurement of gastrointestinal pH profiles in normal ambulant human subjects. *Gut* 29:1035–1041. <http://dx.doi.org/10.1136/gut.29.8.1035>.
- Ibekwe VC, Fadda HM, McConnell EL, Khela MK, Evans DF, Basit AW. 2008. Interplay between intestinal pH, transit time and feed status on the in vivo performance of pH responsive ileo-colonic release systems. *Pharm Res* 25:1828–1835. <http://dx.doi.org/10.1007/s11095-008-9580-9>.
- Slonczewski JL, Rosen BP, Alger JR, Macnab RM. 1981. pH homeostasis in *Escherichia coli*: measurement by  $^{31}\text{P}$  nuclear magnetic resonance of methylphosphonate and phosphate. *Proc Natl Acad Sci U S A* 78:6271–6275. <http://dx.doi.org/10.1073/pnas.78.10.6271>.
- Wilks JC, Slonczewski JL. 2007. pH of the cytoplasm and periplasm of *Escherichia coli*: rapid measurement by green fluorescent protein fluorimetry. *J Bacteriol* 189:5601–5607. <http://dx.doi.org/10.1128/JB.00615-07>.
- Zilberstein D, Agmon V, Schuldiner S, Padan E. 1984. *Escherichia coli* intracellular pH, membrane potential, and cell growth. *J Bacteriol* 158:246–252.
- Foster JW. 2004. *Escherichia coli* acid resistance: tales of an amateur acidophile. *Nat Rev Microbiol* 2:898–907. <http://dx.doi.org/10.1038/nrmicro1021>.
- Lin JS, Lee IS, Frey J, Slonczewski JL, Foster JW. 1995. Comparative analysis of extreme acid survival in *Salmonella typhimurium*, *Shigella flexneri*, and *Escherichia coli*. *J Bacteriol* 177:4097–4104.
- Riggins DP, Narvaez MJ, Martinez KA, Harden MM, Slonczewski JL. 2013. *Escherichia coli* K-12 survives anaerobic exposure at pH 2 without RpoS, Gad, or hydrogenases, but shows sensitivity to autoclaved broth products. *PLoS One* 8:e56796. <http://dx.doi.org/10.1371/journal.pone.0056796>.
- Slonczewski JL, Fujisawa M, Dopson M, Krulwich TA. 2009. Cytoplasmic pH measurement and homeostasis in bacteria and archaea. *Adv Microbiol Physiol* 55:1–79. [http://dx.doi.org/10.1016/S0065-2911\(09\)05501-5](http://dx.doi.org/10.1016/S0065-2911(09)05501-5).
- Krulwich TA, Sachs G, Padan E. 2011. Molecular aspects of bacterial pH sensing and homeostasis. *Nat Rev Microbiol* 9:330–343. <http://dx.doi.org/10.1038/nrmicro2549>.
- Kanjee U, Houry WA. 2013. Mechanisms of acid resistance in *Escherichia coli*. *Annu Rev Microbiol* 67:65–81. <http://dx.doi.org/10.1146/annurev-micro-092412-155708>.
- Audia JP, Webb CC, Foster JW. 2001. Breaking through the acid barrier: an orchestrated response to proton stress by enteric bacteria. *Int J Med Microbiol* 291:97–106. <http://dx.doi.org/10.1078/1438-4221-00106>.
- Meng S-Y, Bennett G. 1992. Nucleotide sequence of the *Escherichia coli cad* operon: a system for neutralization of low extracellular pH. *J Bacteriol* 174:2659–2669.
- Neely MN, Olson ER. 1996. Kinetics of expression of the *Escherichia coli cad* operon as a function of pH and lysine. *J Bacteriol* 178:5522–5528.
- Chang YY, Cronan JE. 1999. Membrane cyclopropane fatty acid content is a major factor in acid resistance of *Escherichia coli*. *Mol Microbiol* 33:249–259. <http://dx.doi.org/10.1046/j.1365-2958.1999.01456.x>.
- Hayes ET, Wilks JC, Sanfilippo P, Yohannes E, Tate DP, Jones BD, Radmacher MD, BonDurant SS, Slonczewski JL. 2006. Oxygen limitation modulates pH regulation of catabolism and hydrogenases, multidrug transporters, and envelope composition in *Escherichia coli* K-12. *BMC Microbiol* 6:89. <http://dx.doi.org/10.1186/1471-2180-6-89>.
- Maurer LM, Yohannes E, Bondurant SS, Radmacher M, Slonczewski JL. 2005. pH regulates genes for flagellar motility, catabolism, and oxidative stress in *Escherichia coli* K-12. *J Bacteriol* 187:304–319. <http://dx.doi.org/10.1128/JB.187.1.304-319.2005>.
- Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, Nielsen T, Pons N, Levenez F, Yamada T. 2010. A human gut microbial gene catalogue established by metagenomic sequencing. *Nature* 464:59–65. <http://dx.doi.org/10.1038/nature08821>.
- Gevers D, Knight R, Petrosino JF, Huang K, McGuire AL, Birren BW, Nelson KE, White O, Methé BA, Huttenhower C. 2012. The Human Microbiome Project: a community resource for the healthy human microbiome. *PLoS Biol* 10:e1001377. <http://dx.doi.org/10.1371/journal.pbio.1001377>.



20. Li W, Pan J, Xie H, Yang Y, Zhou D, Zhu Z. 2012. Pasteurization of fruit juices of different pH values by combined high hydrostatic pressure and carbon dioxide. *J Food Prot* 75:1873–1877. <http://dx.doi.org/10.4315/0362-028X.JFP-12-127>.
21. Lee DH, Palsson BO. 2010. Adaptive evolution of *Escherichia coli* K-12 MG1655 during growth on a nonnative carbon source, L-1,2-propanediol. *Appl Environ Microbiol* 76:4158–4168. <http://dx.doi.org/10.1128/AEM.00373-10>.
22. Blount ZD, Barrick JE, Davidson CJ, Lenski RE. 2012. Genomic analysis of a key innovation in an experimental *Escherichia coli* population. *Nature* 489:513–518. <http://dx.doi.org/10.1038/nature11514>.
23. Kassen R. 2014. Experimental evolution and the nature of biodiversity. Roberts & Company, Greenwood Village, CO.
24. Dragosits M, Mattanovich D. 2013. Adaptive laboratory evolution—principles and applications for biotechnology. *Microb Cell Fact* 12:64. <http://dx.doi.org/10.1186/1475-2859-12-64>.
25. Riehle MM, Bennett AF, Long AD. 2001. Genetic architecture of thermal adaptation in *Escherichia coli*. *Proc Natl Acad Sci U S A* 98:525–530. <http://dx.doi.org/10.1073/pnas.98.2.525>.
26. Goodarzi H, Bennett BD, Amini S, Reaves ML, Hottes AK, Rabinowitz JD, Tavazoie S. 2010. Regulatory and metabolic rewiring during laboratory evolution of ethanol tolerance in *E. coli*. *Mol Syst Biol* 6:378. <http://dx.doi.org/10.1038/msb.2010.33>.
27. Puentes-Tellez PE, Hansen MA, Sorensen SJ, van Elsas JD. 2013. Adaptation and heterogeneity of *Escherichia coli* MC1000 growing in complex environments. *Appl Environ Microbiol* 79:1008–1017. <http://dx.doi.org/10.1128/AEM.02920-12>.
28. Laehnemann D, Peña-Miller R, Rosenstiel P, Beardmore R, Jansen G, Schulenburg H. 2014. Genomics of rapid adaptation to antibiotics: convergent evolution and scalable sequence amplification. *Genome Biol Evol* 6:1287–1301. <http://dx.doi.org/10.1093/gbe/evu106>.
29. Elena SF, Lenski RE. 2003. Evolution experiments with microorganisms: the dynamics and genetic bases of adaptation. *Nat Rev Genet* 4:457–469. <http://dx.doi.org/10.1038/nrg1088>.
30. Conrad TM, Lewis NE, Palsson BO. 2011. Microbial laboratory evolution in the era of genome-scale science. *Mol Syst Biol* 7:11. <http://dx.doi.org/10.1038/msb.2011.42>.
31. Hughes BS, Cullum AJ, Bennett AF. 2007. Evolutionary adaptation to environmental pH in experimental lineages of *Escherichia coli*. *Evolution* 61:1725–1734. <http://dx.doi.org/10.1111/j.1558-5646.2007.00139.x>.
32. Johnson MD, Bell J, Clarke K, Chandler R, Pathak P, Xia Y, Marshall RL, Weinstock GM, Loman NJ, Winn PJ, Lund PA. 2014. Characterization of mutations in the PAS domain of the EvgS sensor kinase selected by laboratory evolution for acid resistance in *Escherichia coli*. *Mol Microbiol* 93:911–927. <http://dx.doi.org/10.1111/mmi.12704>.
33. Smith MW, Neidhardt FC. 1983. Proteins induced by anaerobiosis in *Escherichia coli*. *J Bacteriol* 154:336–343.
34. Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL, Mori H. 2006. Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol Syst Biol* 2:2006.0008. <http://dx.doi.org/10.1038/msb4100050>.
35. Hayashi K, Morooka N, Yamamoto Y, Fujita K, Isono K, Choi S, Ohtsubo E, Baba T, Wanner BL, Mori H, Horiuchi T. 2006. Highly accurate genome sequences of *Escherichia coli* K-12 strains MG1655 and W3110. *Mol Syst Biol* 2:2006.0007. <http://dx.doi.org/10.1038/msb4100049>.
36. Lenski RE, Rose MR, Simpson SC, Tadler SC. 1991. Long-term experimental evolution in *Escherichia coli*. I. Adaptation and divergence during 2,000 generations. *Am Nat* 138:1315–1341.
37. Deininger KNW, Horikawa A, Kitko RD, Tatsumi R, Rosner JL, Wachi M, Slonczewski JL. 2011. A requirement of TolC and MDR efflux pumps for acid adaptation and GadAB induction in *Escherichia coli*. *PLoS One* 6:e18960. <http://dx.doi.org/10.1371/journal.pone.0018960>.
38. Lukas H, Reimann J, Kim OB, Grimpo J, Uden G. 2010. Regulation of aerobic and anaerobic D-malate metabolism of *Escherichia coli* by the LysR-type regulator DmlR (YeaT). *J Bacteriol* 192:2503–2511. <http://dx.doi.org/10.1128/JB.01665-09>.
39. Kim OB, Reimann J, Lukas H, Schumacher U, Grimpo J, Duennwald P, Uden G. 2009. Regulation of tartrate metabolism by TtdR and relation to the DcuS-DcuR-regulated C<sub>4</sub>-dicarboxylate metabolism of *Escherichia coli*. *Microbiology* 155:3632–3640. <http://dx.doi.org/10.1099/mic.0.031401-0>.
40. Stim-Herndon KP, Flores TM, Bennett GN. 1996. Molecular characterization of *adiY*, a regulatory gene which affects expression of the biodegradative acid-induced arginine decarboxylase gene (*adiA*) of *Escherichia coli*. *Microbiology* 142:1311–1320. <http://dx.doi.org/10.1099/13500872-142-5-1311>.
41. Watson N, Donyak D, Rosey E, Slonczewski J, Olson E. 1992. Identification of elements involved in transcriptional regulation of the *Escherichia coli* *cad* operon by external pH. *J Bacteriol* 174:530–540.
42. Yang ZH, Bielawski JP. 2000. Statistical methods for detecting molecular adaptation. *Trends Ecol Evol* 15:496–503. [http://dx.doi.org/10.1016/S0169-5347\(00\)01994-7](http://dx.doi.org/10.1016/S0169-5347(00)01994-7).
43. Opalka N, Brown J, Lane WJ, Twist KA, Landick R, Asturias FJ, Darst SA. 2010. Complete structural model of *Escherichia coli* RNA polymerase from a hybrid approach. *PLoS Biol* 8:e1000483. <http://dx.doi.org/10.1371/journal.pbio.1000483>.
44. Artsimovitch I, Chu C, Lynch AS, Landick R. 2003. A new class of bacterial RNA polymerase inhibitor affects nucleotide addition. *Science* 302:650–654. <http://dx.doi.org/10.1126/science.1087526>.
45. Muzzini O, Campbell EA, Xia LL, Severinova E, Darst SA, Severinov K. 1998. Disruption of *Escherichia coli* HepA, an RNA polymerase-associated protein, causes UV sensitivity. *J Biol Chem* 273:15157–15161. <http://dx.doi.org/10.1074/jbc.273.24.15157>.
46. Jin DJ, Zhou YN, Shaw G, Ji XH. 2011. Structure and function of RapA: a bacterial Swi2/Snf2 protein required for RNA polymerase recycling in transcription. *Biochim Biophys Acta* 1809:470–475. <http://dx.doi.org/10.1016/j.bbarm.2011.03.003>.
47. Vassilyev DG, Sekine S, Laptenko O, Lee J, Vassilyeva MN, Borukhov S, Yokoyama S. 2002. Crystal structure of a bacterial RNA polymerase holoenzyme at 2.6 Å resolution. *Nature* 417:712–719. <http://dx.doi.org/10.1038/nature752>.
48. Malhotra A, Severinova E, Darst SA. 1996. Crystal structure of a  $\sigma^{70}$  subunit fragment from *Escherichia coli* RNA polymerase. *Cell* 87:127–136. [http://dx.doi.org/10.1016/S0092-8674\(00\)81329-X](http://dx.doi.org/10.1016/S0092-8674(00)81329-X).
49. Kannan G, Wilks JC, Fitzgerald DM, Jones BD, BonDurant SS, Slonczewski JL. 2008. Rapid acid treatment of *Escherichia coli*: transcriptomic response and recovery. *BMC Microbiol* 8:37. <http://dx.doi.org/10.1186/1471-2180-8-37>.
50. Baharoglu Z, Lestini R, Duigou S, Michel B. 2010. RNA polymerase mutations that facilitate replication progression in the *rep uvrD recF* mutant lacking two accessory replicative helicases. *Mol Microbiol* 77:324–336. <http://dx.doi.org/10.1111/j.1365-2958.2010.07208.x>.
51. Auger EA, Bennett GN. 1989. Regulation of lysine decarboxylase activity in *Escherichia coli* K-12. *Arch Microbiol* 151:466–468. <http://dx.doi.org/10.1007/BF00416608>.
52. Herring CD, Raghunathan A, Honisch C, Patel T, Applebee MK, Joyce AR, Albert TJ, Blattner FR, van den Boom D, Cantor CR, Palsson BO. 2006. Comparative genome sequencing of *Escherichia coli* allows observation of bacterial evolution on a laboratory timescale. *Nat Genet* 38:1406–1412. <http://dx.doi.org/10.1038/ng1906>.
53. Hindre T, Knibbe C, Beslon G, Schneider D. 2012. New insights into bacterial adaptation through *in vivo* and *in silico* experimental evolution. *Nat Rev Microbiol* 10:352–365. <http://dx.doi.org/10.1038/nrmicro2750>.
54. Weidner U, Geier S, Ptock A, Friedrich T, Leif H, Weiss H. 1993. The gene locus of the proton-translocating NADH ubiquinone oxidoreductase in *Escherichia coli*. Organization of the 14 genes and relationship between the derived proteins and subunits of mitochondrial complex I. *J Mol Biol* 233:109–122. <http://dx.doi.org/10.1006/jmbi.1993.1488>.
55. Bongaerts J, Zoske S, Weidner U, Uden G. 1995. Transcriptional regulation of the proton translocating NADH dehydrogenase genes (*nuoA–N*) of *Escherichia coli* by electron acceptors, electron donors and gene regulators. *Mol Microbiol* 16:521–534. <http://dx.doi.org/10.1111/j.1365-2958.1995.tb02416.x>.
56. Okamura-Ikeda K, Ohmura Y, Fujiwara K, Motokawa Y. 1993. Cloning and nucleotide sequence of the *gcv* operon encoding the *Escherichia coli* glycine cleavage system. *Eur J Biochem* 216:539–548. <http://dx.doi.org/10.1111/j.1432-1033.1993.tb18172.x>.
57. Stauffer LT, Fogarty SJ, Stauffer GV. 1994. Characterization of the *Escherichia coli* *gcv* operon. *Gene* 142:17–22. [http://dx.doi.org/10.1016/0378-1119\(94\)90349-2](http://dx.doi.org/10.1016/0378-1119(94)90349-2).
58. Kikuchi G, Motokawa Y, Yoshida T, Hiraga K. 2008. Glycine cleavage system: reaction mechanism, physiological significance, and hyperglycinemia. *Proc Jpn Acad Ser B Phys Biol Sci* 84:246–263. <http://dx.doi.org/10.2183/pjab.84.246>.

59. Lu PL, Ma D, Chen YL, Guo YY, Chen GQ, Deng HT, Shi YG. 2013. L-Glutamine provides acid resistance for *Escherichia coli* through enzymatic release of ammonia. *Cell Res* 23:635–644. <http://dx.doi.org/10.1038/cr.2013.13>.
60. van der Woude MW, Henderson IR. 2008. Regulation and function of Ag43 (Flu). *Annu Rev Microbiol* 62:153–169. <http://dx.doi.org/10.1146/annurev.micro.62.081307.162938>.
61. Henderson IR, Meehan M, Owen P. 1997. Antigen 43, a phase-variable bipartite outer membrane protein, determines colony morphology and autoaggregation in *Escherichia coli* K-12. *FEMS Microbiol Lett* 149:115–120. <http://dx.doi.org/10.1111/j.1574-6968.1997.tb10317.x>.
62. Doerrler WT, Raetz CRH. 2002. ATPase activity of the MsbA lipid flipase of *Escherichia coli*. *J Biol Chem* 277:36697–36705. <http://dx.doi.org/10.1074/jbc.M205857200>.