



Fate and Persistence of a Pathogenic NDM-1-Positive *Escherichia coli* Strain in Anaerobic and Aerobic Sludge Microcosms

David Mantilla-Calderon,  Pei-Ying Hong

King Abdullah University of Science and Technology (KAUST), Water Desalination and Reuse Center (WDRC), Biological and Environmental Science and Engineering Division (BESE), Thuwal, Saudi Arabia

ABSTRACT The presence of emerging biological pollutants in treated wastewater effluents has gained attention due to increased interest in water reuse. To evaluate the effectiveness of the removal of such contaminants by the conventional wastewater treatment process, the fate and decay kinetics of NDM-1-positive *Escherichia coli* strain PI7 and its plasmid-encoded antibiotic resistance genes (ARGs) were assessed in microcosms of anaerobic and aerobic sludge. Results showed that *E. coli* PI7 decayed at a significantly lower rate under anaerobic conditions. Approximate half-lives were 32.4 ± 1.4 h and 5.9 ± 0.9 h in the anaerobic and aerobic microcosms, respectively. In the aerobic microcosms, after 72 h of operation, *E. coli* PI7 remained detectable, but no further decay was observed. Instead, 1 in every 10,000 *E. coli* cells was identified to be recalcitrant to decay and persist indefinitely in the sludge. ARGs associated with the *E. coli* PI7 strain were detected to have transferred to other native microorganisms in the sludge or were released to the liquid fraction upon host decay. Extracellular DNA quickly degraded in the liquid fraction of the aerobic sludge. In contrast, no DNA decay was detected in the anaerobic sludge water matrix throughout the 24-h sampling period. This study suggests an increased likelihood of environmental dispersion of ARGs associated with anaerobically treated wastewater effluents and highlights the potential importance of persister cells in the dissemination of *E. coli* in the environment during reuse events of treated wastewater.

IMPORTANCE This study examines the decay kinetics of a pathogenic and antibiotic resistant strain of *Escherichia coli* in microcosms simulating biological treatment units of aerobic and anaerobic sludge. The results of this study point at a significantly prolonged persistence of the *E. coli* and the associated antibiotic resistance gene in the anaerobic sludge. However, horizontal transfer of the plasmid encoding the antibiotic resistance gene was detected in the aerobic sludge by a cultivation method. A subpopulation of persister *E. coli* cells was also detected in the aerobic sludge. The findings of this study suggest potential areas of concern arising from pathogenic and antibiotic-resistant *E. coli* during both anaerobic and aerobic sludge treatment processes.

KEYWORDS persister cells, bacterial decay, extracellular DNA decay, antibiotic resistance genes, horizontal gene transfer, wastewater treatment

In most wastewater treatment plants (WWTPs), aerobic activated sludge processes are conventionally used as the main biological unit to achieve removal of organic materials from wastewater but such systems incur a large energy cost (1). Anaerobic digesters are increasingly being considered for use as an alternative process for wastewater treatment due to their various advantages (2). Energy can be recovered from the wastewater in the form of methane produced from anaerobic fermentation of

Received 16 March 2017 Accepted 8 April 2017

Accepted manuscript posted online 14 April 2017

Citation Mantilla-Calderon D, Hong P-Y. 2017. Fate and persistence of a pathogenic NDM-1-positive *Escherichia coli* strain in anaerobic and aerobic sludge microcosms. *Appl Environ Microbiol* 83:e00640-17. <https://doi.org/10.1128/AEM.00640-17>.

Editor Ning-Yi Zhou, Shanghai Jiao Tong University

Copyright © 2017 Mantilla-Calderon and Hong. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Pei-Ying Hong, peiyong.hong@kaust.edu.sa.

the organic carbon, and the amount of sludge produced from anaerobic processes is lower than that from aerobic processes (1).

However, municipal wastewater often contains subtherapeutic levels of antibiotic residues, and the continuous exposure to antibiotics can select for antibiotic-resistant bacteria (ARB) within both aerobic and anaerobic sludge processes. In recent years, WWTPs had been shown to be potential hot spots for ARB and antibiotic resistance gene (ARG) propagation (3). Despite having undergone treatment, the treated municipal wastewater can still contain significant amounts of ARB and ARGs. For example, an earlier study showed that treated effluents that ultimately would be intended for reuse or discharged into receiving water bodies carried about 10^6 to 10^{11} heterotrophic CFU per cubic meter (4, 5), from which 16 to 28% corresponded to ARB (5). In another study, 10^7 to 10^9 copies of diverse tetracycline resistance genes were found in each cubic meter of chlorinated effluents (4). To further compound this problem, genes that confer resistance to carbapenems (i.e., bla_{NDM-1}), which are antibiotics used as a last line of defense against multidrug-resistant infections (6, 7), were also detected at alarming levels in final effluents, approaching 10^9 copies per cubic meter of treated wastewater (8).

This problem is of particular concern in water-scarce countries with pressing needs to reuse the treated wastewater. The incidence of bacterial pathogens carrying ARGs that confer resistance to antibiotics of last resort (e.g., bla_{NDM-1} -positive pathogenic *Escherichia coli*) requires particular attention, as the reuse of such treated wastewater effluents might pose a potential risk to the public health if disseminated into the environment during reuse events (4). As a first step to assess the risk associated with wastewater reuse, it is necessary to understand the differential fates and persistence of ARB and ARGs in the main biological treatment unit of anaerobic and aerobic wastewater treatment systems. Few studies comparing differential aerobic/anaerobic ARG removal have been performed and showed conflicting information. Molecular studies by Diehl and LaPara (9) and Burch et al. (10) found that ARGs tend to be removed more efficiently under anaerobic conditions. However, other studies suggest that combined anaerobic-aerobic (11) or, in contrast, completely aerobic conditions are more efficient at removing ARGs and ARB from wastewater (12).

Although informative, these earlier studies relied purely on molecular-based detection and did not examine the factors potentially shaping the decay or persistence of the ARG and the ARB host in the sludge. The formation of specialized persister cells is a strategy adopted by different types of bacteria, such as *E. coli*, to endure harsh environmental conditions (13). These cells are dormant variations of vegetative cells that occur at low frequencies within the bacterial population and are commonly known for their capacity to withstand supra-lethal concentrations of antibiotics (13–15). The contribution of persisters to the establishment of chronic infections due to several pathogens is well documented (16–18). However, little is known about the contribution of persisters to ARB survival in aerobic and anaerobic sludge. The earlier studies also did not examine the factors shaping the decay or persistence of the ARGs associated with the antibiotic-resistant host in anaerobic and aerobic sludge. These factors include the stability and persistence of ARGs as extracellular DNA and the potential for horizontal gene transfer (HGT) events when the ARG is encoded on a conjugative plasmid.

In this study, *Escherichia coli* strain PI7 was used as a model bacterium to examine the existing knowledge gaps associated with the fate and persistence of ARB and ARGs in anaerobic and aerobic sludge. This bacterium was previously isolated from wastewater and carries an extensive repertoire of ARGs, including a copy of bla_{NDM-1} , in a plasmid of the IncF family identified earlier as pKOX_NDM1 (19). In the last decade, bla_{NDM-1} has undergone a pandemic spread among clinically relevant bacteria (20, 21). It has also been detected at alarming levels in different environmental compartments, including water (22, 23), soil (24), and wastewater (8, 19). Given the importance of wastewater in the environmental mobilization of ARB and ARGs, we aimed to evaluate the fate and persistence of our model bacterium *E. coli* PI7 and its associated IncF plasmid in anaerobic and aerobic sludge microcosms under various trace antibiotic




FIG 1 bla_{NDM-1} decay kinetics in nontreated (black circles) and PMA-treated (white circles) biomass fractions under anaerobic (a) and aerobic (b) conditions at $0 \mu\text{g/liter}$ of meropenem ($n = 3$). Plasmid stability under anaerobic (c) and aerobic (d) conditions was evaluated by comparing the bla_{NDM-1} (white circles) and $uidA$ (white triangles) decay rates in PMA-treated biomass fractions ($n = 3$). Arrows in panel d indicate sampling points at which transconjugants were recovered by plating techniques.

concentrations. Using molecular tools, this study evaluated plasmid stability, host persistence, and HGT events under anaerobic and aerobic conditions. Chromosomal and plasmidic decay rates were measured by quantitative PCR (qPCR) coupled with propidium monoazide (PMA) to discriminate between dead cells or extracellular DNA and cells with intact cell membranes. In addition, the potential stability of extracellular DNA in the liquid fraction of anaerobic and aerobic sludge was also evaluated.

RESULTS

Differential decay of bla_{NDM-1} in anaerobic and aerobic sludge microcosms.

Under anaerobic conditions, bla_{NDM-1} decayed following a first-order decay kinetic model ($R^2 > 0.935$) (Fig. 1a). After 360 h of operation, bla_{NDM-1} copy numbers decreased by 2.5 log from 10^8 to 10^6 copies/g biomass. This decrease in the bla_{NDM-1} copy numbers corresponded to average first-order decay rates of $-0.021 \pm 0.002 \text{ h}^{-1}$ ($t_{1/2} = 32.9 \pm 2.8 \text{ h}$) and $-0.021 \pm 0.001 \text{ h}^{-1}$ ($t_{1/2} = 32.3 \pm 1.7 \text{ h}$) for the nontreated biomass (NTB) and PMA-treated biomass (PTB) fractions, respectively (Fig. 1a, Table 1).

Under aerobic conditions, bla_{NDM-1} decay in the biomass fraction showed a biphasic behavior (Fig. 1b). An initial 4-log decrease in copy numbers from 10^8 to 10^4 copies/g biomass during the first 72 h of reactor operation was observed. Phase I fits a first-order decay kinetic model ($R^2 > 0.83$), with average decay rates of $-0.1049 \pm 0.019 \text{ h}^{-1}$ ($t_{1/2} = 6.6 \pm 1.2 \text{ h}$) and $-0.1196 \pm 0.014 \text{ h}^{-1}$ ($t_{1/2} = 5.8 \pm 0.7 \text{ h}$) (Fig. 1b, Table 1) for the NTB and PTB fractions, respectively. Similar to results observed under anaerobic conditions, differences in the bla_{NDM-1} decay rates of the two fractions were not

TABLE 1 Decay rates of bla_{NDM-1} in PMA-treated biomass fractions of anaerobic and aerobic sludge microcosms under different trace meropenem concentrations

Microcosm	Replicate run	Meropenem concentration ($\mu\text{g/liter}$) ^a	k (h^{-1}) ^b	$t_{1/2}$ (h)	P value ^c
Anaerobic sludge	1	Control	-0.0202	34.3	
		100	-0.0223	31.1	0.33
	2	Control	-0.0219	31.7	
		100	-0.0213	32.5	0.15
	3	Control	-0.0223	31.1	NA
		100	-0.0205	33.8	0.98
Aerobic sludge	1	Control	-0.1361	5.1	
		1	-0.1460	4.7	0.77
		10	-0.1436	4.8	0.82
		100	-0.1386	5.0	0.94
	2	Control	-0.1107	6.3	
		1	-0.1095	6.3	0.70
		10	-0.0993	7.0	0.71
	3	100	-0.0923	7.5	0.46
		Control	-0.1119	6.2	
		1	-0.1253	5.5	0.49
		10	-0.1095	6.3	0.85
		100	-0.1133	6.1	0.96

^aControl corresponds to 0 $\mu\text{g/liter}$ of meropenem.

^b k values for the aerobic microcosms correspond to phase I decay.

^c P values correspond to the comparison of each decay curve derived from meropenem-spiked microcosms (i.e., 1, 10, and 100 $\mu\text{g/liter}$ meropenem), with the decay model obtained in the respective control microcosms (0 $\mu\text{g/liter}$). No significant differences were observed in their bla_{NDM-1} decay at the different meropenem concentrations tested. P value corresponds to the testing of $H_0: k_{\text{control}} = k_{xi}$ $\mu\text{g/liter}$.

statistically significant ($P = 0.92$). After 72 h of aerobic exposure, decay was then followed by a plateau in phase II, with NTB and PTB average decay rates of $-0.007 \pm 0.005 \text{ h}^{-1}$ ($R^2 = 0.917$) and $-0.005 \pm 0.012 \text{ h}^{-1}$ ($R^2 = 0.917$), respectively. No statistically supported differences were observed between NTB and PTB decay rates, suggesting that most of the detected bla_{NDM-1} is harbored within cells with intact cell membranes. PTB decay rates (associated with cells with intact membranes) were statistically undistinguishable from a decay of zero ($P = 0.26$). Instead, bla_{NDM-1} abundance stabilized at 10^5 copies/g biomass.

Decay of bla_{NDM-1} correlates with *E. coli* P17 chromosomal decay. Due to the plasmidic origin of bla_{NDM-1} , the decay kinetics of this ARG might be explained by (i) plasmid loss or (ii) cellular decay of the bacterial host. To determine which of these two factors mainly explained the bla_{NDM-1} decay observed in the microcosms, we compared the PTB decay rates of bla_{NDM-1} (indicative of plasmid decay rates) with the PTB decay rates of a chromosome-carried gene (*uidA*). In the anaerobic microcosms, the two target genes decayed at the same rate ($P = 0.32$) (Fig. 1c). Similarly, under aerobic conditions, phase I and plateau bla_{NDM-1} and *uidA* decay rates were statistically indistinguishable among all tested trace antibiotic concentrations ($0.19 > P > 0.50$) (Fig. 1d). *uidA* was initially spiked in the microcosm experiments at a frequency of 10^9 copies/g biomass and stabilized at 10^5 copies/g biomass during the plateau phase. These results suggest the presence of a recalcitrant subpopulation of *E. coli* cells, occurring at a frequency of 10^{-4} (plateau *uidA* copies/initial *uidA* spiking). The occurring frequency of recalcitrant *uidA* copies coincides with the frequency of persister cells of *E. coli* P17 in pure culture (Fig. 2).

Trace antibiotic concentrations do not influence the persistence of *E. coli* P17 or its associated ARGs. The results from the previous section suggest that bla_{NDM-1} decay is mainly explained by the cellular decay of the *E. coli* P17 host. As residual antibiotics are commonly found in wastewater (25), there is a latent possibility that such trace antibiotic concentrations might provide a selective advantage to *E. coli* P17 in the sludge, and consequently prolong the persistence of bla_{NDM-1} . To evaluate this possibility, the decay of bla_{NDM-1} and *uidA* was assessed in the anaerobic microcosms at 0




FIG 2 *E. coli* PI7 cell count before ($t = 0$ min) and after ($t = 180$ min) meropenem challenge at $640 \mu\text{g/ml}$ ($n = 4$). The persister cell frequency is expressed as the ratio of the cell count at $t = 180$ min and the cell count at $t = 0$ min, which corresponded to a frequency of $3.14 \times 10^{-4} \pm 1.5 \times 10^{-4}$ ($n = 4$).

and $100 \mu\text{g/liter}$ of meropenem and at 0, 1, 10, and $100 \mu\text{g/liter}$ of meropenem in the aerobic microcosms. None of the selected meropenem concentrations influenced $bla_{\text{NDM-1}}$ or *uidA* decay kinetics under either anaerobic or aerobic conditions (see Fig. S1 and S2 in the supplemental material). NTB and PTB decay rates (Table 1) at the different antibiotic concentrations were also not statistically different from their respective controls (i.e., reactors without antibiotic; $P > 0.15$).

Activated sludge as a reservoir for ARGs upon *E. coli* PI7 decay. Due to the plasmidic origin of $bla_{\text{NDM-1}}$, the sludge biomass might act as a sink for this gene through plasmid conjugation. Accordingly, HGT was assessed by culture-based methods. After 72 h of operation, transconjugants were recovered from the aerobic microcosm at three sampling points (Fig. 1d) at an average frequency of 10^3 to 10^4 CFU/g of biomass (see Text SI-1 in the supplemental material). Subsequent Sanger-based sequencing revealed that the transconjugants were $bla_{\text{NDM-1}}$ -positive isolates of *Enterobacteriaceae* belonging to *Shigella* and *Citrobacter* genera. No HGT events were detected by culture-dependent methods in the anaerobic microcosms.

Differential decay of colloidal DNA between anaerobic and aerobic liquid fractions. Upon *E. coli* PI7 cellular decay, $bla_{\text{NDM-1}}$ was detected in the supernatant fraction of the microcosm experiments by endpoint PCR (data not shown). To evaluate the potential persistence of extracellular DNA in liquid fractions of anaerobic and aerobic sludge fractions, a separate experiment was carried out. Naked plasmidic DNA was spiked in a dialysis cassette and its decay was tracked by qPCR and electroporation assays in TOP10 *E. coli* cells. As sequence-dependent DNA flexibility determines the cleave rate mediated by DNase I (26), the decay rate of the cloning vector does not necessarily represent the decay rate of the $bla_{\text{NDM-1}}$ plasmid. Specifically, given the larger size of the $bla_{\text{NDM-1}}$ plasmid (ca. 110 kbp), it may be more inclined to degrade faster than the smaller clone vector. Although the substitution of the clone vector may overestimate the persistence of the actual plasmid decay, this experiment is informative about the persistence potential of DNA/ARGs in anaerobic and aerobic liquid fractions. No decay was detected by qPCR or electroporation assays for plasmids spiked into the anaerobic liquid fraction ($P = 0.5$) (Fig. 3a and b). In contrast, in the aerobic liquid fraction, plasmid decay was detected by both qPCR and electroporation assays at rates of $-0.036 \pm 0.005 \text{ h}^{-1}$ ($t_{1/2} = 19.4 \pm 2.8 \text{ h}$) and $-0.278 \pm 0.03 \text{ h}^{-1}$ ($t_{1/2} = 2.5 \pm 0.25 \text{ h}$), respectively. Decay rates estimated by electroporation assays in the aerobic liquid fraction were significantly higher than those quantified by qPCR ($P \ll 0.05$).




FIG 3 Decay of extracellular colloidal DNA in anaerobic (white diamonds) and aerobic (black diamonds) sludge liquid fractions measured by qPCR (a) and electroporation (b) assays.

DISCUSSION

The removal of ARB from wastewater, particularly strains that are resistant to new classes of antibiotics, is required to protect public health during reuse events (4). Both aerobic- and anaerobic-based wastewater treatment systems are utilized for the treatment of municipal wastewater. However, a key parameter to consider when implementing these systems is the decay rates of ARB and ARGs in the associated sludge and in the treated wastewater effluents.

Using *bla*_{NDM-1}-positive *E. coli* P17 isolated from wastewater as a model bacterium, it was observed that the decay rate of this bacterium was 1 order of magnitude higher in the aerobic sludge than in the anaerobic sludge microcosms. The longer persistence of *E. coli* P17 observed under the anaerobic condition coincides with earlier microcosm studies comparing the differential anaerobic/aerobic survival of *E. coli* in diluted fermenter sludge (27), manure, manure slurry (28, 29), drinking water (30), and activated sludge (31). These studies provide evidence that oxygen is a key factor driving decay of *E. coli* in the secondary habitat, and partly explain the longer persistence of *E. coli* P17 in anaerobic microcosms. Endogenous reactive oxygen species (ROS) are formed as byproducts during aerobic metabolism (32). ROS are highly oxidative molecules that can react with cellular components and lead to cytotoxic (32–34) and mutagenic defects in the cells (35). Several detoxification mechanisms have evolved in aerobic and facultative organisms in order to cope with the deleterious effects of ROS (32, 36). Even though *E. coli* possesses several of those detoxification mechanisms (32), it is possible that the cells experiencing oxidative stress undergo a decrease in fitness and hence decayed faster in the aerobic microcosms.

However, the presence of oxygen is most likely not the only factor driving the decay of *E. coli* P17 in the microcosms, since decay was also observed in the anaerobic microcosms. In addition to ROS, the microbial community has been reported to play an important role in the decay of *E. coli* within the environment (37). The longer persistence of *E. coli* P17 could also be partially explained by the fact that microbial communities in the gut and the anaerobic sludge tend to be more similar to each other, as exemplified by the predominance of *Firmicutes* and *Bacteroidetes* in both ecosystems (38), than are the gut microbiota and aerobic sludge (39–41). Due to these similarities in the microbial communities, *E. coli* P17 might have acclimated better and decayed more slowly in the anaerobic sludge than aerobic sludge.

It was further observed that the decay kinetics of *E. coli* P17 followed a biphasic decay behavior in the aerobic microcosms. This biphasic decay behavior has been documented extensively for *E. coli* and other fecal indicator bacteria (42, 43). After the

initial spiking event, *bla*_{NDM-1} and *uidA* were detected in the aerobic reactors at an average frequency of 10⁹ copies/g biomass. After 72 h, no further decay was observed, and both gene copy numbers plateaued at an approximate frequency of 10⁵ copies/g biomass (Fig. 1b). As only one copy of *bla*_{NDM-1} or *uidA* is present per *E. coli* P17 cell, these results indicate that 1 in 10,000 cells may be recalcitrant to decay (see Text S1-2 in the supplemental material). Previous studies on pure cultures of *E. coli* suggest this pattern resulted from heterogeneity within the bacterial population (44). We further hypothesized that the bacterial subpopulation responsible for the plateau phase phenotype corresponds to persister cells, a phenotypical variant of vegetative cells that exhibit improved tolerance to antimicrobials and other stressful environmental conditions (13). Indeed, the same *E. coli* P17 was observed to upregulate transcriptional responses of genes related to persister cell formation upon solar irradiation (45), suggesting that *E. coli* P17 adopts this strategy to facilitate its persistence during stressful conditions. Persisters typically occur at a fixed frequency between 10⁻⁴ and 10⁻⁶, depending on the *E. coli* strain (14). Consequently, in all aerobic microcosms (*n* = 12), the *E. coli* P17 populations consistently stabilized at a frequency of 10⁵ cells/g biomass after a 4-log cell density decline (Fig. 1). A further evaluation of the frequency of persister cell formation in pure cultures of *E. coli* P17 revealed that for every 10,000 vegetative cells, 1 persister cell is formed (frequency of 10⁻⁴) (Fig. 2), coinciding with the persisting cell numbers obtained in the decay experiments. Further evidence supporting the presence of persister cells in the aerobic sludge is the fact that even though *uidA* was consistently detected in PTB fractions at a frequency of 10⁵ copies/g sludge until the end of the decay experiments, *E. coli* P17 was no longer recovered from the sludge by culture-based methods after 96 h of microcosm establishment (data not shown). These data suggest that *E. coli* P17 possibly transitioned to a viable but nonculturable (VBNC) state that is consistent with the persister cell hypothesis.

Persister cells had been studied in clinical settings, as this dormant state allows antibiotic-susceptible bacterial populations to survive antimicrobial treatments (14). To the best of our knowledge, there is only one report of environmental incidence of a dormant but infective state of the fish pathogen *Pasteurella piscicida* (46). However, the importance of these specialized cells for the survival and dissemination of *E. coli* in the environment, particularly in the wastewater treatment system, has been overlooked (47). The occurrence of persisters of pathogenic strains in the sludge (such as *E. coli* P17) has implications for the management and disposal of the sludge originating from WWTPs and their respective treated effluents. As this subpopulation of bacteria exhibits high tolerance to antimicrobials, this raises the question of whether our current disinfection practices are effective at fully inactivating persisters of pathogenic bacteria that remain in the treated wastewater. This can raise potential concerns when treated wastewater is intended for reuse, as improper inactivation might represent a direct risk to the public health.

An earlier study in reactors treating wastewater suggested that as a general trend, ARGs are removed more efficiently under anaerobic conditions (9). However, in a study by Diehl and LaPara (9), the anaerobic reactors were operated at a longer retention time than the aerobic reactors. This difference in the operational parameters of the systems makes it difficult to determine whether the improved ARG removal corresponded to factors related to the anaerobic/aerobic condition or to differences in the retention times of the two types of systems. Nonetheless, it is important to highlight that Diehl and LaPara (9) observed that some particular ARGs showed prolonged persistence under the anaerobic condition, suggesting that the differential ARG decay is dependent on each particular ARG. In agreement with this observation, Burch and collaborators (10) concluded that the removal rates of ARGs vary substantially depending on the specific ARG. In a more detailed study, Rysz and collaborators (48) indicated that the effect of oxygen availability on the maintenance of tetracycline resistance genes is also dependent on the host cell or, more specifically, on the particular plasmid-host pair. Anaerobic conditions led to complete loss of a plasmid carrying the tetracycline resistance gene (the RP1 plasmid) in *Pseudomonas aeruginosa*, while *E. coli* retained its

tetracycline resistance plasmid (pSC101) for more than 500 generations. The pSC101 and *bla*_{NDM-1} plasmids harbored by *E. coli* P17 (pKOX_NDM-1) carry a toxin-antitoxin system that might improve their retention in the *E. coli* host. Moreover, these two plasmids belong to the IncF family, which shows high stability in *Enterobacteriaceae* (49–52). These two characteristics shared by pSC101 and pKOX_NDM-1 might have accounted for their retention under energy-deprived anaerobic conditions. As the particular interaction of pKOX_NDM-1 and *E. coli* P17 is highly stable, cellular decay of *E. coli* P17 is the main factor driving the decay kinetics of *bla*_{NDM-1} in our microcosm experiments.

In a complex microbial community where competition for resources takes place (42), we expected that the addition of meropenem would have resulted in increased fitness of *E. coli* P17 in the anaerobic and aerobic sludge. On the contrary, such antibiotic concentrations did not provide an evident selective advantage to *E. coli* P17 that resulted in prolonged persistence. It is important to highlight that *bla*_{NDM-1} is fully functional in *E. coli* P17, with an MIC of 64 $\mu\text{g/ml}$ (19), suggesting that the possible adaptive advantage imposed by the meropenem addition might be negligible compared to the other negative pressures experienced by *E. coli* in the sludge environment.

Although we did not detect any effects of trace antibiotic concentrations on the survival of *E. coli* P17 or on *bla*_{NDM-1} persistence, other studies suggested that exposure to sublethal antibiotic and disinfectant concentrations stimulate plasmid conjugation rates in activated sludge (53) and in water (54). Indeed, transconjugants were recovered from the aerobic microcosms. As IncF plasmids are unique to *Enterobacteriaceae* (55), all transconjugants isolated fall within this taxonomical unit, which includes many genera associated with waterborne pathogens, such as *Salmonella*, *Yersinia*, *Klebsiella*, *Shigella*, *Citrobacter*, etc. (56, 57). In contrast, no transconjugants were recovered from the anaerobic microcosms, most likely because such HGT events took place with fastidious bacteria highly prevalent in anaerobic environments (58), and these bacteria cannot be easily cultivated. Similarly, after 120 h of aerobic microcosm establishment, transconjugants were no longer recovered from the aerobic sludge (Fig. 1d). Lower plasmid stability of the new host-plasmid interaction is a plausible explanation for the loss of transconjugant recovery. However, IncF plasmids are highly stable in the *Enterobacteriaceae* (49–52), and such stability is likely improved by the toxin-antitoxin system (19). Due to the phylogenetic proximity of *Shigella* and *Citrobacter* spp. to *E. coli* (59), it is speculated that this plasmid might also be stable in these hosts. Hence, an alternative explanation for the lack of transconjugant recovery is the loss of culturability of such bacteria, since the progression to a viable but nonculturable (VBNC) state is a survival strategy common among diverse groups of bacteria (60).

The contribution of HGT and the factors affecting the dissemination of plasmids in activated sludge under laboratory conditions have been well documented. These factors include plasmid host range (61), sludge retention times (62), stressful environmental conditions (53), and host/donor phylogenetic affiliations (63). Significant enrichment of the ratios of multiple ARGs compared to 16S rRNA genes throughout the wastewater treatment process confirms the potential for mobility and proliferation of ARGs within the activated sludge microbial communities in full-scale treatment systems (64). *bla*_{NDM-1} HGT events involving activated sludge microorganisms and soil bacteria had also been reported (8). In agreement with these studies, our results emphasize the role of activated sludge as an environmental reservoir of plasmid-carried ARGs, as well as its potential role in mediating the dissemination of ARGs into the environment and to other *Enterobacteriaceae*.

Subsequently, upon cellular decay of *E. coli* P17 and the previously identified transconjugants, *bla*_{NDM-1} can be released to the nonsettleable or colloidal fraction of the sludge. The plasmid DNA (pDNA) decay experiments showed that although the decay at the gene level is considerably slower (measured by qPCR), the fragmentation process of naked DNA in the liquid aerobic fraction leads to rapid decay in the replicative structure of the plasmid (measured by electroporation assays). Circular plasmids replicate by rolling-circle, strand displacement, or theta replication. In these

three replication mechanisms, the circular structure of at least one DNA strand must be maintained in order to complete the replication cycle (65). A single double-strand break would linearize the plasmid, resulting in the disruption of the replication process by any of these mechanisms. Ultimately, the lack of replicative functions would compromise the ability of the plasmid to be disseminated and maintained in the transformed bacterial host.

DNA degradation in the environment is a complex multifactorial process involving chemical and biological aspects (66). DNase-mediated degradation is the main biological process driving the decay of extracellular DNA in the environment (67, 68), while low temperatures, high salinity, high levels of organic matter, and anoxic environments are some of the physicochemical factors that contribute to the preservation of environmental DNA (66, 69). Determining the factors influencing the persistence of DNA in sludge liquid fraction is outside the scope of this study. However, it was determined that DNA persisted for a longer time in the liquid fraction of the anaerobic sludge than in that of the aerobic sludge, ultimately increasing the probability of subsequent bla_{NDM-1} uptake and fixation by environmental bacteria. In a previous study, it was found that membranes of 100 kDa and smaller could achieve significant removal of ARGs, attaining up to 4.5-log removal of colloidal DNA (70). Hence, coupling membrane separation with the activated sludge process can serve to mitigate microbial risk associated with the presence of persisters and extracellular DNA in the treated effluent.

In summary, the results from this study highlight the higher potential of dissemination of *E. coli* PI7 and ARGs associated with prolonged host and extracellular DNA persistence in anaerobic sludge. In the aerobic sludge, this study demonstrates transconjugation of plasmids encoding ARGs to compatible bacteria within the sludge, highlighting the likelihood of potential horizontal gene transfer events. Furthermore, this study emphasizes the potential importance of persister cells in the survival and dissemination of enteric pathogens into the natural environment and suggests a certain extent of indirect and direct risk to public health imposed by the presence of persister cells of pathogenic strains remaining in the sludge and effluents of WWTPs.

MATERIALS AND METHODS

Microcosm preparation. Two sets of microcosm experiments, representing anaerobic and aerobic biological reactors, were established in 1-liter sterile Pyrex bottles. Anaerobic and aerobic microcosms were seeded with 800 ml of sludge (mixed liquor suspended solids [MLSS] of 4 to 5 g/liter) from a lab scale anaerobic bioreactor (38) and an aerobic full-scale WWTP, respectively. The sludge in the anaerobic bioreactor was comprised of camel feces and anaerobic sludge from an industrial WWTP in Riyadh, and the anaerobic bioreactor had been in operation for more than 3 years (38, 71, 72). The full-scale WWTP was located at King Abdullah University of Science and Technology in Thuwal, Saudi Arabia, and had a capacity of 1,600 m³/day. Hydraulic retention time (HRT) and sludge retention time (SRT) in the activated sludge tank were 2.5 h and 40 days, respectively. The sludge tanks were operated at an average temperature of 33°C with 1 to 2 mg/liter of dissolved oxygen at pH 7 to 8. Sludge for each replicate run ($n = 3$) was recovered at three different time points scattered along a 1-year period for both anaerobic and aerobic seed sludge. Prior to the *E. coli* PI7 spiking event, the seed sludge was acclimated for 10 days and screened for the presence of bla_{NDM-1} or *uidA*. No bla_{NDM-1} or *uidA* copies were detected in anaerobic and aerobic sludge by either endpoint PCR or cultivation methods before *E. coli* PI7 spiking.

Decay experiments. The *E. coli* PI7 inoculum to be spiked into the microcosms was grown in LB broth at 37°C for 8 h to an optical density at 600 nm (OD_{600}) of 0.7. Once this cell density was reached, 100 ml of culture were spiked into each microcosm, resulting in a final volume of 900 ml per microcosm. The addition of this large quantity of *E. coli* PI7 cells into each individual microcosm, albeit not representative of actual conditions in WWTPs, was similar to approaches undertaken in earlier studies (73, 74). The high cell density spiked into the microcosm was required to ensure final cell density upon decay remained within the limits of detection by qPCR, and also to allow detection of subpopulations (e.g., persister cells) that generally occur in low cell densities. Decay experiments were performed at different trace antibiotic concentrations. In the anaerobic microcosms, decay was evaluated at 0 and 100 μ g/liter of meropenem ($n = 2$ per replicate run) (Sigma-Aldrich, St. Louis, MO, USA). In the aerobic microcosm, concentrations of 0, 1, 10, and 100 μ g/liter of meropenem were tested ($n = 4$ per replicate run). Anaerobic microcosms were not tested with 1 and 10 μ g/liter meropenem because of low sludge production from the anaerobic bioreactor, and this restricted the number of microcosms that could be set up. Concentrations were chosen to represent the concentration range of organic micropollutants commonly reported in municipal wastewater (75, 76). Three independent replicate runs for each aerobic and anaerobic set-up were performed, comprising a total of 6 and 12 anaerobic and aerobic microcosms, respectively. Microcosms were operated as sequencing batch reactors at a constant temperature of 37°C.

TABLE 2 Primers and probes used in this study

Primer or probe	Gene target	Amplicon size (bp)	Sequence ^a	Cycling conditions	Use
Primers					
NDM154-F ^b	<i>bla</i> _{NDM-1}	154	ATTAGCCGCTGCATTGAT	50°C × 2 min; 95°C × 20 s; 40 cycles of 95°C × 1 s	qPCR
NDM154-R ^b			CATGTCGAGATAGGAAGTG	and 60°C × 20 s	
uidA159-F ^c	<i>uidA</i>	159	CGAATCCTTTGCCACGCAAG	50°C × 2 min; 95°C × 20 s; 40 cycles of 95°C × 1 s	qPCR
uidA159-R ^c			TCACAGCCAAAAGCCAGACA	and 60°C × 20 s	
NDM640-F	<i>bla</i> _{NDM-1}	640	TAGTGCTCAGTGTCG	95°C × 3 min; 35 cycles of 95°C × 30 s, 60°C × 30 s	Sequencing
NDM640-R			CATTAGCCGCTGCA	and 72°C × 1 min; final elongation 72°C × 5 min	(transconjugant screening)
11F	16S	1481	GTTYGATYCTGGCTCAG	95°C × 5 min; 35 cycles of 95°C × 1 min, 45°C × 45 s	Sequencing
1492R			GGYTACCTTGTTACGACTT	and 72°C × 2 min; final elongation 72°C × 10 min	(transconjugant identity)
Probes					
NDM-22 ^b	<i>bla</i> _{NDM}		56-FAM/AGACATTCG/ZEN/ GTGCGAGCTGGCGGA/ 3IABkFQ	As described for primer pair	qPCR
uidA-23 ^c	<i>uidA</i>		56-FAM/TCGCCCTTC/ZEN/ ACTGCCACTGACCG/ 3IABkFQ	As described for primer pair	qPCR

^aFAM, 6-carboxyfluorescein.

^bPrimer-probe pair NDM-154 and NDM-22 qPCR amplification efficiency was 98%.

^cPrimer-pair uidA159 and uidA-23 qPCR amplification efficiency was 102%.

The aerobic microcosms were aerated with atmospheric air at 250 ml/min, while the anaerobic microcosms were established in airtight bottles and mechanically stirred at 250 rpm. To prevent oxygen intrusion, anaerobic microcosms were established and sampled in a vinyl anaerobic chamber (Coy Lab Products, MI). After each feeding/sampling event, anaerobic microcosms were purged with 99.9% nitrogen for 15 min. MLSS and pH were measured every 24 h for all microcosms, and pH was maintained at 7.2 ± 0.4 using HEPES buffer at a final concentration of 25 mM. Daily, 100 ml of liquid was removed from the aerobic and anaerobic microcosms and replaced with synthetic wastewater (38) that had the corresponding concentration of antibiotic, achieving a food-to-microorganism ratio (F/M) of 0.2. Biomass was separated from the nonsettling liquid fraction by centrifugation at $9,800 \times g$ for 20 to 30 min in a sterile ultracentrifuge bottle, and reintroduced into the respective microcosm.

Sample processing. At each sampling event, 14 ml of sludge was collected from each microcosm and a subset was used for (i) total biomass DNA isolation (2 ml), (ii) sludge exposure to propidium monoazide (PMA) (0.5 ml), and (iii) detection of horizontal gene transfer (HGT) events by culture-based methods (0.5 ml). Sludge exposure to PMA was performed as previously described (77). Given that PMA can be limited in its accuracy to differentiate between cells with compromised and intact cell membranes, more details on PMA exposure protocol and validation are given in Text S1-3 and Table S1 in the supplemental material.

Sludge samples for total DNA isolation and PMA-treated samples were centrifuged at $10,000 \times g$ for 5 min and the supernatant fraction was discarded. The remaining biomass pellet was immediately frozen at -80°C , and subsequently lyophilized using the Alpha 1-2 LDplus freeze dryer (Martin Christ GmbH, Germany). After completion of the drying cycle, dry biomass weights were recorded and samples were ready for DNA extraction.

Horizontal gene transfer detection by culture-based methods. To detect potential conjugation events, the remaining 0.5 ml of the initial 14 ml sludge sample were serially diluted and plated on MUG-EC medium (Sigma-Aldrich, St. Louis, MO) with 1.5% (wt/vol) agar and 8 $\mu\text{g/ml}$ meropenem. MUG-EC allows the rapid screening of *E. coli* isolates, as MUG (methylumbelliferyl- β -glucuronide) cleavage by the glucuronidase enzyme leads to the formation of colonies that exhibit blue fluorescence under UV (78). Nonfluorescent colonies were selected as potential transconjugants, and subsequently confirmed by colony PCR using *bla*_{NDM-1}-specific primers as shown in Table 2. Strain identity was determined by 16S rRNA gene sequencing using the 11F/1492R primer pair (79). The detection limit of culture-based methods performed in this study was determined to be 10^4 CFU/g sludge (see Text S1-4 in the supplemental material).

Colloidal DNA decay. *E. coli* PI7 carries a 110-kbp plasmid encoding NDM-1 (19). Plasmid integrity was determined by electroporation into Invitrogen TOP10 electrocompetent cells (Thermo Fisher Scientific, Carlsbad, CA, USA). Due to the low electroporation frequencies of this large plasmid, decay experiments were carried out using the pCR2.1 cloning vector (Thermo Fisher Scientific, Carlsbad, CA, USA) that harbors a 640-bp *bla*_{NDM-1} gene insertion (i.e., a total plasmid vector size of ~ 4.6 kbp). Decay experiments for the plasmid were performed in the liquid fraction of either aerobic or anaerobic sludge collected from a local WWTP and a lab scale anaerobic membrane bioreactor (AnMBR), respectively. The sludge liquid fraction was separated from the biomass by centrifugation at $10,000 \times g$ for 20 min. Supernatant was recovered and filtered with cheesecloth to further remove biomass in suspension.

Subsequently, 1.6 ml of plasmidic DNA (10^{10} copies/ μ l) were dosed into a Float-A-Lyzer G2 dialysis device with a molecular mass cutoff of 100 kDa (Spectrum Laboratories, Rancho Dominguez, CA), and the device was submerged in the liquid fraction of either aerobic or anaerobic sludge for a period of 24 h. Anaerobic decay experiments were performed under anaerobic conditions in a vinyl anaerobic chamber with an atmosphere of 95% nitrogen and 5% hydrogen (Coy Lab Products, MI, USA). A total of 50 μ l samples were taken from the dialysis device at each sampling point, and used for (i) plasmidic DNA quantification with qPCR and (ii) plasmid integrity quantification by electroporation in TOP10 cells (see Text SI-5 in the supplemental material).

DNA extraction. Lyophilized PMA-treated sludge and non-PMA-treated sludge fractions were subjected to DNA extraction using the PowerSoil DNA extraction kit (Mo Bio) with slight modifications to the manufacturer's protocol, as previously described (4).

Gene quantification and statistical tests. The *uidA* and *bla*_{NDM-1} genes were used as the chromosomal and plasmidic markers, respectively. β -D-Glucuronidase (*uidA*) was selected as the chromosomal marker since only one copy of this gene is present per *E. coli* genome (19). Differences in the decay of the chromosomal and plasmidic material of *E. coli* P17 were therefore assessed by comparing PMA-treated biomass (PTB) decay rates of *uidA* and *bla*_{NDM-1} genes, respectively. *bla*_{NDM-1} and *uidA* copy numbers were determined by absolute quantification on an Applied Biosystems 7900HT Fast real-time PCR system (Thermo Fisher Scientific, Carlsbad, CA, USA). PCR primers and TaqMan probe sequences are listed in Table 2. MLSS measurements were fairly stable in the microcosm experiments during the whole length of the experiment (see Fig. S3 in the supplemental material), suggesting stability of both aerobic and anaerobic microbial communities in the microcosms. Gene copy numbers were normalized by dry weight of biomass. Decay rates were expressed as $\ln(N/N_0)$, where N corresponds to the copy numbers at $t = X_i$ and N_0 corresponds to the copy numbers at $t = 0$. Half-life was calculated using a first-order decay kinetic model. Linear regressions were performed using the least-squares method and the significance of the slopes of the regression models ($\beta \neq 0$) was evaluated using the t test. Decay curves were compared using the model $Y_i = \beta_0 + \beta_1 X_{i1} + \beta_2 X_{i2} + \beta_3 X_{i1} X_{i2}$ (80). All statistical analysis was done using StatPlus at 95% confidence unless otherwise stated.

Determination of persister cell frequency in pure cultures of *E. coli* P17. Persister cells are dormant variations of vegetative cells that can withstand harsh environmental conditions, including exposure to supra-lethal concentrations of antibiotics. This resistance to antibiotics is not encoded in the chromosome but is rather a consequence of their dormant phenotype. The frequency of persister cells in this study was determined using a modified protocol described by Keren et al. (15). Modifications were made to the type of antibiotic used and the working concentration. As *E. coli* P17 exhibits extremely high tolerance to ampicillin, the antibiotic challenge was performed with meropenem, a carbapenem exhibiting a similar mode of action (7). *E. coli* P17 was challenged with a supra-lethal meropenem concentration of 640 μ g/ml, corresponding to a 10-fold increase of the MIC and a 5-fold increase of the lethal meropenem concentration reported for this strain (19). In summary, overnight cultures of *E. coli* P17 were diluted 1:1,000 in LB broth without meropenem and incubated at 37°C and 200 rpm to a final OD₆₀₀ of 0.2. To provide a baseline cell count ($t = 0$), 10 ml of culture from each flask ($n = 4$) were pooled and placed at 4°C. Subsequently, cell cultures were challenged with meropenem (640 μ g/ml), and incubated at 37°C and 200 rpm for 180 min. Cell counts were determined by serial dilution and plating in MUG-EC agar plates. Persister cell frequency was expressed as the ratio of the cell count at $t = 180$ min to the cell count at $t = 0$.

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/AEM.00640-17>.

SUPPLEMENTAL FILE 1, PDF file, 0.2 MB.

ACKNOWLEDGMENTS

The research reported in this publication was supported by funding from King Abdullah University of Science and Technology (KAUST) (grants BAS/1/1033-01-01 and FCC/1/1971-15-01) awarded to P.-Y. Hong.

REFERENCES

- McCarty PL, Bae J, Kim J. 2011. Domestic wastewater treatment as a net energy producer—can this be achieved? *Environ Sci Technol* 45: 7100–7106. <https://doi.org/10.1021/es2014264>.
- Smith AL, Stadler LB, Love NG, Skerlos SJ, Raskin L. 2012. Perspectives on anaerobic membrane bioreactor treatment of domestic wastewater: a critical review. *Bioresour Technol* 122:149–159. <https://doi.org/10.1016/j.biortech.2012.04.055>.
- Rizzo L, Manaia C, Merlin C, Schwartz T, Dagot C, Ploy MC, Michael I, Fatta-Kassinos D. 2013. Urban wastewater treatment plants as hotspots for antibiotic resistant bacteria and genes spread into the environment: a review. *Sci Total Environ* 447:345–360. <https://doi.org/10.1016/j.scitotenv.2013.01.032>.
- Al-Jassim N, Ansari MI, Harb M, Hong PY. 2015. Removal of bacterial contaminants and antibiotic resistance genes by conventional wastewater treatment processes in Saudi Arabia: is the treated wastewater safe to reuse for agricultural irrigation? *Water Res* 73:277–290. <https://doi.org/10.1016/j.watres.2015.01.036>.
- Novo A, Manaia CM. 2010. Factors influencing antibiotic resistance burden in municipal wastewater treatment plants. *Appl Microbiol Biotechnol* 87:1157–1166. <https://doi.org/10.1007/s00253-010-2583-6>.
- Yong D, Toleman MA, Giske CG, Cho HS, Sundman K, Lee K, Walsh TR. 2009. Characterization of a new metallo-beta-lactamase gene, *bla*_{NDM-1}, and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. *Antimi-*

- croB Agents Chemother 53:5046–5054. <https://doi.org/10.1128/AAC.00774-09>.
7. Papp-Wallace KM, Endimiani A, Taracila MA, Bonomo RA. 2011. Carbapenems: past, present, and future. *Antimicrob Agents Chemother* 55:4943–4960. <https://doi.org/10.1128/AAC.00296-11>.
 8. Luo Y, Yang F, Mathieu J, Mao D, Wang Q, Alvarez PJJ. 2014. Proliferation of multidrug-resistant New Delhi metallo- β -lactamase genes in municipal wastewater treatment plants in northern China. *Environ Sci Technol Lett* 1:26–30. <https://doi.org/10.1021/ez400152e>.
 9. Diehl DL, LaPara TM. 2010. Effect of temperature on the fate of genes encoding tetracycline resistance and the integrase of class 1 integrons within anaerobic and aerobic digesters treating municipal wastewater solids. *Environ Sci Technol* 44:9128–9133. <https://doi.org/10.1021/es102765a>.
 10. Burch TR, Sadowsky MJ, LaPara TM. 2013. Aerobic digestion reduces the quantity of antibiotic resistance genes in residual municipal wastewater solids. *Front Microbiol* 4:17. <https://doi.org/10.3389/fmicb.2013.00017>.
 11. Christgen B, Yang Y, Ahammad S, Li B, Rodriguez DC, Zhang T, Graham DW. 2015. Metagenomics shows that low-energy anaerobic-aerobic treatment reactors reduce antibiotic resistance gene levels from domestic wastewater. *Environ Sci Technol* 49:2577–2584. <https://doi.org/10.1021/es505521w>.
 12. Tao C-W, Hsu B-M, Ji W-T, Hsu T-K, Kao P-M, Hsu C-P, Shen S-M, Shen T-Y, Wan T-J, Huang Y-L. 2014. Evaluation of five antibiotic resistance genes in wastewater treatment systems of swine farms by real-time PCR. *Sci Total Environ* 496:116–121. <https://doi.org/10.1016/j.scitotenv.2014.07.024>.
 13. Lewis K. 2010. Persister cells. *Annu Rev Microbiol* 64:357–372. <https://doi.org/10.1146/annurev.micro.112408.134306>.
 14. Lewis K. 2007. Persister cells, dormancy and infectious disease. *Nat Rev Microbiol* 5:48–56. <https://doi.org/10.1038/nrmicro1557>.
 15. Keren I, Shah D, Spoering A, Kaldalu N, Lewis K. 2004. Specialized persister cells and the mechanism of multidrug tolerance in *Escherichia coli*. *J Bacteriol* 186:8172–8180. <https://doi.org/10.1128/JB.186.24.8172-8180.2004>.
 16. Mulcahy LR, Burns JL, Lory S, Lewis K. 2010. Emergence of *Pseudomonas aeruginosa* strains producing high levels of persister cells in patients with cystic fibrosis. *J Bacteriol* 192:6191–6199. <https://doi.org/10.1128/JB.01651-09>.
 17. Schumacher MA, Balani P, Min J, Chinnam NB, Hansen S, Vulić M, Lewis K, Brennan RG. 2015. HipBA-promoter structures reveal the basis of heritable multidrug tolerance. *Nature* 524:59–64. <https://doi.org/10.1038/nature14662>.
 18. LaFleur MD, Kumamoto CA, Lewis K. 2006. *Candida albicans* biofilms produce antifungal-tolerant persister cells. *Antimicrob Agents Chemother* 50:3839–3846. <https://doi.org/10.1128/AAC.00684-06>.
 19. Mantilla-Calderon D, Jumat MR, Wang T, Ganesan P, Al-Jassim N, Hong P-Y. 2016. Isolation and characterization of NDM-positive *Escherichia coli* from municipal wastewater in Jeddah, Saudi Arabia. *Antimicrob Agents Chemother* 60:5223–5231. <https://doi.org/10.1128/AAC.00236-16>.
 20. Johnson AP, Woodford N. 2013. Global spread of antibiotic resistance: the example of New Delhi metallo-beta-lactamase (NDM)-mediated carbapenem resistance. *J Med Microbiol* 62:499–513. <https://doi.org/10.1099/jmm.0.052555-0>.
 21. Nordmann P, Poirel L, Walsh TR, Livermore DM. 2011. The emerging NDM carbapenemases. *Trends Microbiol* 19:588–595. <https://doi.org/10.1016/j.tim.2011.09.005>.
 22. Ahammad ZS, Sreerishnan TR, Hands CL, Knapp CW, Graham DW. 2014. Increased waterborne bla_{NDM-1} resistance gene abundances associated with seasonal human pilgrimages to the upper Ganges River. *Environ Sci Technol* 48:3014–3020. <https://doi.org/10.1021/es405348h>.
 23. Walsh TR, Weeks J, Livermore DM, Toleman MA. 2011. Dissemination of NDM-1 positive bacteria in the New Delhi environment and its implications for human health: an environmental point prevalence study. *Lancet Infect Dis* 11:355–362. [https://doi.org/10.1016/S1473-3099\(11\)70059-7](https://doi.org/10.1016/S1473-3099(11)70059-7).
 24. Wang B, Sun D. 2015. Detection of NDM-1 carbapenemase-producing *Acinetobacter calcoaceticus* and *Acinetobacter junii* in environmental samples from livestock farms. *J Antimicrob Chemother* 70:611–613. <https://doi.org/10.1093/jac/dku405>.
 25. Michael I, Rizzo L, McArdell C, Manaia C, Merlin C, Schwartz T, Dagot C, Fatta-Kassinos D. 2013. Urban wastewater treatment plants as hotspots for the release of antibiotics in the environment: a review. *Water Res* 47:957–995. <https://doi.org/10.1016/j.watres.2012.11.027>.
 26. Heddi B, Abi-Ghanem J, Lavigne M, Hartmann B. 2010. Sequence-dependent DNA flexibility mediates DNase I cleavage. *J Mol Biol* 395:123–133. <https://doi.org/10.1016/j.jmb.2009.10.023>.
 27. Wagner AO, Malin C, Gstraunthaler G, Illmer P. 2009. Survival of selected pathogens in diluted sludge of a thermophilic waste treatment plant and in NaCl-solution under aerobic and anaerobic conditions. *Waste Manag* 29:425–429. <https://doi.org/10.1016/j.wasman.2008.03.003>.
 28. Pandey PK, Biswas S, Vaddella VK, Soupir ML. 2015. *Escherichia coli* persistence kinetics in dairy manure at moderate, mesophilic, and thermophilic temperatures under aerobic and anaerobic environments. *Bio-process Biosyst Eng* 38:457–467. <https://doi.org/10.1007/s00449-014-1285-3>.
 29. Semenov AV, van Overbeek L, Termorshuizen AJ, van Bruggen AH. 2011. Influence of aerobic and anaerobic conditions on survival of *Escherichia coli* O157:H7 and *Salmonella enterica* serovar Typhimurium in Luria-Bertani broth, farm-yard manure and slurry. *J Environ Manage* 92:780–787. <https://doi.org/10.1016/j.jenvman.2010.10.031>.
 30. Roslev P, Bjergbæk LA, Hesselsoe M. 2004. Effect of oxygen on survival of faecal pollution indicators in drinking water. *J Appl Microbiol* 96:938–945. <https://doi.org/10.1111/j.1365-2672.2004.02209.x>.
 31. Merlin C, Bonot S, Courtois S, Block JC. 2011. Persistence and dissemination of the multiple-antibiotic-resistance plasmid pB10 in the microbial communities of wastewater sludge microcosms. *Water Res* 45:2897–2905. <https://doi.org/10.1016/j.watres.2011.03.002>.
 32. Imlay JA. 2013. The molecular mechanisms and physiological consequences of oxidative stress: lessons from a model bacterium. *Nat Rev Microbiol* 11:443–454. <https://doi.org/10.1038/nrmicro3032>.
 33. Anjem A, Imlay JA. 2012. Mononuclear iron enzymes are primary targets of hydrogen peroxide stress. *J Biol Chem* 287:15544–15556. <https://doi.org/10.1074/jbc.M111.330365>.
 34. Kuo C, Mashino T, Fridovich I. 1987. alpha, beta-Dihydroxyisovalerate dehydratase: a superoxide-sensitive enzyme. *J Biol Chem* 262:4724–4727.
 35. Sakai A, Nakanishi M, Yoshiyama K, Maki H. 2006. Impact of reactive oxygen species on spontaneous mutagenesis in *Escherichia coli*. *Genes Cells* 11:767–778. <https://doi.org/10.1111/j.1365-2443.2006.00982.x>.
 36. Storz G, Imlay JA. 1999. Oxidative stress. *Curr Opin Microbiol* 2:188–194. [https://doi.org/10.1016/S1369-5274\(99\)80033-2](https://doi.org/10.1016/S1369-5274(99)80033-2).
 37. Korajkic A, Wanjugi P, Harwood VJ. 2013. Indigenous microbiota and habitat influence *Escherichia coli* survival more than sunlight in simulated aquatic environments. *Appl Environ Microbiol* 79:5329–5337. <https://doi.org/10.1128/AEM.01362-13>.
 38. Harb M, Xiong Y, Guest J, Amy G, Hong P-Y. 2015. Differences in microbial communities and performance between suspended and attached growth anaerobic membrane bioreactors treating synthetic municipal wastewater. *Environ Sci (Camb)* 1:800–813. <https://doi.org/10.1039/C5EW00162E>.
 39. Riviere D, Desvignes V, Pelletier E, Chaussonnerie S, Guermazi S, Weisenbach J, Li T, Camacho P, Sghir A. 2009. Towards the definition of a core of microorganisms involved in anaerobic digestion of sludge. *ISME J* 3:700–714. <https://doi.org/10.1038/ismej.2009.2>.
 40. Mahowald MA, Rey FE, Seedorf H, Turnbaugh PJ, Fulton RS, Wollam A, Shah N, Wang C, Magrini V, Wilson RK. 2009. Characterizing a model human gut microbiota composed of members of its two dominant bacterial phyla. *Proc Natl Acad Sci U S A* 106:5859–5864. <https://doi.org/10.1073/pnas.0901529106>.
 41. Saunders AM, Albertsen M, Vollertsen J, Nielsen PH. 2016. The activated sludge ecosystem contains a core community of abundant organisms. *ISME J* 10:11–20. <https://doi.org/10.1038/ismej.2015.117>.
 42. Zhang Q, He X, Yan T. 2015. Differential decay of wastewater bacteria and change of microbial communities in beach sand and seawater microcosms. *Environ Sci Technol* 49:8531–8540. <https://doi.org/10.1021/acs.est.5b01879>.
 43. Jeanneau L, Solecki O, Wery N, Jarde E, Gourmelon M, Communal PY, Jadas-Hecart A, Caprais MP, Gruau G, Pourcher AM. 2012. Relative decay of fecal indicator bacteria and human-associated markers: a microcosm study simulating wastewater input into seawater and freshwater. *Environ Sci Technol* 46:2375–2382. <https://doi.org/10.1021/es203019y>.
 44. Hellweger FL, Bucci V, Litman MR, Gu AZ, Onnis-Hayden A. 2009. Biphasic decay kinetics of fecal bacteria in surface water not a density effect. *J Environ Eng* 135:372–376. [https://doi.org/10.1061/\(ASCE\)0733-9372\(2009\)135:5\(372\)](https://doi.org/10.1061/(ASCE)0733-9372(2009)135:5(372)).
 45. Al-Jassim N, Mantilla-Calderon D, Wang T, Hong P. 2017. Inactivation of a virulent wastewater *Escherichia coli* and non-virulent commensal *Esch-*

- erichia coli* DSM1103 strains and their gene expression upon solar irradiation. *Environ Sci Technol* 51:3649–3659. <https://doi.org/10.1021/acs.est.6b05377>.
46. Magarinos B, Romalde J, Barja J, Toranzo A. 1994. Evidence of a dormant but infective state of the fish pathogen *Pasteurella piscicida* in seawater and sediment. *Appl Environ Microbiol* 60:180–186.
 47. van Elsas JD, Semenov AV, Costa R, Trevors JT. 2011. Survival of *Escherichia coli* in the environment: fundamental and public health aspects. *ISME J* 5:173–183. <https://doi.org/10.1038/ismej.2010.80>.
 48. Rysz M, Mansfield WR, Fortner JD, Alvarez PJ. 2013. Tetracycline resistance gene maintenance under varying bacterial growth rate, substrate and oxygen availability, and tetracycline concentration. *Environ Sci Technol* 47:6995–7001. <https://doi.org/10.1021/es3035329>.
 49. Ohtsubo H, Ryder TB, Maeda Y, Armstrong K, Ohtsubo E. 1986. DNA replication of the resistance plasmid R100 and its control. *Adv Biophys* 21:115–133. [https://doi.org/10.1016/0065-227X\(86\)90018-3](https://doi.org/10.1016/0065-227X(86)90018-3).
 50. Nordström K. 2006. Plasmid R1—replication and its control. *Plasmid* 55:1–26. <https://doi.org/10.1016/j.plasmid.2005.07.002>.
 51. Silver L, Chandler M, de la Tour EB, Caro L. 1977. Origin and direction of replication of the drug resistance plasmid R100.1 and of a resistance transfer factor derivative in synchronized cultures. *J Bacteriol* 131:929–942.
 52. Kline BC. 1985. A review of mini-F plasmid maintenance. *Plasmid* 14:1–16. [https://doi.org/10.1016/0147-619X\(85\)90027-7](https://doi.org/10.1016/0147-619X(85)90027-7).
 53. Kim S, Yun Z, Ha UH, Lee S, Park H, Kwon EE, Cho Y, Choung S, Oh J, Medriano CA, Chandran K. 2014. Transfer of antibiotic resistance plasmids in pure and activated sludge cultures in the presence of environmentally representative micro-contaminant concentrations. *Sci Total Environ* 468–469:813–820. <https://doi.org/10.1016/j.scitotenv.2013.08.100>.
 54. Zhang Y, Gu AZ, He M, Li D, Chen J. 2017. Subinhibitory concentrations of disinfectants promote the horizontal transfer of multidrug resistance genes within and across genera. *Environ Sci Technol* 51:570–580. <https://doi.org/10.1021/acs.est.6b03132>.
 55. Carattoli A. 2009. Resistance plasmid families in *Enterobacteriaceae*. *Antimicrob Agents Chemother* 53:2227–2238. <https://doi.org/10.1128/AAC.01707-08>.
 56. Maynard C, Berthiaume F, Lemarchand K, Harel J, Payment P, Bayardelle P, Masson L, Brousseau R. 2005. Waterborne pathogen detection by use of oligonucleotide-based microarrays. *Appl Environ Microbiol* 71:8548–8557. <https://doi.org/10.1128/AEM.71.12.8548-8557.2005>.
 57. Abbott SL. 2011. *Klebsiella*, *Enterobacter*, *Citrobacter*, *Serratia*, *Plesiomonas*, and other *Enterobacteriaceae*, p 639–657. In Versalovic J, Carroll K, Funke G, Jorgensen J, Landry M, Warnock D (ed), *Manual of clinical microbiology*, 10th ed. ASM Press, Washington, DC.
 58. Mori K, Kamagata Y. 2014. The challenges of studying the anaerobic microbial world. *Microbes Environ* 29:335–337. <https://doi.org/10.1264/jsm2.ME2904rh>.
 59. Paradis S, Boissinot M, Paquette N, Bélanger SD, Martel EA, Boudreau DK, Picard FJ, Ouellette M, Roy PH, Bergeron MG. 2005. Phylogeny of the *Enterobacteriaceae* based on genes encoding elongation factor Tu and F-ATPase β -subunit. *Int J Syst Evol Microbiol* 55:2013–2025. <https://doi.org/10.1099/ijs.0.63539-0>.
 60. Li L, Mendis N, Trigui H, Oliver JD, Faucher SP. 2014. The importance of the viable but non-culturable state in human bacterial pathogens. *Front Microbiol* 5:258. <https://doi.org/10.3389/fmicb.2014.00258>.
 61. von Wintersdorff CJ, Penders J, van Niekerk JM, Mills ND, Majumder S, van Alphen LB, Savelkoul PH, Wolffs PF. 2016. Dissemination of antimicrobial resistance in microbial ecosystems through horizontal gene transfer. *Front Microbiol* 7:173. <https://doi.org/10.3389/fmicb.2016.00173>.
 62. Tsutsui H, Anami Y, Matsuda M, Inoue D, Sei K, Soda S, Ike M. 2010. Transfer of plasmid pJP4 from *Escherichia coli* and *Pseudomonas putida* to bacteria in activated sludge developed under different sludge retention times. *J Biosci Bioeng* 110:684–689. <https://doi.org/10.1016/j.jbiosc.2010.07.009>.
 63. De Gelder L, Vandecasteele FP, Brown CJ, Forney LJ, Top EM. 2005. Plasmid donor affects host range of promiscuous IncP-1beta plasmid pB10 in an activated-sludge microbial community. *Appl Environ Microbiol* 71:5309–5317. <https://doi.org/10.1128/AEM.71.9.5309-5317.2005>.
 64. Mao D, Yu S, Rysz M, Luo Y, Yang F, Li F, Hou J, Mu Q, Alvarez PJ. 2015. Prevalence and proliferation of antibiotic resistance genes in two municipal wastewater treatment plants. *Water Res* 85:458–466. <https://doi.org/10.1016/j.watres.2015.09.010>.
 65. del Solar G, Giraldo R, Ruiz-Echevarría MJ, Espinosa M, Díaz-Orejas R. 1998. Replication and control of circular bacterial plasmids. *Microbiol Mol Biol Rev* 62:434–464.
 66. Corinaldesi C, Beolchini F, Dell'Anno A. 2008. Damage and degradation rates of extracellular DNA in marine sediments: implications for the preservation of gene sequences. *Mol Ecol* 17:3939–3951. <https://doi.org/10.1111/j.1365-294X.2008.03880.x>.
 67. Blum SAE, Lorenz MG, Wackernagel W. 1997. Mechanism of retarded DNA degradation and prokaryotic origin of DNases in nonsterile soils. *Syst Appl Microbiol* 20:513–521. [https://doi.org/10.1016/S0723-2020\(97\)80021-5](https://doi.org/10.1016/S0723-2020(97)80021-5).
 68. Dell'Anno A, Corinaldesi C. 2004. Degradation and turnover of extracellular DNA in marine sediments: ecological and methodological considerations. *Appl Environ Microbiol* 70:4384–4386. <https://doi.org/10.1128/AEM.70.7.4384-4386.2004>.
 69. Willerslev E, Cappellini E, Boomsma W, Nielsen R, Hebsgaard MB, Brand TB, Hofreiter M, Bunce M, Poinar HN, Dahl-Jensen D. 2007. Ancient biomolecules from deep ice cores reveal a forested southern Greenland. *Science* 317:111–114. <https://doi.org/10.1126/science.1141758>.
 70. Breazeal MV, Novak JT, Vikesland PJ, Pruden A. 2013. Effect of wastewater colloids on membrane removal of antibiotic resistance genes. *Water Res* 47:130–140. <https://doi.org/10.1016/j.watres.2012.09.044>.
 71. Wei C-H, Harb M, Amy G, Hong P-Y, Leiknes T. 2014. Sustainable organic loading rate and energy recovery potential of mesophilic anaerobic membrane bioreactor for municipal wastewater treatment. *Bioresour Technol* 166:326–334. <https://doi.org/10.1016/j.biortech.2014.05.053>.
 72. Harb M, Wei C-H, Wang N, Amy G, Hong P-Y. 2016. Organic micropollutants in aerobic and anaerobic membrane bioreactors: changes in microbial communities and gene expression. *Bioresour Technol* 218:882–891. <https://doi.org/10.1016/j.biortech.2016.07.036>.
 73. Oliver DM, Haygarth PM, Clegg CD, Heathwaite AL. 2006. Differential *E. coli* die-off patterns associated with agricultural matrices. *Environ Sci Technol* 40:5710–5716. <https://doi.org/10.1021/es0603249>.
 74. Jenkins MB, Fisher DS, Endale DM, Adams P. 2011. Comparative die-off of *Escherichia coli* O157:H7 and fecal indicator bacteria in pond water. *Environ Sci Technol* 45:1853–1858. <https://doi.org/10.1021/es1032019>.
 75. Yang S, Cha J, Carlson K. 2005. Simultaneous extraction and analysis of 11 tetracycline and sulfonamide antibiotics in influent and effluent domestic wastewater by solid-phase extraction and liquid chromatography-electrospray ionization tandem mass spectrometry. *J Chromatogr A* 1097:40–53. <https://doi.org/10.1016/j.chroma.2005.08.027>.
 76. Teerlink J, Hering AS, Higgins CP, Drewes JE. 2012. Variability of trace organic chemical concentrations in raw wastewater at three distinct sewershed scales. *Water Res* 46:3261–3271. <https://doi.org/10.1016/j.watres.2012.03.018>.
 77. Bae S, Wuertz S. 2009. Discrimination of viable and dead fecal *Bacteroides* bacteria by quantitative PCR with propidium monoazide. *Appl Environ Microbiol* 75:2940–2944. <https://doi.org/10.1128/AEM.01333-08>.
 78. Feng P, Hartman PA. 1982. Fluorogenic assays for immediate confirmation of *Escherichia coli*. *Appl Environ Microbiol* 43:1320–1329.
 79. Weisburg WG, Barns SM, Pelletier DA, Lane DJ. 1991. 16S ribosomal DNA amplification for phylogenetic study. *J Bacteriol* 173:697–703. <https://doi.org/10.1128/jb.173.2.697-703.1991>.
 80. Neter J, Kutner MH, Nachtsheim CJ, Wasserman W. 1996. *Applied linear statistical models*, 4th ed. McGraw-Hill/Irwin, Chicago, IL.