



Ectomycorrhizal Fungal Communities in Urban Parks Are Similar to Those in Natural Forests but Shaped by Vegetation and Park Age

Nan Hui,^a Xinxin Liu,^a D. Johan Kotze,^a Ari Jumpponen,^b Gaia Francini,^a Heikki Setälä^a

Department of Environmental Sciences, University of Helsinki, Lahti, Finland^a; Division of Biology, Kansas State University, Manhattan, Kansas, USA^b

ABSTRACT Ectomycorrhizal (ECM) fungi are important mutualists for the growth and health of most boreal trees. Forest age and its host species composition can impact the composition of ECM fungal communities. Although plentiful empirical data exist for forested environments, the effects of established vegetation and its successional trajectories on ECM fungi in urban greenspaces remain poorly understood. We analyzed ECM fungi in 5 control forests and 41 urban parks of two plant functional groups (conifer and broadleaf trees) and in three age categories (10, ~50, and >100 years old) in southern Finland. Our results show that although ECM fungal richness was marginally greater in forests than in urban parks, urban parks still hosted rich and diverse ECM fungal communities. ECM fungal community composition differed between the two habitats but was driven by taxon rank order reordering, as key ECM fungal taxa remained largely the same. In parks, the ECM communities differed between conifer and broadleaf trees. The successional trajectories of ECM fungi, as inferred in relation to the time since park construction, differed among the conifers and broadleaf trees: the ECM fungal communities changed over time under the conifers, whereas communities under broadleaf trees provided no evidence for such age-related effects. Our data show that plant-ECM fungus interactions in urban parks, in spite of being constructed environments, are surprisingly similar in richness to those in natural forests. This suggests that the presence of host trees, rather than soil characteristics or even disturbance regime of the system, determine ECM fungal community structure and diversity.

IMPORTANCE In urban environments, soil and trees improve environmental quality and provide essential ecosystem services. ECM fungi enhance plant growth and performance, increasing plant nutrient acquisition and protecting plants against toxic compounds. Recent evidence indicates that soil-inhabiting fungal communities, including ECM and saprotrophic fungi, in urban parks are affected by plant functional type and park age. However, ECM fungal diversity and its responses to urban stress, plant functional type, or park age remain unknown. The significance of our study is in identifying, in greater detail, the responses of ECM fungi in the rhizospheres of conifer and broadleaf trees in urban parks. This will greatly enhance our knowledge of ECM fungal communities under urban stresses, and the findings can be utilized by urban planners to improve urban ecosystem services.

KEYWORDS ectomycorrhizal fungal community, anthropogenic disturbance, park age, urban ecology, urban park, vegetation type

Received 15 August 2017 Accepted 21 September 2017

Accepted manuscript posted online 29 September 2017

Citation Hui N, Liu X, Kotze DJ, Jumpponen A, Francini G, Setälä H. 2017. Ectomycorrhizal fungal communities in urban parks are similar to those in natural forests but shaped by vegetation and park age. *Appl Environ Microbiol* 83:e01797-17. <https://doi.org/10.1128/AEM.01797-17>.

Editor Donald W. Schaffner, Rutgers, The State University of New Jersey

Copyright © 2017 American Society for Microbiology. All Rights Reserved.

Address correspondence to Nan Hui, nan.hui@helsinki.fi.

N.H. and X.L. contributed equally to this work.

Soils and trees in urban parks improve environmental quality and provide essential ecosystem services (1). Healthy urban trees facilitate rainwater storage, mitigate urban heat island effects, support biodiversity, and provide an esthetically appealing environment for urban residents (2). Ectomycorrhizal (ECM) fungi, necessary mutualists of most boreal trees, are important for host performance as well as for nutrient cycling at an ecosystem level (3). ECM fungi enhance plant growth and performance (4), increase plant nutrient acquisition (5), and protect plants against toxic compounds (6).

In urban parks, soil organisms are subject to anthropogenic disturbances, such as pollution, fertilization, trampling, and the removal of plant litter. ECM fungi are sensitive to urban disturbance (7), and their richness and abundance are lower in urban ecosystems than in rural areas (8, 9). Therefore, ECM fungi may be useful indicators reflecting the disturbance status of the below- and aboveground communities in urban areas, particularly where disturbances influence both soil properties and plant health. Although many factors influence ECM fungal community composition in boreal forest ecosystems, such as edaphic factors, host plant species composition, and stand age (10–12), the factors that impact urban ECM fungi and their succession remain unclear (9).

Most ECM fungal species typically have broad plant host ranges (13), and different hosts associate with divergent ECM fungal communities (14). ECM fungal communities can differ among urban and nonurban trees (9, 15), even among conspecific hosts (16). Recent evidence indicates that soil-inhabiting fungal communities (including ECM and saprotrophic fungi) in urban parks are affected by plant functional type and park age (17). However, ECM fungal diversity and its responses to urban stress, plant functional type, or park age remain unknown. In the current study, we extracted ECM fungal sequence data from a broader data set analyzed by Hui et al. (17). This allowed for exclusive analyses of ECM fungal responses in the rhizospheres of conifer and broadleaf trees in 41 parks in the cities of Lahti and Helsinki, Finland. The selected sites represent different park ages (i.e., time since park construction) and thus provide a means to dissect how ECM fungal communities are modified over time in an urban environment. To compare these park communities to those in a more natural and less disturbed environment, we included 5 minimally disturbed rural forests dominated by *Picea abies* and *Tilia cordata* as nonurban controls.

Here, we focus on (i) the response of ECM fungal communities to plant functional type and park age in urban park soils and (ii) potential differences in ECM fungal communities between nonurban control forests and disturbed urban parks (land use type). Further, we investigated (iii) which ECM fungal genera are particularly responsive to land use, plant functional type, and park age. We hypothesized that (i) ECM fungal communities under conifer and broadleaf trees in urban parks differ from those in control forests. This is because urban soils often have high pH and high concentrations of organic and inorganic pollutants (18), and their microbial communities may be affected by urban management (9). We also predicted that in the forest soil, ECM fungi are more diverse than in urban park soils. This is because of the positive relationship between canopy tree diversity and ECM fungal diversity (9, 11). (ii) ECM fungal community structure in urban parks depends on plant functional type. This is because plant functional types differ fundamentally in terms of effects on soil properties (18), allocation of recent photosynthate (19), litter, and root exudates (14), upon which ECM fungi depend. (iii) ECM fungal communities respond to park age. This is due to the different abilities of early and late-stage fungi to form symbioses with host roots (20).

RESULTS

Comparisons of ECM fungal communities between urban parks and control forests. The control forests and old parks in Lahti, representing roughly similarly aged trees, differed in ECM fungal diversity. Diversity was generally greater in control forests than in the old parks (Fig. 1, and Table S1 in the supplemental material). Operational taxonomic unit (OTU) richness and diversity were lower under conifer trees than broadleaf trees both in old parks and control forests (Fig. 1a and b), whereas evenness




FIG 1 Predicted (mean \pm standard error [SE]) ECM fungal OTU richness (a), diversity (b), and evenness (c) (GLMM results) in parks (both Lahti and Helsinki parks, left side) across plant functional types (symbols) and park ages (x axis), and between old Lahti parks and control forests (right side) across plant functional types.

showed an opposite trend (Fig. 1c). OTU richness, diversity, and evenness correlated positively with soil organic matter (OM). Soil pH correlated with ECM fungal community richness positively and with evenness negatively (Table S1).

ECM fungal community composition differed between (i) the old parks and control forests ($r^2 = 0.425$, $P < 0.001$) and (ii) the two tree functional types ($r^2 = 0.587$, $P < 0.001$; Fig. 2a). ECM fungal OTUs were classified into 51 genera throughout the data set. *Inocybe* was the most dominant genus (13.8% of the ECM fungal sequences [49 OTU]), followed by *Cenococcum* (11.7% [21 OTU]) and *Wilcoxina* (10.1% [4 OTU]). To explore the ECM fungal community distinctions between control forests and old parks in Lahti, we conducted generalized linear mixed models (GLMM) analyses on the 10 most abundant genera. Five genera (*Amphinema*, *Piloderma*, *Russula*, *Tomentella*, and *Tylospora*) were more abundant in control forests than in old parks (Fig. 3 and Table S1), while none of the most abundant genera occurred more frequently in the parks. *Cenococcum* and *Cortinarius* were constantly more abundant under broadleaf trees than conifers, whereas *Wilcoxina* showed an opposite trend. *Russula* and *Tylospora* showed significant plant functional type \times land use type (control forest versus old Lahti parks) interactions. In addition to these analyses, we also included a set of environmental variables in the GLMM analyses: four ECM fungal genera were correlated with soil N content (one positively and three negatively), three with soil C content (one positively and two negatively), three with soil OM content (all positively), four with percent sand (two positively and two negatively), and three with soil pH (two positively and one negatively) (Table S1).

Effects of plant functional group and park age on ECM fungi in urban parks. In parks, ECM fungal OTU richness and evenness had significant plant functional group \times




FIG 2 NMDS plots for ECM fungal communities. ECM fungal communities (a) of soils below broadleaf and conifer trees in old Lahti parks and control forests and ECM fungal communities (b) of soils below broadleaf trees in parks. Statistically significant ($P < 0.05$) vectors (soil pH, %N, %C, organic matter content, and percent sand) are shown. All NMDS plots showed significant differences ($P < 0.05$) either across land use type (control forest versus urban park), plant functional type, or park age by *envfit* analyses.

park age interactions (Table S2). In young parks, both ECM fungal richness and diversity were indistinguishable between the two plant functional groups. However, ca. 50 years after park establishment, soils under broadleaf trees tended to host more diverse ECM fungal communities than those under conifer trees (Fig. 1a and b). The diversity and evenness of the ECM fungal communities in parks were negatively correlated with soil pH. In our case, all soils were acidic, with a maximum pH ~ 6.9 (18). As a result, the diversity and evenness declined as pH approached neutral. All diversity indices correlated negatively with soil C content (Table S2).

ECM fungal community composition differed clearly between conifer and broadleaf trees in old parks in Lahti. As a result, we conducted analyses separately for the two tree functional groups in parks: ECM fungal communities under broadleaf trees responded to park age ($r^2 = 0.110$, $P = 0.029$; Fig. 2b), but this was not the case under conifer trees ($r^2 = 0.041$, $P = 0.697$).

To study the effects of plant functional type and park age on common ECM fungi in parks, we analyzed the 10 most abundant ECM fungal genera using GLMM. The abundances of *Inocybe*, *Wilcoxina*, and *Cenococcum* were similar, and the differences among the two plant functional types became more pronounced in intermediate and old parks than in young parks (Fig. 4). In intermediate and old parks, *Wilcoxina* was more abundant under conifer trees than under broadleaf trees, whereas *Cenococcum* and *Tuber* showed the opposite trend. *Tuber* was also highly abundant under conifer trees in young parks but not under broadleaf trees. *Hebeloma* and *Tomentella* abun-




FIG 3 Predicted count (mean ± SE) of the 10 most abundant ECM fungal genera across land use type (control forest versus old urban park) and plant functional type (GLMM results).

dances differed across park age, with a higher count in young parks than in intermediate and old parks. *Laccaria* and *Cortinarius* were consistently more abundant under broadleaf trees than under conifer trees, especially in old parks. *Cenococcum*, *Scleroderma*, and *Tomentella* showed plant functional type × park age interactions (Table S2). Our GLMM results showed that six ECM fungal genera were correlated with soil N content (all negatively), five with C content (four positively and one negatively), three with OM content (all negatively), seven with percent sand (one positively and six negatively), and two with pH (one positively and one negatively) (Table S2).

DISCUSSION

Our previous research showed that vegetation and park age drive changes in soil properties in urban parks (18), leading to distinct microbial communities (bacteria and fungi) (17). Here, we focused exclusively on ECM fungi and addressed how they respond to land use type (forest versus urban park), plant functional type, and park age under northern climatic conditions. Since different fungal groups have distinct life history strategies (21), we expected that ECM fungal responses would differ from those of the general soil-inhabiting fungi (17).

Differences in ECM fungal communities between old parks and forests. Urbanization likely has negative effects on soil properties (18), microbial communities (22),




FIG 4 Predicted count (mean ± SE) of the 10 most abundant ECM fungal genera across plant functional type and park age (GLMM results).

and soil fauna (23). Our data indicate that ECM fungal OTU richness and diversity were greater in control forests than in urban parks, supporting our first hypothesis and corroborating previous observations. Urban anthropogenic disturbance can reduce ECM fungal diversity and richness (8, 9, 24, 25), particularly in boreal regions where these fungi are most diverse. Our data are in contrast to predictions that increasing urbanization and the concomitant loss of natural forests will lead to the dramatic suppression of ECM fungi in urbanized ecosystems (22). Instead, we conclude that boreal hosts recruit quite diverse ECM fungi in urban greenspaces, suggesting ECM fungal community resistance and resilience to urbanization and cooccurring anthropogenic disturbances.

Despite the possible resistance and resilience, the ECM fungal communities in urban parks and control forests differed. This result was driven by taxon rank reordering, not taxon replacement. Urban disturbances (litter removal and raking, trampling, and mowing) are absent in forests, which likely result in alterations in the relative proportions of ECM fungal taxa between urban park and forest communities (9). Furthermore, unlike natural boreal forests typified by podzol soils with an organic matter layer developed on top of the soil, urban parks (even the oldest ones in our study) lacked

such a clear pedogenesis. The direct and indirect effects of pedogenesis on soil physical-chemical parameters are factors that likely affect ECM fungal communities between the two land use types. However, despite the absence of visible pedogenesis in the urban settings, ECM fungal richness and diversity were surprisingly similar between the natural and urban environments.

At the genus level, the 10 most abundant ECM fungal genera were present in both urban parks and forest stands, but the abundances of many of these genera differed between urban parks and forests. These observations are in line with previous reports (9). For example, the abundances of *Russula* and *Tylospora* were low in urban parks compared to those in control forests. This is in accordance with Hartmann et al., who showed that the abundance of *Russula* is negatively related to soil compaction (26). *Tylospora* occurs in decaying wood (27), which is scarce or absent in urban parks. *Tuber* was more frequent in urban parks than in control forests. This result is in agreement with Wang et al. (28), who predicted that *Tuber* may be “preadapted” to environmental conditions associated with human activities. *Tuber* species tend to prefer alkaline soils (28). A potential explanation for the observed greater abundance is that the acidic soil in control forests largely suppresses *Tuber* species, while they may survive in the neutral or weakly acid soils, which indeed typify urban environments (29).

Generally, our results suggest that ECM fungal richness and diversity were comparable in urban parks and control forests, albeit minimally different. ECM fungal communities shared a number of taxa between the land use types but were reordered and dominants replaced as indicated by our genus-level analyses.

Effects of plant functional type on ECM fungal communities in parks. Supporting previous findings (17, 30) and our second hypothesis, the two tree functional groups hosted distinct ECM fungal communities, both in their diversity and composition. The reasons for the observed compositional differences may lie in the plant-ECM fungus interaction. Plant functional types can influence ECM fungal communities in several ways, including effects through host specificity, modulation of edaphic conditions, litter quality and quantity (labile or recalcitrant), and rhizodeposition (root exudates) (14, 31–33). A recent meta-analysis revealed that host family explained 34% of the variation in ECM fungal community composition (34). This may be a result of specific molecular signaling between the host and its fungal symbionts that determines their compatibility (35).

Our previous studies show that plant functional types modify soils differently, and soils under conifer trees have lower pH but higher %OM, %N, and %C than soils under broadleaf trees in urban parks (18). In the current study, we found that six of the 10 most abundant ECM fungal genera in urban parks were negatively correlated with soil N content. Nitrogen content is a major factor influencing ECM fungal communities (35). High N supply suppresses biomass of ECM fungi, because when the host tree can easily obtain nutrients, there is no need to form such plant-fungus symbionts (36, 37). However, these chemical responses and correlations did not remarkably affect ECM fungal community richness and diversity.

Differences in the ECM fungal communities between the two tree functional types were largely attributable to shifts in the abundances of some ECM fungal genera. For example, *Cenococcum* was more common with broadleaf trees than conifers. Twieg et al. showed that the mean relative abundance of *Cenococcum* on broadleaf tree roots (paper birch [*Betula papyrifera*]) was about four times greater than on conifer trees (Douglas fir) in a mixed temperate forest (12). *Cenococcum*, one of the most common ECM fungal genera in boreal forest soil, seems to respond negatively to high nitrogen content in the soil (11). *Cenococcum* is common in soils with low nitrogen content; thus, as a result of the N deposition and subsequent higher N availability, the taxon declines (38, 39). Indeed, in our study, *Cenococcum* was negatively correlated with N in park soils that accumulate traffic-derived nitrogen (40). *Wilcoxina* spp., in turn, are generalists and are well adapted to a wide range of plant community types. They are often among the dominant ECM fungal taxa in coniferous forest (41, 42). Similarly, in parks, *Wilcoxina*

spp. were clearly associated with conifer trees. These results highlight that, despite the distinct environmental conditions in parks and forests, ECM fungus host preferences operate similarly regardless of land use. This highlights the pivotal role of plant identity in controlling plant-fungus symbiosis.

The effects of park age on the ECM fungal community. ECM fungal diversity seemed rather insensitive to park age. This finding supports neither our third hypothesis nor previous results by Twieg et al., who showed that ECM fungal diversity increased as stand age increased in soils under conifer trees in natural forests (12). Presumably, ECM fungal diversity would increase as plants grow older, because trees in young parks lack an extensive root system for ECM fungal colonization (43). Further, young soils may lack an extensive ECM fungal propagule bank because landfill top soils are common in park construction and because of insufficient propagule dispersal to recently established habitats (44). The relatively stable ECM fungal diversity that we observed across park ages may result from the minimal competition in the urban environments which allows many ECM fungi to rapidly colonize roots of young trees. It appears that young trees are equally suitable hosts for these ECM fungal spores to colonize their roots. Although park soil characteristics change by age (18), the modifications do not influence the colonization of ECM fungi, suggesting that ECM fungal spores are ubiquitously present in the urban environment. However, to our knowledge, studies that explicitly explore the effects of host age on ECM fungal communities in urban soils are nonexistent.

Despite the lack of an overall park age effect on ECM fungal community diversity, community composition responded to park age in the broadleaf tree rhizospheres. Similar responses were absent with the conifer hosts. The lack of this response under conifer trees is surprising, given that soil chemistry changes (lower pH and higher OM, C, and N content) were particularly pronounced underneath conifers in our parks (18). Previous studies on ECM fungal succession in natural forests suggest some context dependency of community responses to stand age. ECM fungal composition and diversity were insensitive to the age of oak (*Quercus ilex*) stands (45), whereas Kyaschenko et al. reported changes in the ECM fungus species composition along an age gradient of managed *Pinus sylvestris* stands (46). Taken together, these suggest that ECM fungal communities are primarily shaped by host-fungus interactions rather than by abiotic habitat conditions, such as soil chemistry.

Conclusion. Our results demonstrate that, in general, ECM fungi respond to land use type (urban park versus nonurban forest stands) and to plant functional types within parks and forests. Although ECM fungal richness was marginally greater in control forests than in urban parks, urban parks still hosted rich and diverse ECM fungal communities. ECM fungal community composition differed between the two habitats, but it was the common taxa that varied in abundance without clear taxon replacements, indicating that key ECM fungi remained mainly the same. In parks, ECM fungal community composition differed between conifer and broadleaf trees. Park age also proved to shape ECM fungal community composition, but this was evident under broadleaf trees only. Interestingly, plant functional group effects tended to be amplified in older parks where ECM fungi have had a longer time to interact with tree roots. We conclude that despite the lack of natural pedogenesis and arrested vegetation succession, as well as anthropogenic disturbance that includes raking leaves, mowing, and trampling, urban parks host a surprisingly diverse set of ECM fungi. Whether these urban ECM fungal communities functionally approximate those in natural forest stands requires further research.

MATERIALS AND METHODS

Study area and sampling design. The study sites have been described in detail previously (17, 18). Briefly, we selected 41 urban parks in the cities of Helsinki and Lahti, southern Finland, and 5 additional control forests in the proximity of Lahti. The urban parks represent different ages: more than 100 years old (the oldest parks were established over 2 centuries ago), 50 ± 10 years old, and 10 years old, referred to as old, intermediate, and young parks, respectively. We considered two plant functional types in these parks: conifer (43.3% of the conifer tree species represents *Picea* spp., 20% represents *Abies* spp., 13.3%

represents *Pseudotsuga menziesii*, 13.3% represents *Pinus sylvestris*, and 10% represents *Larix* sp.) and broadleaf (*Tilia* × *vulgaris* 100%) trees. With a few exceptions, conifer and broadleaf trees commonly existed together within a park. The distance between the two tree types was always greater than the height of the tallest tree. The age of plants within each park age class corresponded with park age, except for the young parks, where trees are commonly planted as ca. 10-year-old saplings at the time of park construction. The ideal experimental design would have included 15 parks per city, represented by five old, five intermediate, and five young parks, with both plant functional types present. However, since some parks did not include both plant functional types, we also selected parks with only one plant functional type. This resulted in a total of 41 urban parks and 58 urban sampling locations with 7 to 11 replicates per park age and plant functional type. Park sizes varied considerably, ranging from ca. 0.1 ha to several hectares, but with no systematic grouping of size with park age and plant functional type.

Soil sampling and edaphic conditions. ECM fungi colonize roots, but they grow from roots into the soil to deliver soil nutrients to the roots. Because of this, soil is a good proxy in studying ECM fungal communities and assigning the detected taxa to ecological roles (47–49). We sampled soils in May 2015 at the edge of the canopy projection so that distance to the nearest tree trunk ranged from 1 m (young parks; samples always collected outside the planting pit) to several meters (old parks). At each sampling point, we subsampled 3 soil cores (top 10 cm) using a steel push corer (10 cm deep, 2.54-cm diameter) and pooled the three subsamples into one composite for a total of 68 samples across the experiment (58 urban park samples and 10 control forest samples). The corer was sterilized using 70% ethanol between samples. Samples were stored in Minigrip bags on ice in the field and frozen at -20°C in the laboratory. Before DNA extraction, the samples were thawed at room temperature and sieved through a 2-mm mesh to remove any remaining large particles. The edaphic conditions (0 to 10 cm deep) of all samples were analyzed in our previous studies (18, 50). Five variables, carbon (C) content, nitrogen (N) content, organic matter (OM) content, percent sand (PS) and pH, were used in our statistical analyses.

DNA extraction, PCR, and Illumina MiSeq sequencing. Total DNA was extracted from ~ 10 -g (8.2 to 10.1 g [fresh weight]) soil samples using the PowerMax Soil DNA isolation kit (Mo Bio, Carlsbad, CA), according to the manufacturer's instructions, and stored at -20°C until PCR amplification. The hyper-variable internal transcribed spacer 2 (ITS2) region of the fungal rRNA gene was amplified with primers flITS7 5'-GTGARTCATCGAATCTTTG-3', incorporating a 5'-GTGACTGGAGTTCAGACGTGTGCTCTTCCGA TCT-3' overhang, and ITS4 5'-TCCTCCGTTATTGATATGC-3', incorporating a 5'-ATCTACACTCTTCCCTA CACGACGCTCTCCGATCT-3' overhang. In the secondary PCR, the full-length P5 and indexed P7 Illumina MiSeq adapters were used. The PCRs were performed as described by Koskinen et al. (51). The samples were analyzed using the Fragment Analyzer (Advanced Analytical, USA) and amplicons sequenced with Illumina MiSeq (version 3, 2 × 300-bp paired-end reads) at the Institute of Biotechnology, University of Helsinki.

Bioinformatics. We extracted the ECM fungal data set from a broader environmental sequence data set described in our previous study (17). In the current contribution, we explicitly focused on mycorrhizal communities, because our previous effort on general fungal communities poorly permitted us to address changes in ECM fungal community composition and diversity. Briefly, we processed the paired-end sequence data (.fastq) using mothur version 1.36.1 (52). The fungal.fastq files were contiged, and any sequences with ambiguous bases, with more than one mismatch to the primers, homopolymers longer than 8 bp, and any without a minimum overlap of 50 bp were removed. The sequences were screened for chimeras using UCHIME (53), and putative chimeras were removed. To permit pairwise alignment of fungal ITS sequences to calculate a pairwise distance matrix, we omitted sequences that were shorter than 300 bp and truncated the remaining sequences to the first 300 bp. These fungal sequences were assigned to taxa using the Naive Bayesian Classifier and the UNITE-curated International Nucleotide Sequence Database reference database (54). Any sequences not assigned to the kingdom Fungi were removed. A pairwise distance matrix was derived from pairwise alignments, and sequences clustered to OTUs at a 97% threshold using nearest-neighbor joining. All low-abundance OTUs were removed (≤ 10 sequences across all experimental units), as they may be PCR or sequencing artifacts (55–57). We assigned OTUs into trophic modes using the FUNGuild database (58) and selected ECM fungal OTUs at the cut value of “highly probable.” This resulted in a total of 216,916 sequences representing 357 ECM fungal OTU. We estimated richness and diversity indices for ECM fungal communities in mothur. Observed OTU richness (S_{obs}), the complement of Simpson's diversity ($1/D = 1/\sum p_i^2$), and Simpson's evenness ($E_D = 1/\sum p_i^2/S$), with p_i representing the abundance of each OTU within a sample, were iteratively calculated and subsampled at 517 sequences per sample.

Statistical analyses. All statistical analyses were performed in R (version 3.2.1, R Development Core Team, 2015) using various packages.

ECM fungal community data were analyzed using two different strategies. First, we evaluated differences between urban parks and the control forests (land use type) using a data set including the 10 controls (five control forests with conifer and broadleaf species in the vicinity of the city of Lahti) and old parks (five parks with conifer and broadleaf species within the city of Lahti), for a total of 20 experimental units. We compared controls to old parks because they have trees of virtually the same age class, enabling a comparison between habitat types and excluding tree age. In this analysis, we specifically explored differences in ECM fungal communities in control forests and comparable park treatments. Differences in ECM fungal diversity indices (Ln-transformed where necessary) and counts (sequence abundance) of the dominant genera (the 10 most abundant genera) between land use types were evaluated using generalized linear mixed models (GLMM) with the *lmer* and *glmer* functions in the *lme4* package in R. Diversity index data were modeled following a Gaussian distribution, while count data (the dominant genera) were modeled following a Poisson error distribution, with an individual-level




FIG 5 NMDS plot for ECM fungal communities under conifer trees in parks. The NMDS plot shows significant differences ($P < 0.05$) across conifer tree species by *envfit* analyses.

random effect included to account for possible overdispersion (59). Predictor variables included plant functional type as a factor, land use type as a factor, and their interaction, as well as C and N content of the soil, OM, percent sand (PS), and soil pH. Since our samples were from two different vegetation treatments that may be located in the same park, park location was added as a random term. We performed model selection by removing nonsignificant terms, starting with the term with the highest P value. C, N, and OM content, PS, and pH were initially subject to model simplification until only terms with P values of <0.1 were left. If the land use type \times plant functional group interaction remained nonsignificant ($P > 0.1$) after this procedure, it was also removed. However, to remain true to our experimental design, the main effects (land use type and plant functional type) were always retained in the model irrespective of their significance. Second, we evaluated the effects of plant functional type and park age on ECM fungi. Here, we analyzed a data set including all park age categories and plant functional types, but we omitted the control stands. The response of individual ECM fungal taxa (the 10 most abundant genera only) (count data) and diversity indices (Ln-transformed when necessary) to park age and plant functional type were tested using GLMM. In these analyses, land use was replaced with park age; otherwise, the analyses were identical to those described for the first strategy.

For each of the two discrete analyses, we also utilized nonmetric multidimensional scaling (NMDS; vegan package in R) to visualize community-wide responses to the factors included, based on Bray-Curtis dissimilarity. Soil carbon and nitrogen content, OM content, PS, and soil pH were correlated with the community structure using permutation tests as the vector-fitting procedure (the *envfit* function in vegan). We did the same ordination analyses on the park ECM fungal communities under conifer and broadleaf trees separately. These analyses were motivated by the distinctions between the two plant functional types, thus permitting a more detailed focus on the effects of park age and multiple tree species within the conifer group. In addition, because the same conifer tree species were not consistently present in our parks, we included 5 tree species in this plant functional group and tested the tree species effect on the ECM fungal communities. Although the five conifer tree species differed in their ECM fungal communities ($r^2 = 0.110$, $P = 0.029$; Fig. 5) in a comparison of “inter-tree type variation” with variation between the two plant functional types, the tree species effect within the conifer group was minor.

Accession number(s). The paired fastq files are available in the Sequence Read Archive at NCBI (<https://www.ncbi.nlm.nih.gov/sra>) under accession number [SRX1584451](https://www.ncbi.nlm.nih.gov/sra).

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/AEM.01797-17>.

SUPPLEMENTAL FILE 1, PDF file, 0.4 MB.

ACKNOWLEDGMENTS

We thank Tuukka Ryyänänen and John Allen for help in the field. Satu Tegel, Tuuli Ylikotila, and Pekka Engblom provided data for the parks in Helsinki, and Markku Saari provided data for the parks in Lahti. Lars Paulin provided DNA sequencing services.

The research was funded by an Academy of Finland grant (grant 268548).

REFERENCES

- Gómez-Baggethun E, Barton DN. 2013. Classifying and valuing ecosystem services for urban planning. *Ecol Econ* 86:235–245. <https://doi.org/10.1016/j.ecolecon.2012.08.019>.
- Grote R, Samson R, Alonso R, Amorim JH, Cariñanos P, Churkina G, Fares S, Thiec DL, Niinemets Ü, Mikkelsen TN, Paoletti E, Tiwary A, Calapietra C. 2016. Functional traits of urban trees: air pollution mitigation potential. *Front Ecol Environ* 14:543–550. <https://doi.org/10.1002/fee.1426>.
- Smith S, Read D. 2008. Mineral nutrition, toxic element accumulation and water relations of arbuscular mycorrhizal plants, p 145–148. In Smith SE, Read DJ (ed), *Mycorrhizal symbiosis*, 3rd ed. Academic Press, London, United Kingdom.
- Jonsson LM, Nilsson MC, Wardle DA, Zackrisson O. 2001. Context dependent effects of ectomycorrhizal species richness on tree seedling productivity. *Oikos* 93:353–364. <https://doi.org/10.1034/j.1600-0706.2001.930301.x>.
- Velmala SM, Rajala T, Heinonsalo J, Taylor AF, Pennanen T. 2014. Profiling functions of ectomycorrhizal diversity and root structuring in seedlings of Norway spruce (*Picea abies*) with fast- and slow-growing phenotypes. *New Phytol* 201:610–622. <https://doi.org/10.1111/nph.12542>.
- Luo Z-B, Wu C, Zhang C, Li H, Lipka U, Polle A. 2014. The role of ectomycorrhizas in heavy metal stress tolerance of host plants. *Environ Exp Bot* 108:47–62. <https://doi.org/10.1016/j.envexpbot.2013.10.018>.
- Dighton J, Tuininga AR, Gray DM, Huskins RE, Belton T. 2004. Impacts of atmospheric deposition on New Jersey pine barrens forest soils and communities of ectomycorrhizae. *For Ecol Manage* 201:131–144. <https://doi.org/10.1016/j.foreco.2004.07.038>.
- Bainard LD, Kironomos JN, Gordon AM. 2011. The mycorrhizal status and colonization of 26 tree species growing in urban and rural environments. *Mycorrhiza* 21:91–96. <https://doi.org/10.1007/s00572-010-0314-6>.
- Jumpponen A, Jones KL, David Mattox J, Yaeger C. 2010. Massively parallel 454-sequencing of fungal communities in *Quercus* spp. ectomycorrhizas indicates seasonal dynamics in urban and rural sites. *Mol Ecol* 19:41–53. <https://doi.org/10.1111/j.1365-294X.2009.04483.x>.
- Visser S. 1995. Ectomycorrhizal fungal succession in jack pine stands following wildfire. *New Phytol* 129:389–401. <https://doi.org/10.1111/j.1469-8137.1995.tb04309.x>.
- Kernaghan G, Widden P, Bergeron Y, Légaré S, Paré D. 2003. Biotic and abiotic factors affecting ectomycorrhizal diversity in boreal mixed-woods. *Oikos* 102:497–504. <https://doi.org/10.1034/j.1600-0706.2003.12415.x>.
- Twieg BD, Durall DM, Simard SW. 2007. Ectomycorrhizal fungal succession in mixed temperate forests. *New Phytol* 176:437–447. <https://doi.org/10.1111/j.1469-8137.2007.02173.x>.
- Tedersoo L, Sadam A, Zambrano M, Valencia R, Bahram M. 2010. Low diversity and high host preference of ectomycorrhizal fungi in Western Amazonia, a neotropical biodiversity hotspot. *ISME J* 4:465–471. <https://doi.org/10.1038/ismej.2009.131>.
- Prescott CE, Grayston SJ. 2013. Tree species influence on microbial communities in litter and soil: current knowledge and research needs. *Forest Ecol Manage* 309:19–27. <https://doi.org/10.1016/j.foreco.2013.02.034>.
- Baxter JW, Pickett ST, Carreiro MM, Dighton J. 1999. Ectomycorrhizal diversity and community structure in oak forest stands exposed to contrasting anthropogenic impacts. *Can J Bot* 77:771–782. <https://doi.org/10.1139/b99-039>.
- Timonen S, Kauppinen P. 2008. Mycorrhizal colonisation patterns of *Tilia* trees in street, nursery and forest habitats in southern Finland. *Urban For Urban Gree* 7:265–276. <https://doi.org/10.1016/j.ufug.2008.08.001>.
- Hui N, Jumpponen A, Francini G, Kotze DJ, Liu X, Romantschuk M, Strömmer R, Setälä H. 2017. Soil microbial communities are shaped by vegetation type and park age in cities under cold climate. *Environ Microbiol* 19:1281–1295. <https://doi.org/10.1111/1462-2920.13660>.
- Setälä HM, Francini G, Allen JA, Hui N, Jumpponen A, Kotze DJ. 2016. Vegetation type and age drive changes in soil properties, nitrogen and carbon sequestration in urban parks under cold climate. *Front Ecol Evol* 4:93. <https://doi.org/10.3389/fevo.2016.00093>.
- Sato H, Morimoto S, Hattori T. 2012. A thirty-year survey reveals that ecosystem function of fungi predicts phenology of mushroom fruiting. *PLoS One* 7:e49777. <https://doi.org/10.1371/journal.pone.0049777>.
- Last FT, Dighton J, Mason PA. 1987. Successions of sheathing mycorrhizal fungi. *Trends Ecol Evol* 2:157–161. [https://doi.org/10.1016/0169-5347\(87\)90066-8](https://doi.org/10.1016/0169-5347(87)90066-8).
- Hobbie EA, Macko SA, Shugart HH. 1999. Insights into nitrogen and carbon dynamics of ectomycorrhizal and saprotrophic fungi from isotopic evidence. *Oecologia* 118:353–360. <https://doi.org/10.1007/s004420050736>.
- Schmidt DJE, Pouyat R, Szlavetz K, Setälä H, Kotze DJ, Yesilonis I, Cliiers S, Hornung E, Dombos M, Yarwood SA. 2017. Urbanization erodes ectomycorrhizal fungal diversity and may cause microbial communities to converge. *Nat Ecol Evol* 1:0123. <https://doi.org/10.1038/s41559-017-0123>.
- Amossé J, Dózsa-Farkas K, Boros G, Rochat G, Sandoz G, Fournier B, Mitchell EAD, Le Bayon R-C. 2016. Patterns of earthworm, enchytraeid and nematode diversity and community structure in urban soils of different ages. *Eur J Soil Biol* 73:46–58. <https://doi.org/10.1016/j.ejsobi.2016.01.004>.
- Baxter JW, Dighton J. 2005. Diversity-functioning relationships in ectomycorrhizal fungal communities, p 383–398. In Dighton J, White JF, Jr, Oudemans P (ed), *The fungal community: its organization and role in the ecosystem*. CRC Press, Boca Raton, FL.
- Ochimarou T, Fukuda K. 2007. Changes in fungal communities in evergreen broad-leaved forests across a gradient of urban to rural areas in Japan. *Can J For Res* 37:247–258. <https://doi.org/10.1139/X06-293>.
- Hartmann M, Niklaus PA, Zimmermann S, Schmutz S, Kremer J, Abarenkov K, Lüscher P, Widmer F, Frey B. 2014. Resistance and resilience of the forest soil microbiome to logging-associated compaction. *ISME J* 8:226–244. <https://doi.org/10.1038/ismej.2013.141>.
- Mäkipää R, Rajala T, Schigel D, Rinne KT, Pennanen T, Abrego N, Ovaskainen O. 2017. Interactions between soil- and dead wood-inhabiting fungal communities during the decay of Norway spruce logs. *ISME J* 11:1964–1974. <https://doi.org/10.1038/ismej.2017.57>.
- Wang X-H, Benucci GMN, Xie X-D, Bonito G, Leisola M, Liu P-G, Shamekh S. 2013. Morphological, mycorrhizal and molecular characterization of Finnish truffles belonging to the Tuber anniae species-complex. *Fungal Ecol* 6:269–280. <https://doi.org/10.1016/j.funeco.2013.03.002>.
- Pouyat RV, Yesilonis I, Russell-Anelli J, Neerchal N. 2007. Soil chemical and physical properties that differentiate urban land-use and cover types. *Soil Sci Soc Am J* 71:1010–1019. <https://doi.org/10.2136/sssaj2006.0164>.
- Tedersoo L, Mett M, Ishida TA, Bahram M. 2013. Phylogenetic relationships among host plants explain differences in fungal species richness and community composition in ectomycorrhizal symbiosis. *New Phytol* 199:822–831. <https://doi.org/10.1111/nph.12328>.
- Korkkama T, Pakkanen A, Pennanen T. 2006. Ectomycorrhizal community structure varies among Norway spruce (*Picea abies*) clones. *New Phytol* 171:815–824. <https://doi.org/10.1111/j.1469-8137.2006.01786.x>.
- Hartmann A, Schmid M, Van Tuinen D, Berg G. 2009. Plant-driven selection of microbes. *Plant Soil* 321:235–257. <https://doi.org/10.1007/s11104-008-9814-y>.
- Urbanová M, Šnajdr J, Baldrian P. 2015. Composition of fungal and bacterial communities in forest litter and soil is largely determined by dominant trees. *Soil Biol Biochem* 84:53–64. <https://doi.org/10.1016/j.soilbio.2015.02.011>.
- Tedersoo L, Bahram M, Toots M, Diedhiou AG, Henkel TW, Kjoller R, Morris MH, Nara K, Nouhra E, Peay KG, Polme S, Ryberg M, Smith ME, Koljalg U. 2012. Towards global patterns in the diversity and community

- structure of ectomycorrhizal fungi. *Mol Ecol* 21:4160–4170. <https://doi.org/10.1111/j.1365-294X.2012.05602.x>.
35. Cox F, Barsoum N, Lilleskov EA, Bidartondo MI. 2010. Nitrogen availability is a primary determinant of conifer mycorrhizas across complex environmental gradients. *Ecol Lett* 13:1103–1113. <https://doi.org/10.1111/j.1461-0248.2010.01494.x>.
 36. Högberg MN, Högberg P, Myrold DD. 2007. Is microbial community composition in boreal forest soils determined by pH, C-to-N ratio, the trees, or all three? *Oecologia* 150:590–601.
 37. Avis PG. 2012. Ectomycorrhizal iconoclasts: the ITS rDNA diversity and nitrophilic tendencies of fetid Russula. *Mycologia* 104:998–1007. <https://doi.org/10.3852/11-399>.
 38. Lilleskov EA, Wargo PM, Vogt KA, Vogt DJ. 2008. Mycorrhizal fungal community relationship to root nitrogen concentration over a regional atmospheric nitrogen deposition gradient in the northeastern USA. *Can J Forest Res* 38:1260–1266. <https://doi.org/10.1139/X07-211>.
 39. Lilleskov EA, Hobbie EA, Fahey TJ. 2002. Ectomycorrhizal fungal taxa differing in response to nitrogen deposition also differ in pure culture organic nitrogen use and natural abundance of nitrogen isotopes. *New Phytol* 154:219–231. <https://doi.org/10.1046/j.1469-8137.2002.00367.x>.
 40. Pouyat RV, Russell-Anelli J, Yesilonis ID, Groffman PM. 2003. Soil carbon in urban forest ecosystems. CRC Press, Boca Raton, FL.
 41. Aučina A, Rudawska M, Leski T, Rylíškiš D, Pietras M, Riepišas E. 2011. Ectomycorrhizal fungal communities on seedlings and conspecific trees of *Pinus mugo* grown on the coastal dunes of the Curonian Spit in Lithuania. *Mycorrhiza* 21:237–245. <https://doi.org/10.1007/s00572-010-0341-3>.
 42. Hui N, Jumpponen A, Niskanen T, Liimatainen K, Jones KL, Koivula T, Romantschuk M, Strömmer R. 2011. EcM fungal community structure, but not diversity, altered in a Pb-contaminated shooting range in a boreal coniferous forest site in southern Finland. *FEMS Microbiol Ecol* 76:121–132. <https://doi.org/10.1111/j.1574-6941.2010.01038.x>.
 43. Kranabetter JM, Friesen J, Gamiet S, Kroeger P. 2005. Ectomycorrhizal mushroom distribution by stand age in western hemlock–lodgepole pine forests of northwestern British Columbia. *Can J For Res* 35: 1527–1539. <https://doi.org/10.1139/x05-095>.
 44. Huang J, Nara K, Zong K, Lian C. 2015. Soil propagule banks of ectomycorrhizal fungi along forest development stages after mining. *Microb Ecol* 69:768. <https://doi.org/10.1007/s00248-014-0484-4>.
 45. Richard F, Millot S, Gardes M, Selosse MA. 2005. Diversity and specificity of ectomycorrhizal fungi retrieved from an old-growth Mediterranean forest dominated by *Quercus ilex*. *New Phytol* 166:1011–1023. <https://doi.org/10.1111/j.1469-8137.2005.01382.x>.
 46. Kyaschenko J, Clemmensen KE, Hagenbo A, Karlton E, Lindahl BD. 2017. Shift in fungal communities and associated enzyme activities along an age gradient of managed *Pinus sylvestris* stands. *ISME J* 11:863–874. <https://doi.org/10.1038/ismej.2016.184>.
 47. Coince A, Caël O, Bach C, Lenggellé J, Cruaud C, Gavory F, Morin E, Murat C, Marçais B, Buée M. 2013. Below-ground fine-scale distribution and soil versus fine root detection of fungal and soil oomycete communities in a French beech forest. *Fungal Ecol* 6:223–235. <https://doi.org/10.1016/j.funeco.2013.01.002>.
 48. Lothamer K, Brown SP, Mattox J, Jumpponen A. 2014. Comparison of root-associated communities of native and non-native ectomycorrhizal hosts in an urban landscape. *Mycorrhiza* 24:267–280. <https://doi.org/10.1007/s00572-013-0539-2>.
 49. Baldrian P. 2017. Forest microbiome: diversity, complexity and dynamics. *FEMS Microbiol Rev* 41:109–130.
 50. Setälä H, Francini G, Allen J, Jumpponen A, Hui N, Kotze D. 2017. Urban parks provide ecosystem services by retaining metals and nutrients in soils. *Environ Pollut* 231:451–461. <https://doi.org/10.1016/j.envpol.2017.08.010>.
 51. Koskinen K, Hultman J, Paulin L, Auvinen P, Kankaanpää H. 2011. Spatially differing bacterial communities in water columns of the northern Baltic Sea. *FEMS Microbiol Ecol* 75:99–110. <https://doi.org/10.1111/j.1574-6941.2010.00987.x>.
 52. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber CF. 2009. Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol* 75:7537–7541. <https://doi.org/10.1128/AEM.01541-09>.
 53. Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R. 2011. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27:2194–2200. <https://doi.org/10.1093/bioinformatics/btr381>.
 54. Abarenkov K, Henrik Nilsson R, Larsson KH, Alexander IJ, Eberhardt U, Erland S, Høiland K, Kjølter R, Larsson E, Pennanen T, Sen R, Taylor AFS, Tedersoo L, Ursing BM, Vralstad T, Liimatainen K, Peintner U, Koljalg U. 2010. The UNITE database for molecular identification of fungi—recent updates and future perspectives. *New Phytol* 186:281–285. <https://doi.org/10.1111/j.1469-8137.2009.03160.x>.
 55. Tedersoo L, Nilsson RH, Abarenkov K, Jairus T, Sadam A, Saar I, Bahram M, Bechem E, Chuyong G, Kõljalg U. 2010. 454 pyrosequencing and Sanger sequencing of tropical mycorrhizal fungi provide similar results but reveal substantial methodological biases. *New Phytol* 188:291–301. <https://doi.org/10.1111/j.1469-8137.2010.03373.x>.
 56. Brown SP, Veach AM, Rigdon-Huss AR, Grond K, Lickteig SK, Lothamer K, Oliver AK, Jumpponen A. 2015. Scraping the bottom of the barrel: are rare high throughput sequences artifacts? *Fungal Ecol* 13:221–225. <https://doi.org/10.1016/j.funeco.2014.08.006>.
 57. Oliver AK, Brown SP, Callahan MA, Jr, Jumpponen A. 2015. Polymerase matters: non-proofreading enzymes inflate fungal community richness estimates by up to 15%. *Fungal Ecol* 15:86–89. <https://doi.org/10.1016/j.funeco.2015.03.003>.
 58. Nguyen NH, Song Z, Bates ST, Branco S, Tedersoo L, Menke J, Schilling JS, Kennedy PG. 2016. FUNGuild: an open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecol* 20:241–248. <https://doi.org/10.1016/j.funeco.2015.06.006>.
 59. Harrison XA. 2014. Using observation-level random effects to model overdispersion in count data in ecology and evolution. *PeerJ* 2:e616. <https://doi.org/10.7717/peerj.616>.